

Towards a mechanistic understanding of the drivers of plant-pollinator interactions

Jose B. Lanuza^{1,2}, Alfonso Allen-Perkins³, Henriette F. Morgenroth¹, Raymond Umazekabiri¹, Will Glenny¹, Marc Hoffmann¹, Panagiotis Theodorou^{1,4}, Robert Paxton^{1,4}, Isabell Hensen^{1,4}, Robert Rauschkolb^{1,5}, Christine Römermann^{1,5}, Oliver Schweiger^{1,7}, Martin Freiberg^{1,6}, Tiffany Knight¹

Affiliations

1. German Centre for Integrative Biodiversity Research (iDiv) Halle-JenaLeipzig, Puschstr. 4, 04103 Leipzig, Germany
2. Mediterranean Institute for Advanced Studies (IMEDEA, UIB-CSIC), Mallorca, Balearic Islands, Spain
3. Departamento de Ingeniería Eléctrica, Electrónica, Automática y Física Aplicada, ET-SIDI, Universidad Politécnica de Madrid, 28040 Madrid, Spain.
4. Institute for Biology, Martin-Luther-UniversityHalle-Wittenberg, Hoher Weg 8, 06120 Halle (Saale), Germany
5. Institute of Ecology and Evolution with Herbarium Haussknecht and Botanical Garden, Friedrich Schiller University Jena, Jena, Germany
6. Systematic Botany and Functional Biodiversity/Botanical Garden, Institute of Biology, Leipzig University, Johannisallee 21-23, 04103, Leipzig, Germany
7. UFZ-Helmholtz Centre for Environmental Research, Department of Community Ecology, 06120 Halle, Germany

Abstract

Flowering plants exhibit an astonishing diversity of reproductive structures, which shapes how they interact with their pollinators. However, to what extent reproductive traits or other drivers like phenology and species abundances shape plant-pollinator interactions is still an open question. To disentangle the relative effects of species phenology, abundances and traits in plant-pollinator interactions, we compared pollinator visitation rates, diet breadth, specialization from plant species that vary in reproductive traits, phenology and abundances. To

do this, we observed plant-pollinator interactions for a whole flowering season in three different botanical gardens located in an urban setting. We anticipate that the ecological role of the different plant and pollinator species are largely driven by their phenological overlap and abundance, and only certain taxonomic groups are strongly influenced by trait matching processes. Our results highlight the relevance of considering multiple mechanisms to understand plant-pollinator interactions.

Introduction

1st paragraph

General introductory paragraph

Ecological networks... Introduce drivers End with limitations and state of the art, temporal and spatial components, sampling biases and data limitations from rarity.

2nd paragraph

Highlight the importance of abundance:

There is consensus that including abundances in our view of ecological processes shaping network patterns yields consistently better results [REFS, eg., annual reviews]. However, most ecological studies...

Highlight the importance of phenology: Highlight the importance of traits/phylogeny:

Last paragraph, introduce what we test!

By conducting this experiment in botanical gardens, we have as objective reducing the “noise” associated with the context-dependent nature of ecological interactions [REF].

What use and what temporal resolution of networks are best for improving our predictive understanding of the structure and dynamics of ecological communities? by Caradonna and Waser 2020

Explore how traits, phenology and abundances shape species interactions in three botanical gardens from Germany at different levels of temporal complexity

Pending tasks:

Question 1:

- Explore relative contribution of ecological drivers (abundances, phenology, traits, phylogeny (plants and bees)).
- Associated with this, how increasing sampling impact our view of those processes.

- Temporal grain, how does it affect network patterns. Relative contribution of temporality to network structure or patterns (e.g., day 1, only adds 1 spp versus day 10 that adds 30).

Methods

Study site and sampling method

Plant-pollinator interactions were monitored during the entire flowering season of 2023 in the botanical gardens of Halle, Jena, and Leipzig (Germany). These gardens belong to PhenObs initiative (Nordt et al., 2021), which consists of a network of botanical gardens across the Northern Hemisphere that records phenological observations on a large set of herbaceous species. The objective of PhenObs is to evaluate the impacts of climate change on species performance (e.g., Sporbert et al., 2022). In this study, plant-pollinator interactions were recorded for a subset of species from the PhenObs project within these three gardens. Each garden was visited weekly, and plant-pollinator interactions were sampled during 15 mins on each focal plant. On each sampling day, the different focal plants were sampled in a random order and only pollinator individuals that contacted the reproductive organs of the plant were recorded as interacting species. In addition, to create a comprehensive inventory of pollinators and their interactions within the gardens, 10 random sites were selected each sampling day, and plant-pollinator interactions were monitored for three minutes at each random site, resulting in a total of 30 minutes per sampling day and garden. Therefore, two sampling methodologies were implemented, one that recorded interactions on PhenObs plants, and another that consisted of random stops within the garden, with the specific goal of describing all potential pollinators present in the garden.

Sampling effort and plant taxonomic coverage

This study monitored a total of 78 focal plant species that belonged to PhenObs initiative, which accounted for 68 different genera, 33 families and 22 different orders. In total, we sampled pollinators on 46 plants species in Halle (52.3 hours), 43 in Jena (41.3 hours) and 47 in Leipzig (49.8 hours), and only 12 species were shared across the three gardens. In addition, with the random stops across the garden we sampled a total of XXXX species that comprised XXXX genera, XXXXX families and XXXXX orders. This study accounted for a total of 134 plant species in Halle (10.1 hours), 124 in Jena (11.3 hours), and 125 in Leipzig (10.4 hours). Explain accumulation curves here!!!

Data description

To evaluate the different mechanisms that drive plant-pollinator interactions, detailed information on abundance, phenology and species traits for both plants and pollinators was recorded as follows:

- (i) Floral abundance was annotated for each focal plant species, and given that botanical gardens are a semi-control setting with relatively low abundance of pollinators, we were also able to annotate the different number of pollinator individuals on each sampled plant when documenting interactions. Despite not being fully independent of interactions, this approach offers a more realistic estimate of pollinator abundance than the standard method of estimating abundances from interactions, which can bias our understanding of the mechanisms driving network structure (Blüthgen et al., 2006; Vizentin-Bugoni et al., 2014). Note that flowering species with tiny densely packed flowers were considered at inflorescence level (15.2% of the total flowering species considered in this study).
- (ii) Given that plant phenology was already monitored on the PhenObs project, we have weekly phenological records with flowering percent for all focal plants. For 10 species with missing phenology data, we conducted generative additive models (GAM) with our records of flower number to build the phenology on the same sampling dates as the other species by using the *mgcv* package. Unlike plants, pollinator phenology relies on ‘stochastic’ interaction observations that may not represent their complete flying period. For this reason, we opted to complement our observational data from the Global Biodiversity Information Facility (GBIF). Germany is one of the countries with highest records of occurrence data (Rocha-Ortega et al., 2021), and in particular, their cities (e.g., botanical gardens) contain most of those records (Sweet et al., 2022). Thus, we opted to extract occurrence records around the three cities for the different pollinator species to complement the different pollinator phenologies. We first created a centroid within the 3 cities, and then, created a rectangular shape around it that expanded approximately 1000 kms in longitude and 500 kms in latitude (Figure S1). We used this shape because we expect environmental variables to be more consistent and phenology to be more similar across a longitudinal range than a latitudinal one. Thus, our download resulted in 2,933,772 occurrence records (<https://doi.org/10.15468/dl.k5cnuf>).
- (iii) Plant reproductive traits were selected following the main traits driving the trait spectrum of flowering plants as shown in Lanuza et al. (2023) (i.e., flower number, plant height, style length, flower size, ovule number and selfing). In addition, we also included phenological traits (i.e., flower lifespan and flowering length), floral rewards (nectar volume and number of pollen grains per flower) and seed weight as a measurement of the reproductive investment. Pollinator traits included intertegular distance, body length and proboscis length. Both intertegular distance and body length were measured from images using ImageJ, and proboscis length was extrapolated for bee species by using the pollimetry package (Kendall et al., 2019). Non-bee species were complemented with proboscis values from the literature and images when possible.

Plant and pollinator phylogeny

*Phylogeny... Plant phylogeny is easy, bee phylogeny too...

*compute phenology matrices: presence absence and probability ones based by calculating the probability of observing both if they are independent events divided by total number of sampling days

CONCEPTUAL GRAPH. **Figure 1**

Analysess

To unravel the role of time in our understanding of the cological drivers of plant-pollinator networks, this study analyses the relative importance of abundance, phenology and traits at the whole season level, by dividing in early-mid-and late season and by analysing at the sampling day level.

Results

PROVIDE: Percentage of forbidden links due to phenological mismatches by garden.

References

- Blüthgen, N., Menzel, F., & Blüthgen, N. (2006). Measuring specialization in species interaction networks. *BMC Ecology*, 6, 1–12.
- Kendall, L. K., Rader, R., Gagic, V., Cariveau, D. P., Albrecht, M., Baldock, K. C., Freitas, B. M., Hall, M., Holzschuh, A., Molina, F. P., et al. (2019). Pollinator size and its consequences: Robust estimates of body size in pollinating insects. *Ecology and Evolution*, 9(4), 1702–1714.
- Lanuza, J. B., Rader, R., Stavert, J., Kendall, L. K., Saunders, M. E., & Bartomeus, I. (2023). Covariation among reproductive traits in flowering plants shapes their interactions with pollinators. *Functional Ecology*, 37(7), 2072–2084.
- Nordt, B., Hensen, I., Bucher, S. F., Freiberg, M., Primack, R. B., Stevens, A.-D., Bonn, A., Wirth, C., Jakubka, D., Plos, C., et al. (2021). The PhenObs initiative: A standardised protocol for monitoring phenological responses to climate change using herbaceous plant species in botanical gardens. *Functional Ecology*, 35(4), 821–834.
- Rocha-Ortega, M., Rodriguez, P., & Córdoba-Aguilar, A. (2021). Geographical, temporal and taxonomic biases in insect GBIF data on biodiversity and extinction. *Ecological Entomology*, 46(4), 718–728.

- Sporbert, M., Jakubka, D., Bucher, S. F., Hensen, I., Freiberg, M., Heubach, K., König, A., Nordt, B., Plos, C., Blinova, I., et al. (2022). Functional traits influence patterns in vegetative and reproductive plant phenology—a multi-botanical garden study. *New Phytologist*, 235(6), 2199–2210.
- Sweet, F. S., Apfelbeck, B., Hanusch, M., Garland Monteagudo, C., & Weisser, W. W. (2022). Data from public and governmental databases show that a large proportion of the regional animal species pool occur in cities in germany. *Journal of Urban Ecology*, 8(1), juac002.
- Vizentin-Bugoni, J., Maruyama, P. K., & Sazima, M. (2014). Processes entangling interactions in communities: Forbidden links are more important than abundance in a hummingbird–plant network. *Proceedings of the Royal Society B: Biological Sciences*, 281(1780), 20132397.

Supporting information

Figure S1: Some sort of combination of the code below and the graph of location from the poster