

# Covariation among reproductive traits in flowering plants shapes interactions with floral visitors

Jose B. Lanuza<sup>1,2</sup> , Romina Rader<sup>1</sup> , Jamie Staver<sup>3</sup> , Liam K. Kendall<sup>4</sup> , Manu  
E. Saunders<sup>1</sup>  and Ignasi Bartomeus<sup>2</sup> 

<sup>1</sup> School of Environmental and Rural Science, University of New England, Armidale, New South Wales 2350, Australia. <sup>2</sup> Estación Biológica de Doñana (EBD-CSIC), E-41092 Seville, Spain. <sup>3</sup> Department of Conservation | Te Papa Atawhai, Auckland, New Zealand. <sup>4</sup> Centre for Environmental and Climate Science, Lund University, Sölvegatan 37, S-223 62 Lund, Sweden.

**Keywords:** life-history strategies | plant reproductive traits | plant-pollinator interactions

**Author for correspondence:** Jose B. Lanuza | [barragansljose@gmail.com](mailto:barragansljose@gmail.com)

## Abstract

Globally, plants display enormous variation in life history strategies and trait combinations. However, evidence suggests that evolutionary and physiological constraints limit the number of plant ecological strategies. Although there have been recent advances in understanding correlations among plant traits, reproductive traits are rarely considered, despite their key role in shaping plant life-history strategies and interactions with floral visitors. Here, using a global dataset of 16 reproductive traits for 1,506 species, we investigate the reproductive spectrum of flowering plants to identify how it shapes interactions with floral visitors. We show that over 50% of all trait variation is explained by the first two reproductive axes, which represent the negative correlation between flower number and flower size, and the positive correlation between pollinator dependence and floral display size. In empirical plant-pollinator networks, the plant position within these axes was associated with the number of interacting partners and the visitation rates of the distinct floral visitor guilds, but the unexplained variability found highlights the need to incorporate other factors along with reproductive traits to fully

explain plant-pollinator interactions. Our study identifies the major reproductive trait correlations in flowering plants and their role in shaping networks of plant-pollinator interactions at a global scale.

## 1. Introduction

Flowering plants display an astonishing diversity of reproductive structures [1,2] that shape their interactions with animal pollinators [3,4]. The variety of reproductive strategies used by different plant species, from specialized reliance on a single pollinator, to full self-pollination, is incredibly rich. However, not all reproductive trait combinations in plants are possible due to evolutionary and ecological constraints [5,6]. Despite recent advances in theoretical and empirical understanding of correlations between plant reproductive traits at large ecological scales [7–11], studies typically focus on just a few traits, often in isolation. Thus, to develop a comprehensive understanding of the plant reproductive spectrum of trait variation, as done recently for root [12], leaf [13] and wood [14] related traits, we require a multi-trait perspective with broad geographical coverage. Importantly, characterization of the plant reproductive spectrum could reveal key reproductive strategies among different species [6] and improve knowledge of associations between specific reproductive traits and pollinators [10].

With the recent availability of large trait databases [e.g., TRY 15,COMPADRE 16], there has been increased research on plant ecological strategies, which has produced identification of global patterns in plant form and function [7,17–19]. However, studies that look at multiple traits tend to overlook plant reproductive traits [10,20,21], and focus on trait correlations in other plant organs such as morphological and physiological leaf traits [13,22–24]. Despite the lack of a holistic understanding that depicts reproductive trait covariation patterns, there are widely recognized associations between plant reproductive trait pairs (e.g., the negative correlation between flower size and flower number [25,26]; the positive association between flower size and outcrossing rate [27];

or the positive correlation between outcrossing rate and lifespan [28–30]). Although identification of these trait correlations has allowed progress toward a conceptual framework that integrates different floral trait relationships [10], we still lack empirical understanding of how these associations among floral and other reproductive traits (e.g., reproductive biology) shape plant life history strategies.

Trait-based approaches [3,31] and trait-matching analyses [32,33] are powerful for exploring the underlying mechanisms driving plant-pollinator interactions. For example, plant traits can define species' network roles [e.g., specialists vs generalists 34] and plant species that occupy reproductive trait space extremes are likely to be more specialized and display tighter trait-matching with pollinators [35,36]. Indeed, the degree of morphological matching between plants and floral visitors often determines whether or not interactions occur, and consequently influences the broader network structure [32,37]. However, it is still unclear how specific reproductive traits (e.g., mating or compatibility system) influence plant-pollinator associations [34,38,39]. Indeed, both morphology and mating system can determine a species' functional role within pollination networks, and the combination of several traits increases power for predicting network structure [40]. Because the species position within the multidimensional trait space can result in different interaction patterns or adaptations to pollinators [see 41 for a purely morphological approach], the use of the reproductive trait space to explore plant-pollinator interactions could help to identify species functional roles and associations of traits with specific pollinators (i.e., pollination syndromes).

Here, we explore the spectrum of reproductive traits in flowering plants, within a global set of plant-pollinator networks, to identify reproductive trait covariation and how this influences the network of interactions between plants and floral visitors. First, we identify the major axes of reproductive trait variation and correlations among traits for plant species. Then, we investigate the associations between plant species' positions in reproductive trait space and the strength of their interactions with different floral visitor guilds within networks. Finally, we determine how the main axes of reproductive trait variation influence plant species' functional roles within pollination networks, using a

84 range of complementary interaction network metrics.

## 85 **2. Methods**

### 86 **(a) Plant-pollinator network studies**

87 We selected 28 studies from 18 different countries that constituted a total of 64 plant-  
88 pollinator networks (see table S1 and figure S1). These studies recorded plant-pollinator  
89 interactions in natural systems and were selected so that we had broad geographical  
90 representation across different ecological communities. Although these studies differ  
91 in sampling effort and methodology, all studies provided information about plant-  
92 pollinator interactions (weighted and non-weighted), which we used to build a database  
93 of plant species that are likely to benefit, to some degree, from animal pollination. Many  
94 of these networks are freely available either as published studies [e.g., 42,43,44] or in  
95 online archives [e.g., Mangal 45, The Web of Life 44]. In total, our network dataset  
96 constituted 60 weighted (number of visits) and four unweighted (interaction presence-  
97 absence) networks, each sampled at a unique location and year, as well as eight meta-  
98 networks where interactions were pooled across several locations and multiple years.

### 99 **(b) Taxonomy of plants and pollinators**

100 All species names, genera, families and orders were retrieved and standardized from  
101 the taxonomy data sources NCBI (<https://www.ncbi.nlm.nih.gov/taxonomy>) for  
102 plants and ITIS (<https://www.itis.gov/>) for pollinators, using the R package *taxize* [46].  
103 We filled the 'not found' searches manually using <http://www.theplantlist.org/> and  
104 <http://www.mobot.org/> for plants and <http://www.catalogueoflife.org/> for floral  
105 visitors.

### **(c) Plant traits**

We selected a total of 19 quantitative and categorical functional traits that included reproductive traits and traits related with plant form and size (see table 1). Reproductive traits consisted of those related to floral and reproductive biology: (i) floral traits included those related to the size of floral organs (e.g., style length), floral display size (e.g., flower number) and floral rewards (e.g., pollen quantity), while (ii) reproductive biology traits represented the plant reproductive system (i.e., the breeding, mating and compatibility system). The remaining traits were related to plant size and form (i.e., plant height, lifespan and life form), which are commonly used to characterize the fast-slow continuum of plant trait variation [7]. For each plant species, we undertook an extensive literature and online search for all traits across a wide range of resources (plant databases, online floras, books, journals and images). From a matrix comprising a total of 28,614 possible cells (considering all traits and plant species: 19 columns  $\times$  1,506 species), we were able to fill 23,354 cells (81.6% of the dataset, see figure S2 for missing value information for each trait). An extended description of each trait and how it was obtained can be found in Appendix S1.

### **(d) Phylogenetic Distance**

We calculated the phylogenetic distance between all plant species considered using the function *get\_tree* from the package *rtrees* (<https://github.com/daijiang/rtrees>), which downloads phylogenetic distances from the extended R implementation of the Open Tree of Life [47,48].

### **(e) Data Imputation**

To avoid excluding species with trait missing values, we imputed trait missing values with the help of the function *missForest* [49] which allows imputation of datasets with continuous and categorical variables. We accounted for the phylogenetic distance

**Table 1.** Quantitative and categorical traits used in this study.

<b>Traits</b>	<b>Numerical range/Categories</b>
<b>Quantitative</b>	
Plant height	0.01-30m
Flower width	0.50-205mm
Flower length	0.20-195mm
Inflorescence width	0.80-300mm
Style length	0-138mm
Ovules per flower	1-25000
Flowers per plant	1-1.8 × 10 <sup>5</sup>
Microliters of nectar	0-160μl
Milligrams of nectar	0-6.1mg
Nectar concentration	0-100%
Pollen grains per flower	13-2 × 10 <sup>8</sup>
Autonomous selfing (fruit set)	0-100%
<b>Categorical</b>	
Lifepan	Short-lived and perennial
Life form	Herb, shrub and tree
Flower shape	Brush, campanulate, capitulum, open, papilionaceous and tube
Flower symmetry	Actinomorphic and zygomorphic
Nectar	Presence and absence
Compatibility system	Self-incompatible, partially self-compabtile and self-compatible
Breeding system	Hermaphrodite, monoecious and dioecious

among species on the imputation process by including the eigenvectors from principal component analysis of the phylogenetic distance (PCoA), which has been shown to improve the performance of *missForest* [50]. To extract the eigenvectors, we used the function *PVRdecomp* from the package *PVR* [51] based on a previous conceptual framework that considers phylogenetic eigenvectors [52]. We conducted two different imputations, one for the full set of species (1,506 species: 5.79% of missing values), and excluded nectar and pollen traits because of the high percentage of missing values (figure S2), and a second one for the subset of species with data for pollen per flower and microliters of nectar (755 species: 8.01% of missing values). To corroborate that our imputation of missing values did not affect our results, we evaluated the reproductive spectrum (see section below) with and without imputed values and results for the two datasets were consistent (Fig S3 and Fig S4).

#### (f) Plant strategies

We explored the association between the different quantitative plant traits using a phylogenetically informed Principal Component Analysis (pPCA). We did not include the quantitative variables of flower length and inflorescence width because they were highly and moderately correlated to flower width, respectively (Pearson's correlation = 0.72,  $P < 0.01$  and Pearson's correlation = 0.36,  $P < 0.01$ ), and thus we avoided overemphasizing flower size on the spectrum of trait variation. Prior to the analyses, we excluded outliers and standardized the data. Specifically, due to the high sensitivity of dimensionality reduction to outliers [53,54], we excluded values outside the 2.5th–97.5th percentile range, and thus our final dataset had 1,236 species. Then, we log transformed all continuous variables to reduce the influence of extreme values and then z-transformed ( $X = 0$ ,  $SD = 1$ ) these variables so that they were within the same numerical range, as recommended for principal component analysis [54]. Although qualitative traits were not included in the dimensionality reduction analysis, we also investigated the statistical association of the different qualitative traits with the main axes of trait variation using an ANOVA and Tukey test. We performed the pPCA using

the function *phyl.pca* from the package *phytools* [55] with the method lambda ( $\lambda$ ) that calculates the phylogenetic correlation between 0 (phylogenetic independence) and 1 (shared evolutionary history) and we implemented the mode “covariance” because values for each variables were on the same scale following transformation [56]. Finally, we conducted an additional phylogenetic informed principal component analysis for the subset of species with pollen and nectar quantity data. For this, we included all quantitative traits included in the main pPCA, plus pollen grains and microlitres of nectar per flower.

### **(g) Phylogenetic signal**

We calculated the phylogenetic signal of the different quantitative traits on the imputed dataset with the full set of species ( $N = 1,506$ ) with the package *phytools* [55] and used Pagel’s  $\lambda$  as a measurement of the phylogenetic signal. However, for pollen and nectar traits, phylogenetic signal was calculated only on the subset of species that had quantitative information for these traits ( $N = 755$ ).

### **(h) Network analyses**

#### **(i) Visitation patterns**

We used Bayesian modeling (see details below) to investigate how the different guilds of floral visitors interacted along the main axes of reproductive trait variation using qualitative and quantitative measures of plant-pollinator interactions. Specifically, interactions between plants and floral visitors in binary networks were defined as the presence or absence of floral visitation, while in weighted networks, interactions were measured as the the number of visits from each insect species to different plant species. Although floral visitors are not always pollinators and the number of visits does not consider the pollination efficiency of each species [57], the number of visits does provide valuable information about the relative contributions different floral visitors make to pollination [58,59]. We divided floral visitors into six main taxonomical



guilds that differ in life form, behaviour and thus likely play similar ecological roles: (i) bees (Hymenoptera-Anthophila), (ii) non-bee-Hymenoptera (Hymenoptera-non-Anthophila), (iii) syrphids (Diptera-Syrphidae), (iv) non-syrphid-Diptera (Diptera-non-Syrphidae), (v) Lepidoptera, and (vi) Coleoptera. Moreover, because bees had the greatest number of occurrences in networks (2,256 records) and the highest frequency of visits, we further broke this group down into the main bee families (Andrenidae, Apidae, Colletidae, Halictidae, and Megachilidae). In addition, we found that *Apis mellifera* was the floral visitor with the largest proportion of occurrences in networks (7.55% of the total). This is consistent with previous research showing that *A. mellifera* was the most frequent floral visitor in a similar dataset of 80 plant-pollinator networks in natural ecosystems [60]. Hence, to control for the effect of *A. mellifera* on observed visitation patterns of bees, we conducted an analogous analysis with presence-absence of the interaction and number of visits excluding *A. mellifera*. We found that *A. mellifera* was partly driving some of the observed trends on PC1 (figure S5). However, we did not detect major differences on PC2 and PC3.

We used Bayesian generalized linear mixed models in the R package *brms* [61] to model the presence-absence of observed interactions and number of visits as a function of the main axes of plant trait variation and their interactions with floral visitor guilds (e.g., number of visits  $\sim$  PC1  $\times$  FGs + PC2  $\times$  FGs + PC3  $\times$  FGs). Because we were interested in possible differences in visitation patterns among floral visitor guilds to plants with different reproductive strategies, we included interactions between the main axes of trait variation (PC1, PC2 and PC3) and the different floral visitor guilds in the model. We added a nested random effect of networks within the study system to capture the variation in networks among studies and within networks. Moreover, we included the phylogenetic covariance matrix as a random factor due to the possible shared evolutionary histories of species and therefore lack of independence among them. We specified for the presence or absence of an interaction and number of visits with Bernoulli and a zero inflated negative binomial distributions, respectively. The models were run with 3,000 iterations with previous 1,000 warm up iterations and

with non or very weakly informative priors from the `brm` function so the priors had negligible influence on the results [61]. We set delta ( $\Delta$ ) to 0.99 to avoid divergent transitions and visualized the posterior predictive checks with the function `pp_check` using the *bayesplot* package [62].

## (ii) Plant species network roles

We also used Bayesian modeling to investigate how the main axes of trait variation determined different plant species' functional roles within pollination networks. For this, we selected complementary species-level network metrics, commonly applied in bipartite network studies [63], with clear ecological interpretations and are relevant for defining species' functional roles in networks. Specifically, the species-level metrics we included were: (i) sum of visits per plant species; (ii) normalized degree, calculated as the number of links per plant species divided by the total possible number of partners; and (iii) specialization ( $d'$ ) [64], which measures the deviation of an expected random choice of the available interaction partners and ranges between 0 (maximum generalization) and 1 (maximum specialization). Normalized degree and specialization were calculated with the *specieslevel* function in the R package *bipartite* [63]. We modeled each plant species metric (sum of visits, normalized degree and plant specialization) as a function of the three main axes of trait variation (plant species level metric  $\sim$  PC1 + PC2 + PC3). For the different response variables in each model (i.e., each plant species level metric), we used different distribution families (zero inflated negative binomial for the sum of visits, weibull for normalized degree, and zero-one inflated beta for specialization). Finally, we used the same random factors, model settings and conducted the same posterior predictive checks for each model as detailed above in the 'visitation patterns' section.

Our analyses were conducted on the subset of 60 weighted networks (out of a total of 64 networks) sampled in a unique flowering season and site, which included 556 plant and 1,126 pollinator species. In total, our network dataset (excluding meta- and non-weighted networks) included 2,256 interactions for bees, 1,768 for non-syrphid-


dipterans, 845 for syrphids, 437 for lepidopterans, 432 for coleopterans, and 362 for non-bee-hymenopterans. Sampling methods varied across studies but this was accounted for in analyses by including study identity as a random effect in our models. All analyses were conducted in R version 4.0.3.

### 3. Results

#### (a) Plant strategies

Our phylogenetically informed principal component analysis captured 51.8% and 70.97% of the reproductive trait variation in the first two and three axes, respectively (figure 1 and figure S6). The first principal component (PC1; the flower number - flower size axis) represented 26.72% of the trait variation and indicated a negative correlation between flower number and flower size. The main contributing traits to PC1 were plant height, flower number, ovule number and flower size (loadings  $> |0.5|$ ; table S3) but style length also contributed moderately (loading = -0.33). One end of this axis comprised species with high investment in flower number and plant height but small flower size, short style length and low ovule number. For instance, *Cornus florida* occurred at this end of the spectrum, which, on average, has approximately 10,000 flowers, is 7.5 m high, has flowers that are 3 mm wide, styles 3.5 mm long and two ovules per flower. The other end of this spectrum had species that were short in height, with few large flowers that had long styles and many ovules. For example, *Petunia axillaris* on average, is 0.5 m high and has approximately 10 flowers per plant that are over 50 mm wide with 25 mm long styles and over 200 ovules. The second principal component (PC2; the pollinator dependence axis) encompassed 25.05% of the trait variation and represented variation from low to high autonomous selfing (i.e., high to low pollinator dependence). The main driver of trait variation on PC2 was autonomous selfing (loading = 0.85) but all other traits (except ovule number) made moderate contributions (loadings ranging between 0.27 to 0.4; table S3). Generally speaking, species with high pollinator dependence had larger and many flowers with long styles,


and were tall. *Zuccagnia punctata* occurred at this extreme of the spectrum, which is a self-incompatible species, completely dependent on animal pollination for seed production, producing approximately 1,500 flowers per plant with an average height of 3 m and 20 mm long styles. In contrast, species with low pollinator dependence tended to have fewer and smaller flowers with shorter styles, and were shorter in height. *Veronica peregrina* occurred at this end of the spectrum, and is a self-compatible plant that requires little or no pollination by animals, it has approximately 20 flowers per plant, is 0.2 m high and has 0.25 mm long styles. The third principal component (PC3; the style length - pollinator dependence axis) explained a considerable amount of trait variability (19.17%) and represented a negative correlation between style length (loading = -0.66) and pollinator dependence (autonomous selfing loading =  $|-0.51|$ ), where species with short styles had high pollinator dependence and species with long styles low pollinator dependence. The remaining traits, apart from ovule number, were all positively correlated with style length and negatively correlated with pollinator dependence (loadings from -0.23 to -0.46; table S3). In addition, the pPCA with the subset of species that we had nectar and pollen quantity data for showed that nectar quantity (microlitres of nectar per flower) and pollen grains per flower were positively associated with floral display size. Floral rewards were particularly strongly correlated with flower size but weakly correlated with flower number (figure S4). The first two principal components for this pPCA explained similar variance (45.52%) and similar associations of traits, despite some variability in the loadings (table S4).



**Figure 1.** Phylogenetically informed principal component analysis (pPCA) of 1,236 plant species from 28 plant-pollinator network studies. The solid arrows indicate the direction of the different quantitative traits (flower number, plant height, style length, flower size, ovule number and level of autonomous selfing) across the two main axes of trait variation. The length of the arrows indicate the weight of the variables on each principal component and the dashed lines show the opposed direction of trait variation. The icons at both ends of arrows and dashed lines illustrate the extreme form of the trait continuum.

We found that most categorical traits were statistically associated with the first two axes of trait variation (figure 2 and table S2). Flower symmetry, which was only associated with PC2 (Sum of squares = 8.51, F-value = 14.72,  $P < 0.01$ ), and nectar provision, which was independent of PC1 and PC2 (PC1: Sum of squares = 0.37, F-value = 0.29,  $P = 0.59$ ; PC2: Sum of squares = 0.83, F-value = 1.43,  $P = 0.23$ ) showed a lack of statistical association. In addition, we found statistical differences between the different categorical trait levels within the trait space (figure S7). Regarding self-compatibility,

we found larger differences on PC2. For instance, species with unisexual flowers that were self-incompatible, were statistically differentiated from species with partial or full self-compatibility on this axis (figure S7a and figure S7b; Tukey test  $P < 0.05$ ). Life forms differed statistically across both axes of trait variation and followed a gradient of larger life forms (trees and shrubs) with higher pollinator dependence to smaller ones (herbs) with lower pollinator dependence (figure S7c and figure S7d; Tukey test  $P < 0.05$ ). Consequently, lifespan also followed this gradient but perennial and short lived species only differed statistically on PC2 (figure S7e and figure S7f; Tukey test  $P < 0.05$ ). Species with unisexual flowers (monoecious and dioecious) were clustered on both extremes of the first two principal components and had both the highest pollinator dependence and highest number of flowers (figure S7g and figure S7h; Tukey test  $P < 0.05$ ). Moreover, we found that campanulate and capitulum flower shapes were differentiated from tube, papilionaceous, open and brush shapes in the trait space; the former had larger flowers and high pollinator dependence, while the latter had more flowers and lower pollinator dependence (figure S7i and figure S7j; Tukey test  $P < 0.05$ ). Finally, in terms of flower symmetry, zygomorphic flowers were associated with lower pollinator dependence, whereas actinomorphic flowers were more dependent on pollinators (figure S7k and figure S7l; Tukey test  $P < 0.05$ ).



**Figure 2.** Location of the different qualitative traits in the trait space. The figure panel is composed of the traits that showed statistical association with the first two axes of trait variation: compatibility system (a), life form (b), lifespan (c), breeding system (d), flower shape (e) and flower symmetry (f). The solid arrows indicate the direction of variation of the different quantitative traits shown in figure 1: flower number (FN), plant height (PH), style length (SL), flower size (FS), ovule number (ON) and the level of autonomous selfing (S).

## **(b) Phylogenetic signal**

We found a strong phylogenetic signal ( $P < 0.01$ ) for most quantitative traits (table S5). The traits that showed the highest phylogenetic signal were ovule number ( $\lambda = 1$ ), pollen grains per flower ( $\lambda = 1$ ) and plant height ( $\lambda = 0.96$ ), followed by flower length ( $\lambda = 0.75$ ), flower width ( $\lambda = 0.73$ ), number of flowers per plant ( $\lambda = 0.69$ ) and nectar concentration ( $\lambda = 0.65$ ). The traits that had a moderate phylogenetic signal included inflorescence width ( $\lambda = 0.57$ ), style length ( $\lambda = 0.49$ ) and autonomous selfing ( $\lambda = 0.34$ ). Finally, microliters of nectar per flower showed the lowest phylogenetic signal of all the traits we measured ( $\lambda = 0.14$ ).


## **(c) Visitation patterns**

The main axes of trait variation (PC1, PC2 and PC3) partly explained the presence-absence of interaction partners (conditional  $R^2 = 0.26$ ; marginal  $R^2 = 0.20$ ) but little of the total number of visits (conditional  $R^2 = 0.31$ ; marginal  $R^2 = 0.06$ ). However, we found relevant differences across the different floral visitor guilds for both the presence-absence of interactions and number of visits (figure 3). On the flower number - flower size axis, we found that plants with many small flowers tended to have interaction partners that were Coleoptera, non-bee-Hymenoptera and Diptera (figure 3a), whereas plants with few large flowers mostly had interaction partners that were bees and Lepidoptera. We found similar trends for visitation rates on the flower number - flower size axis, but bees and syrphids visited plant species with many small flowers more frequently (figure 3d). On the pollinator dependence axis, we found that all plant species with higher pollinator dependence had many interaction partners and received many visits from all floral visitor guilds (figure 3b and figure 3e). Finally, on the style length - pollinator dependence axis, plant species with short styles and high pollinator dependence had more interaction partners for all guilds, except bees which interacted more frequently with plant species with long styles and low pollinator dependence (figure 3c). Plants with long styles and low pollinator dependence interacted more



342 frequently with Lepidoptera and non-bee-Hymenoptera (figure 3f).

343 The additional models for the most represented bee families, for both presence-absence  
344 of interactions (marginal  $R^2 = 0.29$ ; conditional  $R^2 = 0.19$ ) and the number of visits  
345 (marginal  $R^2 = 0.30$ ; conditional  $R^2 = 0.03$ ), showed that the Apidae family was the main  
346 driver of the observed patterns. The contrast between interaction presence/absence  
347 and number of visits for bees on PC1 (figure 3a and figure 3d) was driven by the  
348 Andrenidae family, which had more interaction partners but made fewer visits to plant  
349 species with few large flowers (figure S8).



**Figure 3.** Fitted posterior estimates of the presence-absence of interaction (a, b and c) and number of visits (d, e and f) of the different floral visitor guilds in relation to the main axes of plant reproductive trait variation (PC1, PC2 and PC3). PC1 represents the flower number - flower size axis, PC2 represents the pollinator dependence axis and PC3 represents the style length - pollinator dependence axis. For visualization purposes due to large differences between number of visits from bees and the rest of the pollinator guilds, the number of visits was log-transformed (Y-axis of lower panel).

#### (d) Plant species network roles

Variance within plant species-level network metrics was poorly explained by the three main axes of trait variation (figure S9; number of visits ~ PCs, conditional  $R^2 = 0.11$ , marginal  $R^2 = 0.02$ ; normalized degree ~ PCs, conditional  $R^2 = 0.24$ , marginal  $R^2 = 0.02$ ; and, specialization ~ PCs, conditional  $R^2 = 0.37$ , marginal  $R^2 = 0.03$ ). Overall, the most notable trends were found on the flower number - flower size axis (PC1) and the style length - pollinator dependence axis (PC3) for number of visits and specialization. On PC1, the number of visits per flower was higher for plant species with more flowers but was lower for plant species with larger flowers (figure S9a). However, specialization (d') was higher for plant species with large flowers but lower for plant species with small flowers (figure S9g). On PC3, the number of visits was lower for plants with shorter styles and higher pollinator dependence but higher for plant species with longer styles and lower pollinator dependence (figure S9c). Again, specialization showed the opposite trend to the number of visits (figure S9i).

## 4. Discussion

We found strong covariation among traits within the flowering plant reproductive spectrum, which shapes interactions with floral visitors at a global scale. Despite the enormous variability in plant reproductive structures, the first two and three axes captured over 50% and 70% of trait variation, respectively, and these axes encapsulated correlations among flower number and flower size, pollinator dependence and floral display, and style length and pollinator dependence. Although explained trait variation on the first two axes was less than previous studies of morphological and physiological traits [17,19], our results align with the largest and most recent study of plant life strategies that included reproductive, form and size related traits [7]. The different plant reproductive axes that we identified were associated with different numbers of interacting partners and visitation rates of the distinct floral visitor guilds. However, the unexplained variability found highlights the need to account for other local factors,

such as species relative abundances [33], along with reproductive traits, to better explain plant-pollinator associations.

Our study provides strong evidence for several widely discussed trait associations with the help of a great diversity of phylogenetic lineages (170 families and over 1,200 species). Specifically, by using a multitrait approach, we are able to link the reproductive spectrum of plant trait variation with the previously described “fast-slow continuum” in plant [7] and animal [65] life-history strategies. This fast-slow continuum was illustrated by the associations between reproductive traits and plant height, life form and lifespan traits. For instance, although the negative correlation between flower number and flower size has received substantial attention [25,26,66], it has only been investigated across a few hundred Angiosperm species in the absence of a multitrait perspective. We found support for this trait association and show that other reproductive traits (e.g., style length or ovule number) covary along this flower number - flower size axis. Our study also complements previous work linking plant breeding systems and life forms [8,67] by showing that most species with unisexual flowers also have woody life forms and many flowers. Moreover, we identify the previously documented positive association between outcrossing rate and floral display size [27], where species with high pollinator dependence also allocate large resource quantities to attract floral visitors. Although we did not include floral rewards in our main analysis due to insufficient data, we did find that plant species with large floral displays and, in particular, large flowers, tended to have higher pollen and nectar quantity. Thus, we demonstrate that globally, reproductive trait covariation in flowering plants likely imposes constraints on the broader spectrum of plant trait variation.

Overall, plant reproductive traits explained interaction patterns among and within floral visitor guilds suggesting that plant reproductive strategies partly mediate these associations across spatial large scales. For example, we found that all floral visitor guilds interacted more frequently with large flowered and pollinator dependent plant species, which aligns with previous findings [68–70]. As predicted by evolutionary life-history theory [71], larger investment in floral display and rewards to attract

pollinators occurs when pollinator dependence is high. However, the possible range of reproductive strategies in flowering plants is constrained by the cost of reproductive organs, trait associations and selective pressures exerted by different floral visitor guilds [72]. For instance, smaller flowers are visited less overall, but they are inexpensive to produce (e.g., have a low ovule number) and low visitation rate at the individual flower level can be compensated by overall visitation at the plant level by producing many flowers resulting in a larger quantity of small fruits [26]. In addition, because floral traits can be associated with their animal pollinators to optimise pollen transfer, the correlation between distinct suites of reproductive traits and floral visitor guilds could indicate the existence of pollination syndromes [3,4,31,73]. For instance, bees had a higher number of interacting partners on plant species with large flowers but interacted more frequently with species with small flowers while syrphid flies showed the opposite trend. This could be reflective of differences in the ecology and behaviour between these pollinator guilds [74,75] that ultimately exert different selective pressures, leading divergent plant reproductive strategies. In any case, it is worth noting that local factors (e.g., species relative abundances or phenologies) likely contribute to the variability in the association between the plant reproductive spectrum and floral visitors [33,76,77].

Our study provides a comprehensive assessment of reproductive trait covariation patterns in flowering plants, using a global dataset of traits for plant species within plant-pollinator networks. We show the importance of floral visitors in mediating reproductive trait correlations and identify the need to consider broader plant reproductive strategies, rather than just flower morphology, to understand patterns of plant-pollinator interactions. Further, we provide empirical evidence of reproductive trait associations that complements trait covariation studies for other plant organs, highlighting the need to include reproductive traits in the global spectrum of plant form and function. Future work incorporating underrepresented areas of the world and additional relevant reproductive traits (e.g., flower life span and colour) will be valuable for enhancing our understanding of the plant reproductive spectrum and

435 associated plant-pollinator interactions.

## 436 References

- 437 1. Barrett SCH. 2002 The evolution of plant sexual diversity. *Nat. Rev. Genet.* **3**,  
438 274–284. (doi:[10.1038/nrg776](https://doi.org/10.1038/nrg776))
- 439 2. Schiestl FP, Johnson SD. 2013 Pollinator-mediated evolution of floral signals.  
440 *Trends Ecol. Evol.* **28**, 307–315. (doi:[10.1016/j.tree.2013.01.019](https://doi.org/10.1016/j.tree.2013.01.019))
- 441 3. Fenster CB, Armbruster WS, Wilson P, Dudash MR, Thomson JD. 2004 Polli-  
442 nation Syndromes and Floral Specialization. *Annu. Rev. Ecol. Evol. Syst.* **35**,  
375–403. (doi:[10.1146/annurev.ecolsys.34.011802.132347](https://doi.org/10.1146/annurev.ecolsys.34.011802.132347))
- 443 4. Dellinger AS. 2020 Pollination syndromes in the 21st century: Where do we stand  
444 and where may we go? *New Phytol.* **228**, 1193–1213. (doi:[10.1111/nph.16793](https://doi.org/10.1111/nph.16793))
- 445 5. Stearns SC. 1989 Trade-offs in life-history evolution. *Funct. Ecol.* **3**, 259–268.  
446 (doi:[10.2307/2389364](https://doi.org/10.2307/2389364))
- 447 6. Agrawal AA. 2020 A scale-dependent framework for trade-offs, syndromes, and  
448 specialization in organismal biology. *Ecology* **101**, e02924. (doi:[10.1002/ecy.2924](https://doi.org/10.1002/ecy.2924))
- 449 7. Salguero-Gómez R, Jones OR, Jongejans E, Blomberg SP, Hodgson DJ, Mbeau-  
Ache C, Zuidema PA, de Kroon H, Buckley YM. 2016 Fast-slow continuum and  
reproductive strategies structure plant life-history variation worldwide. *Proc.*  
450 *Natl. Acad. Sci. U.S.A.* **113**, 230–235. (doi:[10.1073/pnas.1506215112](https://doi.org/10.1073/pnas.1506215112))
- 451 8. Friedman J. 2020 The evolution of annual and perennial plant life histories:  
Ecological correlates and genetic mechanisms. *Annu. Rev. Ecol. Evol. Syst.* **51**,  
452 461–481. (doi:[10.1146/annurev-ecolsys-110218-024638](https://doi.org/10.1146/annurev-ecolsys-110218-024638))
- 453 9. Paterno GB, Silveira CL, Kollmann J, Westoby M, Fonseca CR. 2020 The male-  
ness of larger angiosperm flowers. *Proc. Natl. Acad. Sci.* **117**, 10921–10926.  
454 (doi:[10.1073/pnas.1910631117](https://doi.org/10.1073/pnas.1910631117))

- 455 10. Roddy AB, Martínez-Perez C, Teixido AL, Cornelissen TG, Olson ME, Oliveira  
RS, Silveira FAO. 2021 Towards the flower economics spectrum. *New Phytol.* **229**,  
456 665–672. (doi:[10.1111/nph.16823](https://doi.org/10.1111/nph.16823))
- 457 11. Song B *et al.* 2022 Global analysis of floral longevity reveals latitudinal gradients  
458 and biotic and abiotic correlates. *New Phytologist* (doi:[10.1111/nph.18271](https://doi.org/10.1111/nph.18271))
- 459 12. Laughlin DC *et al.* 2021 Root traits explain plant species distributions along  
climatic gradients yet challenge the nature of ecological trade-offs. *Nat. Ecol.*  
460 *Evol.*, 1–12. (doi:[10.1038/s41559-021-01471-7](https://doi.org/10.1038/s41559-021-01471-7))
- 461 13. Wright IJ *et al.* 2004 The worldwide leaf economics spectrum. *Nature* **428**, 821–827.  
462 (doi:[10.1038/nature02403](https://doi.org/10.1038/nature02403))
- 463 14. Chave J, Coomes D, Jansen S, Lewis SL, Swenson NG, Zanne AE. 2009 To-  
wards a worldwide wood economics spectrum. *Ecol. Lett.* **12**, 351–366.  
464 (doi:[10.1111/j.1461-0248.2009.01285.x](https://doi.org/10.1111/j.1461-0248.2009.01285.x))
- 465 15. Kattge J *et al.* 2011 TRY – a global database of plant traits. *Glob. Chang. Biol.* **17**,  
466 2905–2935. (doi:[10.1111/j.1365-2486.2011.02451.x](https://doi.org/10.1111/j.1365-2486.2011.02451.x))
- 467 16. Salguero-Gómez R *et al.* 2015 The compadre Plant Matrix Database: An open  
online repository for plant demography. *J. Ecol.* **103**, 202–218. (doi:[10.1111/1365-2745.12334](https://doi.org/10.1111/1365-2745.12334))  
468
- 469 17. Díaz S *et al.* 2016 The global spectrum of plant form and function. *Nature* **529**,  
470 167–171. (doi:[10.1038/nature16489](https://doi.org/10.1038/nature16489))
- 471 18. Bruelheide H *et al.* 2018 Global trait–environment relationships of plant commu-  
472 nities. *Nat. Ecol. Evol.* **2**, 1906–1917. (doi:[10.1038/s41559-018-0699-8](https://doi.org/10.1038/s41559-018-0699-8))
- 473 19. Carmona CP *et al.* 2021 Erosion of global functional diversity across the tree of  
474 life. *Sci. Adv.* **7**, eabf2675. (doi:[10.1126/sciadv.abf2675](https://doi.org/10.1126/sciadv.abf2675))
- 475 20. E-Vojtkó A, Bello F de, Durka W, Kühn I, Götzenberger L. 2020 The neglected  
importance of floral traits in trait-based plant community assembly. *J. Veg. Sci.*  
476 **31**, 529–539. (doi:[10.1111/jvs.12877](https://doi.org/10.1111/jvs.12877))

21. Rüger N, Comita LS, Condit R, Purves D, Rosenbaum B, Visser MD, Wright SJ, Wirth C. 2018 Beyond the fast-slow continuum: Demographic dimensions structuring a tropical tree community. *Ecol. Lett.* **21**, 1075–1084. (doi:[10.1111/ele.12974](https://doi.org/10.1111/ele.12974))
22. Shipley B, Lechowicz MJ, Wright I, Reich PB. 2006 Fundamental trade-offs generating the worldwide leaf economics spectrum. *Ecology* **87**, 535–541. (doi:[10.1890/05-1051](https://doi.org/10.1890/05-1051))
23. Donovan LA, Maherali H, Caruso CM, Huber H, Kroon H de. 2011 The evolution of the worldwide leaf economics spectrum. *Trends in Ecology & Evolution* **26**, 88–95. (doi:[10.1016/j.tree.2010.11.011](https://doi.org/10.1016/j.tree.2010.11.011))
24. Osnas JL, Lichstein JW, Reich PB, Pacala SW. 2013 Global leaf trait relationships: Mass, area, and the leaf economics spectrum. *Science* **340**, 741–744. (doi:[10.1126/science.1231574](https://doi.org/10.1126/science.1231574))
25. Sargent RD, Goodwillie C, Kalisz S, Ree RH. 2007 Phylogenetic evidence for a flower size and number trade-off. *Am. J. Bot.* **94**, 2059–2062. (doi:[10.3732/ajb.94.12.2059](https://doi.org/10.3732/ajb.94.12.2059))
26. Kettle CJ, Maycock CR, Ghazoul J, Hollingsworth PM, Khoo E, Sukri RSH, Burslem DFRP. 2011 Ecological Implications of a Flower Size/Number Trade-Off in Tropical Forest Trees. *PLoS One* **6**, e16111. (doi:[10.1371/journal.pone.0016111](https://doi.org/10.1371/journal.pone.0016111))
27. Goodwillie C *et al.* 2010 Correlated evolution of mating system and floral display traits in flowering plants and its implications for the distribution of mating system variation. *New Phytol.* **185**, 311–321. (doi:[10.1111/j.1469-8137.2009.03043.x](https://doi.org/10.1111/j.1469-8137.2009.03043.x))
28. Barrett SCH. 2003 Mating strategies in flowering plants: The outcrossing-selfing paradigm and beyond. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **358**, 991–1004. (doi:[10.1098/rstb.2003.1301](https://doi.org/10.1098/rstb.2003.1301))
29. Munoz F, Violle C, Cheptou P-O. 2016 CSR ecological strategies and plant mating systems: Outcrossing increases with competitiveness but stress-tolerance is related to mixed mating. *Oikos* **125**, 1296–1303. (doi:[10.1111/oik.02328](https://doi.org/10.1111/oik.02328))



- 495 30. Moeller DA *et al.* 2017 Global biogeography of mating system variation in seed  
496 plants. *Ecol. Lett.* **20**, 375–384. (doi:[10.1111/ele.12738](https://doi.org/10.1111/ele.12738))
- 497 31. Rosas-Guerrero V, Aguilar R, Martén-Rodríguez S, Ashworth L, Lopezaraiza-  
Mikel M, Bastida JM, Quesada M. 2014 A quantitative review of pollination  
syndromes: Do floral traits predict effective pollinators? *Ecol. Lett.* **17**, 388–400.  
498 (doi:[10.1111/ele.12224](https://doi.org/10.1111/ele.12224))
- 499 32. Stang M, Klinkhamer PGL, Waser NM, Stang I, van der Meijden E. 2009 Size-  
specific interaction patterns and size matching in a plant–pollinator interaction  
500 web. *Ann. Bot.* **103**, 1459–1469. (doi:[10.1093/aob/mcp027](https://doi.org/10.1093/aob/mcp027))
- 501 33. Bartomeus I, Gravel D, Tylianakis JM, Aizen MA, Dickie IA, Bernard-Verdier M.  
2016 A common framework for identifying linkage rules across different types  
502 of interactions. *Funct. Ecol.* **30**, 1894–1903. (doi:[10.1111/1365-2435.12666](https://doi.org/10.1111/1365-2435.12666))
- 503 34. Tur C, Castro-Urgal R, Traveset A. 2013 Linking Plant Specialization to Depen-  
dence in Interactions for Seed Set in Pollination Networks. *PLoS One* **8**, e78294.  
504 (doi:[10.1371/journal.pone.0078294](https://doi.org/10.1371/journal.pone.0078294))
- 505 35. Junker RR, Blüthgen N, Brehm T, Binkenstein J, Paulus J, Schaefer HM, Stang  
M. 2013 Specialization on traits as basis for the niche-breadth of flower visitors  
and as structuring mechanism of ecological networks. *Funct. Ecol.* **27**, 329–341.  
506 (doi:[10.1111/1365-2435.12005](https://doi.org/10.1111/1365-2435.12005))
- 507 36. Coux C, Rader R, Bartomeus I, Tylianakis JM. 2016 Linking species functional  
roles to their network roles. *Ecol. Lett.* **19**, 762–770. (doi:[10.1111/ele.12612](https://doi.org/10.1111/ele.12612))
- 509 37. Ibanez S. 2012 Optimizing size thresholds in a plant-pollinator interaction web:  
Towards a mechanistic understanding of ecological networks. *Oecologia* **170**,  
510 233–242. (doi:[10.1007/s00442-012-2290-3](https://doi.org/10.1007/s00442-012-2290-3))
- 511 38. Sargent RD, Ackerly DD. 2008 Plant–pollinator interactions and the assembly of  
512 plant communities. *Trends Ecol. Evol.* **23**, 123–130. (doi:[10.1016/j.tree.2007.11.003](https://doi.org/10.1016/j.tree.2007.11.003))

39. Devaux C, Lepers C, Porcher E. 2014 Constraints imposed by pollinator behaviour on the ecology and evolution of plant mating systems. *J. Evol. Biol.* **27**, 1413–1430. (doi:[10.1111/jeb.12380](https://doi.org/10.1111/jeb.12380))
40. Eklöf A *et al.* 2013 The dimensionality of ecological networks. *Ecol. Lett.* **16**, 577–583. (doi:[10.1111/ele.12081](https://doi.org/10.1111/ele.12081))
41. Dehling DM, Jordano P, Schaefer HM, Böhning-Gaese K, Schleuning M. 2016 Morphology predicts species' functional roles and their degree of specialization in plant–frugivore interactions. *Proc. Royal Soc. B* **283**, 20152444. (doi:[10.1098/rspb.2015.2444](https://doi.org/10.1098/rspb.2015.2444))
42. Carvalheiro LG *et al.* 2014 The potential for indirect effects between co-flowering plants via shared pollinators depends on resource abundance, accessibility and relatedness. *Ecol. Lett.* **17**, 1389–1399. (doi:[10.1111/ele.12342](https://doi.org/10.1111/ele.12342))
43. Olesen JM, Bascompte J, Dupont YL, Jordano P. 2007 The modularity of pollination networks. *PNAS* **104**, 19891–19896. (doi:[10.1073/pnas.0706375104](https://doi.org/10.1073/pnas.0706375104))
44. Fortuna MA, Stouffer DB, Olesen JM, Jordano P, Mouillot D, Krasnov BR, Poulin R, Bascompte J. 2010 Nestedness versus modularity in ecological networks: Two sides of the same coin? *J. Anim. Ecol.* **79**, 811–817. (doi:[10.1111/j.1365-2656.2010.01688.x](https://doi.org/10.1111/j.1365-2656.2010.01688.x))
45. Poisot T *et al.* 2016 Mangal – making ecological network analysis simple. *Ecography* **39**, 384–390. (doi:[10.1111/ecog.00976](https://doi.org/10.1111/ecog.00976))
46. Chamberlain S *et al.* 2020 Taxize: Taxonomic information from around the web. R package version 0.9.99. At <https://CRAN.r-project.org/package=taxize>.
47. Smith SA, Brown JW. 2018 Constructing a broadly inclusive seed plant phylogeny. *Am. J. Bot.* **105**, 302–314. (doi:[10.1002/ajb2.1019](https://doi.org/10.1002/ajb2.1019))
48. Jin Y, Qian H. 2019 V.PhyloMaker: An R package that can generate very large phylogenies for vascular plants. *Ecography* **42**, 1353–1359. (doi:[10.1111/ecog.04434](https://doi.org/10.1111/ecog.04434))

49. Stekhoven DJ, Bühlmann P. 2012 MissForest—non-parametric missing value imputation for mixed-type data. *Bioinformatics* **28**, 112–118. (doi:[10.1093/bioinformatics/btr597](https://doi.org/10.1093/bioinformatics/btr597))
50. Penone C, Davidson AD, Shoemaker KT, Marco MD, Rondinini C, Brooks TM, Young BE, Graham CH, Costa GC. 2014 Imputation of missing data in life-history trait datasets: Which approach performs the best? *Methods Ecol. Evol.* **5**, 961–970. (doi:[10.1111/2041-210X.12232](https://doi.org/10.1111/2041-210X.12232))
51. Chamberlain S *et al.* 2018 Package “PVR”. R package version 0.3. At <https://CRAN.r-project.org/package=PVR>.
52. Diniz-Filho JAF, Bini LM, Rangel TF, Morales-Castilla I, Olalla-Tárraga MÁ, Rodríguez MÁ, Hawkins BA. 2012 On the selection of phylogenetic eigenvectors for ecological analyses. *Ecography* **35**, 239–249. (doi:[10.1111/j.1600-0587.2011.06949.x](https://doi.org/10.1111/j.1600-0587.2011.06949.x))
53. Serneels S, Verdonck T. 2008 Principal component analysis for data containing outliers and missing elements. *Comput. Stat. Data Anal.* **52**, 1712–1727. (doi:[10.1016/j.csda.2007.05.024](https://doi.org/10.1016/j.csda.2007.05.024))
54. Legendre P, Legendre L. 2012 *Numerical ecology*. 2nd edn. Elsevier, Amsterdam.
55. Revell LJ. 2012 Phytools: An R package for phylogenetic comparative biology (and other things). *Methods Ecol. Evol.* **3**, 217–223. (doi:[10.1111/j.2041-210X.2011.00169.x](https://doi.org/10.1111/j.2041-210X.2011.00169.x))
56. Abdi H, Williams LJ. 2010 Principal component analysis. *WIREs Comp. Stats.* **2**, 433–459. (doi:[10.1002/wics.101](https://doi.org/10.1002/wics.101))
57. Ballantyne G, Baldock KCR, Willmer PG. 2015 Constructing more informative plant–pollinator networks: Visitation and pollen deposition networks in a heathland plant community. *Proc. Royal Soc. B* **282**, 20151130. (doi:[10.1098/rspb.2015.1130](https://doi.org/10.1098/rspb.2015.1130))

58. Vázquez DP, Morris WF, Jordano P. 2005 Interaction frequency as a surrogate for the total effect of animal mutualists on plants. *Ecol. Lett.* **8**, 1088–1094. (doi:[10.1111/j.1461-0248.2005.00810.x](https://doi.org/10.1111/j.1461-0248.2005.00810.x))
59. Vázquez DP, Lomáscolo SB, Maldonado MB, Chacoff NP, Dorado J, Stevani EL, Vitale NL. 2012 The strength of plant–pollinator interactions. *Ecology* **93**, 719–725. (doi:[10.1890/11-1356.1](https://doi.org/10.1890/11-1356.1))
60. Hung K-LJ, Kingston JM, Albrecht M, Holway DA, Kohn JR. 2018 The worldwide importance of honey bees as pollinators in natural habitats. *Proc. Royal Soc. B* **285**, 20172140. (doi:[10.1098/rspb.2017.2140](https://doi.org/10.1098/rspb.2017.2140))
61. Bürkner P-C. 2017 Brms: An R package for Bayesian multilevel models using Stan. *J. Stat. Softw.* **80**, 1–28. (doi:[10.18637/jss.v080.i01](https://doi.org/10.18637/jss.v080.i01))
62. Gabry J, Simpson D, Vehtari A, Betancourt M, Gelman A. 2019 Visualization in Bayesian workflow. *J. R. Stat. Soc. Ser. A Stat. Soc.* **182**, 389–402. (doi:[10.1111/rssa.12378](https://doi.org/10.1111/rssa.12378))
63. Dormann CF, Gruber B, Fründ J. 2008 Introducing the bipartite package: Analysing ecological networks. *R News* **8/2**.
64. Blüthgen N, Menzel F, Blüthgen N. 2006 Measuring specialization in species interaction networks. *BMC Ecol.* **6**, 9. (doi:[10.1186/1472-6785-6-9](https://doi.org/10.1186/1472-6785-6-9))
65. Healy K, Ezard THG, Jones OR, Salguero-Gómez R, Buckley YM. 2019 Animal life history is shaped by the pace of life and the distribution of age-specific mortality and reproduction. *Nat. Ecol. Evol.* **3**, 1217–1224. (doi:[10.1038/s41559-019-0938-7](https://doi.org/10.1038/s41559-019-0938-7))
66. Worley AC, Baker AM, Thompson JD, Barrett SC. 2000 Floral display in narcissus: Variation in flower size and number at the species, population, and individual levels. *Int. J. Plant Sci.* **161**, 69–79. (doi:[10.1086/314225](https://doi.org/10.1086/314225))
67. Petit RJ, Hampe A. 2006 Some evolutionary consequences of being a tree. *Annu. Rev. Ecol. Evol. Syst.* **37**, 187–214. (doi:[10.1146/annurev.ecolsys.37.091305.110215](https://doi.org/10.1146/annurev.ecolsys.37.091305.110215))

- 571 68. Hegland SJ, Totland Ø. 2005 Relationships between species' floral traits  
and pollinator visitation in a temperate grassland. *Oecologia* **145**, 586–594.  
572 (doi:[10.1007/s00442-005-0165-6](https://doi.org/10.1007/s00442-005-0165-6))
- 573 69. Lázaro A, Jakobsson A, Totland Ø. 2013 How do pollinator visitation rate and  
seed set relate to species' floral traits and community context? *Oecologia* **173**,  
574 881–893. (doi:[10.1007/s00442-013-2652-5](https://doi.org/10.1007/s00442-013-2652-5))
- 575 70. Kaiser-Bunbury CN, Vázquez DP, Stang M, Ghazoul J. 2014 Determinants  
of the microstructure of plant–pollinator networks. *Ecology* **95**, 3314–3324.  
576 (doi:[10.1890/14-0024.1](https://doi.org/10.1890/14-0024.1))
- 577 71. Stebbins GL. 1970 Adaptive radiation of reproductive characteristics in an-  
giosperms, i: Pollination mechanisms. *Annual Review of Ecology and Systematics*,  
578 307–326. (doi:[10.1146/annurev.es.01.110170.001515](https://doi.org/10.1146/annurev.es.01.110170.001515))
- 579 72. Galen C. 1999 Why do flowers vary? The functional ecology of variation in  
flower size and form within natural plant populations. *Bioscience* **49**, 631–640.  
580 (doi:[10.2307/1313439](https://doi.org/10.2307/1313439))
- 581 73. Faegri K, Van Der Pijl L. 2013 *Principles of pollination ecology*. 3rd edn. Elsevier.  
582
- 583 74. Rader R, Cunningham S, Howlett B, Inouye D. 2020 Non-bee insects as visitors  
and pollinators of crops: Biology, ecology, and management. *Annu. Rev. Entomol.*  
584 **65**, 391–407. (doi:[10.1146/annurev-ento-011019-025055](https://doi.org/10.1146/annurev-ento-011019-025055))
- 585 75. Földesi R, Howlett BG, Grass I, Batáry P. 2021 Larger pollinators deposit more  
pollen on stigmas across multiple plant species—a meta-analysis. *J. Appl. Ecol.*  
586 **58**, 699–707. (doi:[10.1111/1365-2664.13798](https://doi.org/10.1111/1365-2664.13798))
- 587 76. Vázquez DP, Melián CJ, Williams NM, Blüthgen N, Krasnov BR, Poulin R. 2007  
Species abundance and asymmetric interaction strength in ecological networks.  
588 *Oikos* **116**, 1120–1127. (doi:[10.1111/j.0030-1299.2007.15828.x](https://doi.org/10.1111/j.0030-1299.2007.15828.x))

77. Encinas-Viso F, Revilla TA, Etienne RS. 2012 Phenology drives mutualistic network structure and diversity. *Ecol. Lett.* **15**, 198–208. (doi:[10.1111/j.1461-0248.2011.01726.x](https://doi.org/10.1111/j.1461-0248.2011.01726.x))

**Data accessibility.** All data and code used to conduct this study are available in the online repositories of Zenodo (<https://doi.org/10.5281/zenodo.6705621>) and Github (<https://github.com/JoseBSL/Reproductive-traits>).

**Authors' contributions.** JBL, RR and IB designed the study. JBL collated the data and conducted analysis with guidance of JS, LKK and IB. JBL wrote the manuscript with contributions of all authors.

**Conflict of interest declaration.** We declare we have no competing interests.

**Funding.** This study was supported by the European project SAFEGUARD (101003476 H2020-SFS-2019-2) and JBL was supported by a University of New England IPRA postgraduate scholarship.

**Acknowledgements.** We thank all researchers that made their data openly available and sent it upon request. We also thank Bryony Wilcox, Greg Bible, Mercedes Sanchez-Lanuza and David Ragel for their help with data collection. We thank Jason Tylianakis for his insightful comments on the manuscript before submission. Finally, we thank Marcos Méndez, Susan Kalisz, Amanda Benoit, May Berenbaum and an anonymous reviewer for thoughtful feedback and useful comments on earlier versions of this manuscript.