

Trade-offs among plant reproductive traits determine interactions with floral visitors

Jose B. Lanuza^{1,2*}, Romina Rader¹, Jamie Stavert³, Liam K. Kendall⁴,
Manu E. Saunders¹ and Ignasi Bartomeus²

Plant life strategies are constrained by cost-benefit trade-offs that determine plant form and function. Despite the recent advances in the understanding of plant trade-offs with vegetative and physiological traits, little is known about the plant reproductive economics and how they constrain plant life strategies and shape interactions with floral visitors. Here, we investigated plant reproductive trade-offs and their association with floral visitors with a dataset of 1506 plant species with 17 reproductive and 3 vegetative traits from 28 plant-pollinator network studies across 18 countries. Then, we tested whether a species reproductive strategy predicted plant-pollinator interactions (presence-absence and visitation rate) and if the interaction strategy (its role in a pollination network) was predicted by individual traits. We found that 51.8% of trait variation was explained by two independent axes that encompassed plant form and function. Specifically, the first axis indicated the presence of a trade-off between flower number and flower size (PC1, 26.72%) and the second axis indicated a trade-off for the level of pollinator dependency (PC2, 25.08%). The reproductive trade-offs explained some of the qualitative interaction with pollinators (presence-absence) but none of the variability of visitation rates. However, noteworthy differences across pollinator guilds were found in both cases. Finally, we found that specific traits with special relevance of floral rewards were key in the understanding of the species network role. Our results highlight the need to consider plant reproductive trade-offs to improve our understanding of plant life strategies and plant-pollinator interactions at broader spatial scales.

¹ School of Environmental and Rural Science, University of New England, Armidale, New South Wales 2350, Australia. ² Estación Biológica de Doñana (EBD-CSIC), E-41092 Seville, Spain. ³ Department of Conservation | Te Papa Atawhai, Auckland, New Zealand. ⁴ Centre for Environmental and Climate Science, Lund University, Sölvegatan 37, S-223 62 Lund, Sweden. * e-mail: barragansljose@gmail.com

Despite the astonishing diversity of floral structures among flowering plants^{1,2} and their relevance to understand plant-pollinator interactions^{3,4}, a unified framework that explores plant reproductive compromises is currently lacking⁵. In addition, macroecological studies regarding specific reproductive traits are scarce⁶⁻⁹ and as a consequence studies that link reproductive traits to plant-pollinator interactions at large

scales are poorly explored^{10–13}. Linking the position in the trait-space with the different pollinator groups could help to improve our understanding of plant-pollinator associations¹⁴. At the same time, there is increasing interest in the understanding of the determinants of plant-pollinator interactions via trait-based approaches^{3,15} and trait-matching analyses^{16,17}. However, reproductive traits have been overlooked beyond highly specialised systems⁴ despite the apparent generalist nature of plant-pollinator interactions^{18,19}. In addition, it is yet unclear how the different reproductive biology traits influence plant-pollinator associations^{20,21}.

Species can optimize their fitness through various life-history traits, yet trade-offs among those traits constrain the range of potential strategies that a species can employ. With the recent availability of large trait databases (e.g., TRY²² and COMPADRE²³), plant ecological strategies are increasingly being examined, and are facilitating the identification of global patterns and constraints of plant form and function^{12,24,25}. However, the main focus has been on vegetative traits such as leaf²⁶, wood²⁷, or root²⁸ trade-offs with little or no attention given to reproductive traits^{5,29}, also critical to plant life strategies that shape plant interactions with pollinators and ultimately impact plant reproductive success. For instance, short lived versus perennial species tend to have low versus high levels of outcrossing, respectively^{9,30}. Further, outcrossing levels are positively correlated with flower size³¹. In addition, the presence of costly rewards (e.g., pollen or nectar) and showy flowers or floral displays can only be understood through consideration of plant species' reliance upon animal pollination (pollinator dependence) and their role in attracting pollinators³². However, it is still unknown to what extent these different reproductive compromises determine plant-pollinator associations.

Several studies have identified links between plant traits and plant-pollinator network properties^{33–35}. Moreover, plant traits can also define species' network roles^{20,36} (e.g., specialists vs generalists). For example, species that occupy trait space extremes are more likely to exhibit higher specialization and be more reliant on trait-matching^{37,38}.

This morphological matching between plant and floral visitors can determine plant-pollinator interactions, and thus shape their interaction network structure^{16,39}. Remarkably, the combination of traits have shown to increase the predictive power of the network interactions⁴⁰. Therefore, considering the different plant reproductive trade-offs which represent the species strategy within the network¹⁴ could progress our understanding of plant-pollinator interactions. Further, we know little if the different network patterns generally studied at the community level are maintained when we scale up at macroecological scale.

Here, we explore the potential trade-offs among plant floral and reproductive biology traits and how these influence plant-pollinator interactions. First, we identify the major axes of reproductive trait variation and trade-offs that determine plant form and function. Second, we investigate how plant species' position in trait-space influences plant-pollinator interactions (presence-absence and visitation rate). Finally, we investigate how the main axes of trait variation and individual traits influence plant species roles within networks using complementary interaction network metrics (i.e., interaction strength, normalized degree and specialization).

RESULTS

Plant strategies. The phylogenetically informed principal component analysis (pPCA) captured by the first two and three axes 51.8% and 70.97% of trait variation, respectively (Fig. 1 and Supplementary Fig. S5) and had a phylogenetic correlation (λ) of 0.76. The first principal component (PC1) represented 26.72% of the trait variation and indicated a trade-off between flower number and flower size. We refer to this axis as the 'flower number - flower size trade-off', as already described in previous studies^{41,42}. Hence, one end of the spectrum comprised species with high investment in flower number and plant height but small flower size, short style length and low ovule number. The other end of this spectrum comprised species that were short in height and invested in large

92 flowers, long styles, many ovules, but few flowers. The main contributing traits to PC1
 93 were plant height, flower number, ovule number and flower size (loadings $> |0.5|$;
 94 Supplementary Table S3) but style length also contributed moderately on PC1 (loading
 95 $= -0.33$). The second principal component (PC2) represented 25.05% of the trait variation
 96 and indicated a trade-off between low and high pollinator dependence. We refer to
 97 this axis as the 'pollinator dependence trade-off'. The main driver of trait variation on
 98 PC2 was autonomous selfing (loading $= 0.85$) but the other traits (except ovule number)
 99 also made moderate contributions (loadings from 0.27 to 0.4; Supplementary Table S3).
 100 We found that high pollinator dependence was associated with larger and a higher
 101 number of flowers, greater plant height and longer styles. In contrast, species with high
 102 levels of autonomous selfing tended to have fewer and smaller flowers, had shorter
 103 styles and were shorter in height. Further, PC3 explained a considerable amount of trait
 104 variability (19.17%) and the main contributors to this axis were style length (loading
 105 $= -0.66$) and the degree of autonomous selfing (loading $= -0.51$). The remaining traits,
 106 apart from ovule number, were moderately correlated to changes on PC3 (loadings
 107 from -0.23 to -0.46; Supplementary Table S3). Thus, because style length was correlated
 108 with all traits on PC3 and was the main driver of trait variation, we refer to this axis
 109 as the 'style length trade-off'. Further, the pPCA with the subset of species that had
 110 nectar and pollen quantity data showed that nectar quantity (microlitres of nectar per
 111 flower) was positively associated with flower size, style length and ovule number (PC1,
 112 23.40%); and pollen quantity (pollen grains per flower) was positively correlated with
 113 flower number and plant height and negatively associated with autonomous selfing
 114 (PC2, 21.67%; Supplementary Fig. S6). This pPCA explained similar variance with the
 115 first two principal components (45.07%) and similar associations of traits despite some
 116 variability in the loadings (Supplementary Table S4).



Fig. 1 | Plant life-history strategies. Phylogenetically informed principal component analysis (pPCA) of 1,236 plant species from 28 plant-pollinator network studies. The solid arrows indicate the direction of the different quantitative traits (flower number, plant height, style length, flower size, ovule number and level of autonomous selfing) across the two main axes of trait variation. The length of the arrows indicate the weight of the variables on each principal component and the dashed lines show the opposed direction of trait variation. The icons at both ends of arrows and dashed lines illustrate the extreme form of the trait continuum.

117 We found that most categorical traits were statistically associated with the first two axes
 118 of trait variation (Fig. 2 and Supplementary Table S2). Flower symmetry, which was
 119 only associated with PC2 (Sum of squares = 8.51, F-value = 14.72, $P < 0.01$), and nectar
 120 provision, which was independent of PC1 and PC2 (PC1: Sum of squares = 0.37, F-value

121 = 0.29 , $P = 0.59$; PC2: Sum of squares = 0.83, F-value = 1.43, $P = 0.23$) showed lack of
 122 statistical association. In addition, we found (with a Tukey test) statistical differences
 123 between the different levels of categorical traits in the trait space (Supplementary Fig.
 124 S7). Regarding self compatibility, we found larger differences on PC2 (i.e., species
 125 with unisexual flowers that were self incompatible were statistically differentiated from
 126 species with partial or full self compatibility; Supplementary Fig. S7a and Fig. S7b). Life
 127 forms differed statistically across both axes of trait variation and followed a gradient
 128 of larger life forms (trees and shrubs) with higher pollinator dependence to smaller
 129 ones (herbs) with lower pollinator dependence (Supplementary Fig. S7c and Fig. S7d).
 130 Consequently, lifespan also followed this gradient but perennial and short lived species
 131 only differed statistically on PC2 (Supplementary Fig. S7e and Fig. S7f). Species with
 132 unisexual flowers (monoecious and dioecious) were clustered on both extremes of
 133 the first two principal components and had the highest pollinator dependence and
 134 highest number of flowers (Supplementary Fig. S7g and Fig. S7h). Moreover, we
 135 found that the campanulate and capitulum flower shapes were differentiated from tube,
 136 papilionaceous, open and brush shapes in the trait space. The former morphologies
 137 had larger flowers and greater pollinator dependence, while the latter had higher
 138 flower number and greater autonomous selfing (Supplementary Fig. S7i and Fig. S7j).
 139 Regarding flower symmetry, zygomorphic flowers were associated with lower levels of
 140 pollinator dependence, whereas actinomorphic flowers had higher levels of pollinator
 141 dependence (Supplementary Fig. S7k and Fig. S7l).




Fig. 2 | Location of the different qualitative traits on the trait space. The panel is composed by the traits that showed statistical association with the first two axes of trait variation: compatibility system (a), life form (b), lifespan (c), breeding system (d), flower shape (e) and flower symmetry (f).

Phylogenetic signal of traits. We found a strong phylogenetic signal ($P < 0.01$) in all quantitative traits (Supplementary Table S5). The traits that showed the highest phylogenetic signal were ovule number ($\lambda = 1$), pollen grains per flower ($\lambda = 1$) and plant height ($\lambda = 0.96$), followed by flower length ($\lambda = 0.75$), flower width ($\lambda = 0.73$), number of flowers per plant ($\lambda = 0.69$) and nectar concentration ($\lambda = 0.65$). The traits that showed a moderate phylogenetic signal were inflorescence width ($\lambda = 0.57$), style length ($\lambda = 0.49$) and autonomous selfing ($\lambda = 0.34$). Finally, microliters of nectar per flower showed the lowest phylogenetic signal of all traits ($\lambda = 0.14$).

Visitation patterns. The main axes of trait variation explained partly presence-absence interactions between plant and floral visitors (*conditional* $R^2 = 0.26$; *marginal* $R^2 = 0.20$) but little of the overall visitation rates (*conditional* $R^2 = 0.31$; *marginal* $R^2 = 0.06$). However, we found relevant trends across the different floral visitor guilds on both presence-absence and visitation interactions (Fig. 3). On the pollinator dependence trade-off, all floral visitor guilds interacted more frequently with plant species with higher pollinator dependence (PC2; Fig. 3b and Fig. 3e). For presence-absence interactions we found that all Diptera, Coleoptera and non-bee Hymenoptera guilds interacted more frequently with plants with high flower number and small flowers (flower number - flower size trade-off, PC1; Fig. 3a) but bees and Lepidoptera interacted slightly more frequently with plant species with low flower number but large flowers. For presence-absence interactions on PC3 (style length trade-off; Fig. 3c), we found that bees interacted clearly more with plant species with long styles and high selfing and the rest of the guilds interacted slightly more with plant species with short styles and low selfing. In addition, all guilds other than Syrphids and Lepidoptera (i.e., all Hymenoptera, non-Syrphidae-Diptera and Coleoptera) showed greater visitation rates on species with small numerous flowers (PC1; Fig. 3d). On the style length trade-off, bees, Lepidoptera and Non-Anthophila-Hymenoptera showed greater visitation rates on plant species with larger styles and higher levels of selfing; while Syrphidae, Non-Syrphidae-Diptera and Coleoptera showed higher visitation rates on species with shorter styles and lower selfing (Fig. 3f).

171 The additional model for both presence-absence (*marginalR2* = 0.29; *conditionalR2* =
172 0.19) and visitation rate (*marginalR2* = 0.30; *conditionalR2* = 0.03) for the most rep-
173 resented families of bees showed that the family Apidae was the main driver of the
174 observed patterns and that the contrasting differences between presence-absence and
175 visitation rate for bees on PC1 (Fig. 3a and Fig. 3d) were driven by the family An-
176 drenidae which interacted more frequently on presence-absence interactions with plant
177 species with low number of flower but large (Supplementary Fig. S8).




Fig. 3 | Interaction (yes/no) and visitation rates across the three main axes of trait variation per floral visitor guild. Fitted posterior estimates of the presence/absence of interaction (a, b and c) and number of visits (d, e and f) made by the different floral visitors guilds in relation to PC1, PC2 and PC3. PC1 represents the flower number - flower size trade-off, PC2 represents the pollinator dependence trade-off and PC3, the style length trade-off. For visualization purposes, due to large differences between the visitation rates of bees and the rest of guilds, the number of visits was log-transformed (Y-axis of lower panel).

Plant species functional roles. The variance of the different plant species-level network metrics was poorly explained by the three main axes of trait variation (Supplementary Fig. S9; interaction frequency ~ PCs, *conditionalR2* = 0.11, *marginalR2* = 0.02; normalized degree ~ PCs, *conditionalR2* = 0.24, *marginalR2* = 0.02; and, specialization ~ PCs, *conditionalR2* = 0.37, *marginalR2* = 0.03). Overall, the most notable trends were found on PC1 and PC3 for interaction frequency and specialization. On the flower number - flower size trade-off (PC1), interaction frequency was higher for plant species with more flowers but was lower for plant species with larger flowers. On PC1, specialization showed the opposite trend. On the style length trade-off (PC3), interaction frequency was lower for plants with shorter styles and lower autonomous selfing and higher for species with longer styles and higher autonomous selfing. Again, specialization showed the opposite trend to interaction frequency.

When we further investigate which combination of traits drive plant network roles, we show that the regression tree for visitation frequency was best explained by plant height, nectar concentration and style length (Fig. 4a). Specifically, species taller than 3.9 m had the highest interaction frequency, while species that were shorter than 3.9 m and had a nectar concentration lower than 16% had the lowest interaction frequency. Normalized degree was best explained by nectar concentration, pollen grains per flower, plant height, flower width and autonomous selfing (Fig. 4b). Species with a nectar concentration over 49% had the highest levels of normalized degree, whereas species with nectar concentration lower than 49%, more than 21,000 pollen grains per flower and height less than 0.78 m had the lowest normalized degree. Finally, specialization was best explained by plant height, ovule number, pollen grains per flower and autonomous selfing (Fig. 4c). Overall, plant species with the highest specialization were shorter than 1.3 m, had more than 14,000 pollen grains per flower and autonomously self-pollinated less than 11% of their fruits. In contrast, species taller or equal than 5.1 m and with lower than 14 ovules per flower had the lowest specialization values.




Fig. 4 | Contribution of traits in plant's network roles. Regression tree analysis of interaction frequency (log-transformed), normalized degree and specialization for the subset of species with quantitative data for pollen and nectar traits. The superior value inside the node indicates the mean value of the different species-level metric and the lower value, the percentage of species that are considered in each node. Thus, the top node has the mean value of the named trait for the 100% of species. Each node has a yes/no question and when the condition is fulfilled, the branch turns to the 'yes' direction and when not, to the 'no' direction. This rationale is followed in all the regression trees as indicated in the first branch division of the topmost node of each tree.

DISCUSSION

Here, we show that plant species exhibit clear trade-offs in their floral, reproductive biology and vegetative traits. These trade-offs are differentiated on three main axes of trait variation: (i) flower number - flower size, (ii) pollinator dependence and (iii) style length. These axes of trait variation explained partly the presence-absence interactions between plants and floral visitors but did not explain the overall visitation rates. However, on both qualitative and quantitative interactions floral visitor guilds showed associations with the different reproductive trade-offs. Interestingly, pollen and nectar related traits were better than all other traits for characterizing plant species functional roles within their interaction networks.

Over 50% of plant trait variation was captured by the flower number - flower size and pollinator dependence trade-offs. Variation in these two axes of trait variation appears to be associated with the commonly described 'fast-slow continuum' in plant¹² and animal⁴³ life-history strategies as indicated by the different floral and reproductive biology traits associated with plant height, life form and lifespan. The 'slow' part of the continuum (i.e., tall trees and shrubs) had plant species with many flowers, few ovules, higher pollinator dependence, greater occurrence of self-incompatibility and more complex breeding systems (e.g., monoecious and dioecious species). In contrast, species that employed the 'fast' strategy (i.e., short herbs), had fewer flowers, more ovules, greater occurrence of self compatibility and lower pollinator dependence. Further, on the first two axes of trait variation, we found additional support for the positive association already described in a previous study between outcrossing rate and floral display³¹. This trend was further confirmed when looking at the specific correlation between floral display and autonomous selfing for our dataset (see Supplementary Fig. S10).

Despite the low predictive power of the main axes of trait variation on overall interac-

tion patterns (presence-absence and visitation rate), we found commonalities among and within floral visitors guilds across these axes that indicate the influence of life-history strategies on plant-pollinator interactions. For example, all guilds showed an increasing trend of visitation on plant species with higher pollinator dependence, which were also associated with larger floral displays and greater pollen quantities (Fig. 1 and Supplementary Fig. S6). This trend is consistent with previous plant-pollinator studies that show higher visitation rates on species that make greater reproductive investment^{36,44,45}. In regard to the flower number - flower size and style length trade-offs, different guilds showed contrasting visitation rates, which could be associated with different pollination syndromes at a macroecological scale. For instance, the guilds of bees and syrphids were clearly associated with opposing life-strategies on PC1 and PC3 (Fig. 3) indicating a possible niche partitioning among these two guilds^{46,47}. However, these plant-pollinator associations do not account for some of the traits (i.e., pollen and nectar quantity) that had greater relevance on the species network roles (Fig. 4) because of insufficient data availability and lack of pollinator efficiency measurements for more complete descriptions of the pollination syndromes⁴. In any case, it is worth noting that this general pattern emerges in field-measured plant-pollinator networks, where other local factors such as species relative abundances, will surely explain part of the observed variability^{17,48,49}.

To conclude, we provide the first description of plant floral and reproductive biology trade-offs using a large global dataset of plant traits and interactions with floral visitors. This allowed us to identify the major reproductive strategies of flowering plants and how plant life strategies determine their interaction with floral visitor guilds. Although the percentage of explained variation found with the first two axes (~50%) is lower than other studies that have evaluated life histories with vegetative traits^{24,25}, this percentage is consistent with the largest and most recent study that has characterized life strategies with both vegetative and reproductive (i.e., semelparity and iteroparity) traits¹². Future work needs to integrate these floral and reproductive compromises with the already described vegetative and physiological trade-offs²⁴ to create a more

comprehensive spectrum of trait variation⁵. Further, the varying level of phylogenetic signal among traits deserves further attention to understand evolutionary changes on mating and flower morphology in response to pollinators^{50,51}. Finally, a more complete description of the macroecological patterns of plant-pollinator interactions including unrepresented areas of the world⁵² and traits that have been largely ignored (e.g., floral rewards and plant mating system) are needed to progress in our understanding of the global patterns in plant-pollinator interactions.

MATERIALS AND METHODS

Plant-pollinator network studies. We selected 28 studies from 18 different countries that constituted a total of 64 plant-pollinator networks. These studies recorded plant-pollinator interactions in natural systems and were selected so that we had broad geographical representation. Although these studies differ in sampling effort and methodology, all studies provided information about plant-pollinator interactions (weighted and non-weighted), which we used to build a database of plant species that are likely to be animal pollinated. Many of these networks are freely available either as published studies^{53,54} or available in online archives (e.g., ‘The Web of Life’⁵³ and ‘Mangal’⁵⁵). In total, our network dataset (see Supplementary Table S1) constituted 60 weighted (interaction frequency) and 4 unweighted (presence/absence of the interaction) networks, each sampled at a unique location and year, as well as eight meta-webs where interactions were pooled across several locations and multiple years.

Taxonomy of plants and pollinators. All species names, genera, families and orders were retrieved and standardized from the taxonomy data sources NCBI (<https://www.ncbi.nlm.nih.gov/taxonomy>) for plants and ITIS (<https://www.itis.gov/>) for pollinators, using the R package *taxize*⁵⁶. We filled the ‘not found’ searches manually using <http://www.theplantlist.org/> and <http://www.mobot.org/> for plants and <http://www.catalogueoflife.org/> for floral visitors.

Functional traits. We selected 20 different functional traits based on their relevance to plant reproduction and data availability (Table 1). These included twelve quantitative and eight categorical traits belonging to three broader trait groupings (13 floral, 4 reproductive biology and 3 vegetative, Supplementary Information). For each plant species, we undertook an extensive literature and online search across a wide range of resources (plant databases, online floras, books, journals and images). From a total of 30,120 cells (20 columns \times 1,506 species) we were able to fill 24,341 cells (80.8% of the dataset, see Supplementary Fig. S1 for missing values information for each trait).

Phylogenetic Distance. We calculated the phylogenetic distance between different plant species using the function *get_tree* from the package *rtrees* (<https://github.com/daijiang/rtrees>), which downloads phylogenetic distances from the extended R implementation of the Open Tree of Life^{57,58}.

Data Imputation. Trait missing values were imputed with the function *missForest*⁵⁹ which allows imputation of data sets with continuous and categorical variables. We accounted for the phylogenetic distance among species on the imputation process by including the eigenvectors of a principal component analysis of the phylogenetic distance (PCoA) which has been shown to improve the performance of *missForest*⁶⁰. To extract the eigenvectors, we used the function *PVRdecomp* from the package *PVR*⁶¹ based on a previous conceptual framework that considers phylogenetic eigenvectors⁶². Although the variable of autonomous selfing had a high percentage of missing values (68%), we were able to solve this by back transforming the qualitative column of autonomous selfing to numerical. The categories of ‘none’, ‘low’, ‘medium’ and ‘high’ were converted to representative percentages of each category 0%, 13%, 50.5% and 88% respectively. This reduced the percentage of missing values for this column from 68% to 35% and allowed the imputation of this variable. However, we were unable to include nectar and pollen traits on the imputation process because of the high percentage of missing values (Supplementary Fig. S1). Hence, the imputed dataset had 1,506 species,

Table 1 | Quantitative and categorical traits used in this study.

Quantitative traits		Categorical traits		
Type	Traits	Type	Traits	Categories
Vegetative	Plant height (m)	Vegetative	Lifepan	Short-lived Perennial
Floral	Flower width (mm)	Vegetative	Life form	Herb Shrub Tree
Floral	Flower length (mm)	Floral	Flower shape	Brush Campanulate Capitulum Open Papilionaceous Tube
Floral	Inflorescence width (mm)	Floral	Flower symmetry	Actinomorphic Zygomorphic
Floral	Style length (mm)	Floral	Nectar	Presence Absence
Floral	Ovules per flower	Reproductive biology	Autonomous selfing	None Low Medium High
Floral	Flowers per plant	Reproductive biology	Compatibility system	Self-incomp. Part. self-comp. Self-comp.
Floral	Nectar (μ l)	Reproductive biology	Breeding system	Hermaphrodite Monoecious Dioecious
Floral	Nectar (mg)			
Floral	Nectar concentration (%)			
Floral	Pollen grains per flower			
Reproductive biology	Autonomous selfing (fruit set)			

seven categorical and eight numerical variables and 5.79% of missing values. Further, we conducted an additional imputation on the filtered raw data by filled cells of either pollen grains per flower or microlitres of nectar. This subset comprised 755 species, 8.01% missing values and all traits but milligrams of nectar (~50% of missing values) were included in the imputation process.

Plant strategies. We explored the trade-offs between different quantitative plant functional traits with a phylogenetically informed Principal Component Analysis (pPCA). We did not include the quantitative variables of flower length and inflorescence width because they were highly and moderately correlated to flower width respectively (Pearson's correlation = 0.72, $P < 0.01$ and Pearson's correlation = 0.36, $P < 0.01$ respectively), and thus we avoided overemphasizing flower size on the spectrum of trait variation. Although qualitative traits were not included in the dimensionality reduction analysis, we also investigated the association of the different qualitative traits with the main axes of trait variation. Prior to the analyses, we excluded outliers and standardized the data. Due to the high sensitivity of dimensionality reduction to outliers, we excluded values within the 2.5th–97.5th percentile range⁶³, and thus our final dataset had 1,236 species. Then, we log transformed the variables to reduce the influence of outliers and z-transformed ($X = 0$, $SD = 1$) so that all variables were within the same numerical range. We performed the pPCA using the function *phyl.pca* from the package *phytools*⁶⁴ with the method lambda (λ) that calculates the phylogenetic correlation between 0 (phylogenetic independence) and 1 (shared evolutionary history) and we implemented the mode covariance because values for each variables were on the same scale following transformation⁶⁵. Moreover, to corroborate that our imputation of missing values did not affect our results, we conducted a pPCA on the full dataset without missing values (see Supplementary Fig. S2). We found little difference between the explained variance with the imputed dataset (51.08%) and the dataset without missing values (52.87%). In addition, the loadings on each principal component had a similar contribution and correlation patterns, with the exception of plant height which showed slight variations between the imputed and non-imputed dataset. Finally, we conducted an additional

phylogenetic informed principal component analysis for the subset of species with pollen and nectar quantity. For this, we included all quantitative traits considered in the main pPCA plus pollen grains and microlitres of nectar per flower.

Phylogenetic signal of traits. We calculated the phylogenetic signal of the different quantitative traits on the imputed dataset with the full set of species ($N = 1,506$) with the package *phytools* version 0.7-70⁶⁴ and we used Pagel's λ as a measurement of the phylogenetic signal. However, for pollen and nectar traits, phylogenetic signal was calculated only on the subset of species that had quantitative information for these traits ($N = 755$).

Network analyses. Analyses were conducted on the subset of 60 weighted networks sampled in a unique flowering season and site, which included 556 plant and 1,126 pollinator species. These networks were analysed in their qualitative (presence-absence) and quantitative (interaction frequency) form. First, we analysed the binary version of these weighted networks with presence-absence information that assumes equal weight across interactions. Second, we analysed the untransformed weighted networks with interaction frequency that accounts for the intensity of the interaction. Although floral visitors are not always pollinators and interaction frequency does not consider each pollinator species efficiency⁶⁶, interaction frequency can provide valuable information of the contribution of floral visitors to pollination^{67,68}. In total, our network dataset (excluding meta-webs and non-weighted networks) included 2,256 interactions of Anthophila-Hymenoptera (i.e., bees) with plants, 1,768 non-Syrphidae-Diptera interactions, 845 Syrphidae interactions, 437 Lepidoptera interactions, 432 Coleoptera interactions and 362 non-Anthophila-Hymenoptera interactions. Sampling methods varied across networks but this was accounted for in analyses by considering them in the random effects of the modelling process. All analyses were conducted in R version 4.0.3.

Visitation patterns. We used Bayesian modelling (see below for details) to explore

the effect of floral visitor groups and the main axes of trait variation (pPCA with imputed dataset) on both qualitative (presence/absence) and quantitative (visitation rate) floral interactions per plant species. For this, we divided floral visitors into six main guilds that differ in life form, behaviour and are likely to play a similar ecological role: (i) Hymenoptera-Anthophila (bees), (ii) Hymenoptera-non-Anthophila (non-bee Hymenoptera), (iii) Syrphidae-Diptera, (iv) non-Syrphidae-Diptera, (v) Lepidoptera and (vi) Coleoptera. Moreover, because Hymenoptera-Anthophila was the most represented group with 2,256 records and had the highest frequency of visits of all groups, we also explored the presence-absence of interaction and visitation rate of the main Hymenoptera-Anthophila families (Andrenidae, Apidae, Colletidae, Halictidae and Megachilidae) on the trait space. In addition, we found that *Apis mellifera* was the floral visitor with the largest proportion of records counted (7.55% of the total). This finding is consistent with previous research showing that *A. mellifera* was the most frequent floral visitor in a similar dataset of 80 plant-pollinator networks in natural ecosystems⁶⁹. Hence, to control for the effect of *A. mellifera* on the observed visitation patterns of bees, we conducted an analogous analysis with presence-absence of interaction and visitation rate excluding *A. mellifera*. We found that *A. mellifera*, was partly driving some of the observed trends on PC1 (Supplementary Fig. S3). However, we did not detect major differences on PC2 and PC3.

We implemented Bayesian generalized linear mixed models using the R package *brms*⁷⁰. We modelled the frequency of visits as a function of the main axes of plant trait variation and their interactions with floral visitor functional groups (Visits ~ PC1 x FGs + PC2 x FGs + PC3 x FGs). Because we were interested in possible differences in the visitation patterns among floral visitors groups to plants with different strategies, we included interactions between the main axes of trait variation (PC1, PC2 and PC3) and the floral visitor guilds. In this model, we added a nested random effect of networks nested within the study system to capture the variation in networks among studies and within networks. Moreover, we included the phylogenetic covariance matrix as a random factor due to the possible shared evolutionary histories of species and therefore lack

of independence across them. We specified this model with a zero inflated negative binomial distribution and weakly informative priors from the brms function. We run this model for 3,000 iterations and with previous 1,000 warm up iterations. We set delta (Δ) to 0.99 to avoid divergent transitions and visualized the posterior predictive checks with the function *pp_check* using the *bayesplot* package⁷¹.

Plant species functional roles. We investigated whether different quantitative traits determined plant species functional roles using Bayesian modelling and regression trees. For this, we selected simple and complementary species-level network metrics commonly applied in bipartite network studies⁷² with a straightforward ecological interpretation relevant to our research goals. The different plant species-level metrics were: (i) sum of visits per plant species; (ii) normalized degree, calculated as the number of links per plant species divided by the total possible number of partners; and (iii) specialization⁷³, which measures the deviation of an expected random choice of the available interaction partners and ranges between 0 (maximum generalization) and 1 (maximum specialization). Normalized degree and specialization were calculated with the *specieslevel* function from the R package *bipartite*⁷².

First, we modelled the distinct plant species metrics (sum of visits, normalized degree and plant specialization) as a function of the three main axes of trait variation (plant species level metric ~ PC1 + PC2 + PC3). For each response variable (i.e., each plant species level metric), we used different distribution families (zero inflated negative binomial for the sum of visits, weibull for normalized degree and zero one inflated beta for specialization). Finally, we used the same random factors, model settings and conducted the same posterior predictive checks for each model as detailed above in the ‘visitation patterns section’.

Second, to better understand these complex trait relationships, we used regression trees. Regression trees are recursive algorithms which can detect complex relationships among predictors and allow identification of the relevance of specific trait combinations

on species functional roles. We focused exclusively on quantitative traits because almost all categorical traits were statistically associated with the first two axes of trait variation (Supplementary Table S2). We conducted this analysis using the *rpart* package⁷⁴ version 4.1-15 with method ‘*anova*’ with a minimum of 50 observations per terminal node and we used the *rpart.plot* package⁷⁵ version 3.0.9 to plot the regression trees. We considered the species level indices as response variables (interaction frequency, normalized degree and specialization) and we performed one regression tree per metric using the different quantitative traits as predictors. We calculated two regression trees per plant species-level metric, one for the full set of species and another for the subset of species for which we had pollen and nectar traits. We focused on regression trees that included floral rewards because they consistently showed pollen and nectar traits as being the best for explaining the different species-level metrics (see Supplementary Fig. S4).

References

1. Barrett, S. C. H. The evolution of plant sexual diversity. *Nat. Rev. Genet.* **3**, 274–284 (2002).
2. Schiestl, F. P. & Johnson, S. D. Pollinator-mediated evolution of floral signals. *Trends Ecol. Evol.* **28**, 307–315 (2013).
3. Fenster, C. B., Armbruster, W. S., Wilson, P., Dudash, M. R. & Thomson, J. D. Pollination Syndromes and Floral Specialization. *Annu. Rev. Ecol. Evol. Syst.* **35**, 375–403 (2004).
4. Dellinger, A. S. Pollination syndromes in the 21st century: Where do we stand and where may we go? *New Phytol.* **228**, 1193–1213 (2020).
5. Roddy, A. B. *et al.* Towards the flower economics spectrum. *New Phytol.* **229**, 665–672

449 (2021).

450 6. Baude, M. *et al.* Historical nectar assessment reveals the fall and rise of floral resources
451 in Britain. *Nature* **530**, 85–88 (2016).

452 7. Munoz, F., Violle, C. & Cheptou, P.-O. CSR ecological strategies and plant mating
453 systems: Outcrossing increases with competitiveness but stress-tolerance is related to
454 mixed mating. *Oikos* **125**, 1296–1303 (2016).

455 8. Grossenbacher, D. L. *et al.* Self-compatibility is over-represented on islands. *New*
456 *Phytol.* **215**, 469–478 (2017).

457 9. Moeller, D. A. *et al.* Global biogeography of mating system variation in seed plants.
458 *Ecol. Lett.* **20**, 375–384 (2017).

459 10. Sargent, R. D. & Ackerly, D. D. Plant–pollinator interactions and the assembly of
460 plant communities. *Trends Ecol. Evol.* **23**, 123–130 (2008).

461 11. Rech, A. R. *et al.* The macroecology of animal versus wind pollination: Ecological
462 factors are more important than historical climate stability. *Plant Ecol. Divers.* **9**, 253–262
463 (2016).

464 12. Salguero-Gómez, R. *et al.* Fast-slow continuum and reproductive strategies structure
465 plant life-history variation worldwide. *Proc. Natl. Acad. Sci. U.S.A.* **113**, 230–235 (2016).

466 13. Rüger, N. *et al.* Beyond the fastSlow continuum: Demographic dimensions structur-
467 ing a tropical tree community. *Ecol. Lett.* **21**, 1075–1084 (2018).

468 14. Dehling, D. M., Jordano, P., Schaefer, H. M., Böhning-Gaese, K. & Schleuning, M.
469 Morphology predicts species' functional roles and their degree of specialization in
470 plant–frugivore interactions. *Proc. Royal Soc. B* **283**, 20152444 (2016).

471 15. Rosas-Guerrero, V. *et al.* A quantitative review of pollination syndromes: Do floral

- 472 traits predict effective pollinators? *Ecol. Lett.* **17**, 388–400 (2014).
- 473 16. Stang, M., Klinkhamer, P. G. L., Waser, N. M., Stang, I. & van der Meijden, E.
474 Size-specific interaction patterns and size matching in a plantPollinator interaction web.
475 *Ann. Bot.* **103**, 1459–1469 (2009).
- 476 17. Bartomeus, I. *et al.* A common framework for identifying linkage rules across
477 different types of interactions. *Funct. Ecol.* **30**, 1894–1903 (2016).
- 478 18. Waser, N. M., Chittka, L., Price, M. V., Williams, N. M. & Ollerton, J. Generalization
479 in pollination systems, and why it matters. *Ecology* **77**, 1043–1060 (1996).
- 480 19. Olesen, J. M. & Jordano, P. Geographic patterns in plant–pollinator mutualistic
481 networks. *Ecology* **83**, 2416–2424 (2002).
- 482 20. Tur, C., Castro-Urgal, R. & Traveset, A. Linking Plant Specialization to Dependence
483 in Interactions for Seed Set in Pollination Networks. *PLoS One* **8**, e78294 (2013).
- 484 21. Devaux, C., Lepers, C. & Porcher, E. Constraints imposed by pollinator behaviour
485 on the ecology and evolution of plant mating systems. *J. Evol. Biol.* **27**, 1413–1430 (2014).
- 486 22. Kattge, J. *et al.* TRY a global database of plant traits. *Glob. Chang. Biol.* **17**, 2905–2935
487 (2011).
- 488 23. Salguero-Gómez, R. *et al.* The compadre Plant Matrix Database: An open online
489 repository for plant demography. *J. Ecol.* **103**, 202–218 (2015).
- 490 24. Díaz, S. *et al.* The global spectrum of plant form and function. *Nature* **529**, 167–171
491 (2016).
- 492 25. Carmona, C. P. *et al.* Erosion of global functional diversity across the tree of life. *Sci.*
493 *Adv.* **7**, eabf2675 (2021).
- 494 26. Wright, I. J. *et al.* The worldwide leaf economics spectrum. *Nature* **428**, 821–827

495 (2004).

496 27. Chave, J. *et al.* Towards a worldwide wood economics spectrum. *Ecol. Lett.* **12**,
497 351–366 (2009).

498 28. Laughlin, D. C. *et al.* Root traits explain plant species distributions along climatic
499 gradients yet challenge the nature of ecological trade-offs. *Nat. Ecol. Evol.* 1–12 (2021).

500 29. E-Vojtkó, A., Bello, F. de, Durka, W., Kühn, I. & Götzenberger, L. The neglected
501 importance of floral traits in trait-based plant community assembly. *J. Veg. Sci.* **31**,
502 529–539 (2020).

503 30. Barrett, S. C. H. Mating strategies in flowering plants: The outcrossing-selfing
504 paradigm and beyond. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **358**, 991–1004 (2003).

505 31. Goodwillie, C. *et al.* Correlated evolution of mating system and floral display traits
506 in flowering plants and its implications for the distribution of mating system variation.
507 *New Phytol.* **185**, 311–321 (2010).

508 32. Ollerton, J., Winfree, R. & Tarrant, S. How many flowering plants are pollinated by
509 animals? *Oikos* **120**, 321–326 (2011).

510 33. Bartomeus, I. Understanding Linkage Rules in Plant-Pollinator Networks by Using
511 Hierarchical Models That Incorporate Pollinator Detectability and Plant Traits. *PLoS*
512 *One* **8**, e69200 (2013).

513 34. Olito, C. & Fox, J. W. Species traits and abundances predict metrics of plantPollinator
514 network structure, but not pairwise interactions. *Oikos* **124**, 428–436 (2015).

515 35. Rowe, L. *et al.* Flower traits associated with the visitation patterns of bees. *Oecologia*
516 **193**, 511–522 (2020).

517 36. Lázaro, A., Jakobsson, A. & Totland, Ø. How do pollinator visitation rate and seed

- 518 set relate to species' floral traits and community context? *Oecologia* **173**, 881–893 (2013).
- 519 37. Junker, R. R. *et al.* Specialization on traits as basis for the niche-breadth of flower
520 visitors and as structuring mechanism of ecological networks. *Funct. Ecol.* **27**, 329–341
521 (2013).
- 522 38. Coux, C., Rader, R., Bartomeus, I. & Tylianakis, J. M. Linking species functional
523 roles to their network roles. *Ecol. Lett.* **19**, 762–770 (2016).
- 524 39. Ibanez, S. Optimizing size thresholds in a plant-pollinator interaction web: Towards
525 a mechanistic understanding of ecological networks. *Oecologia* **170**, 233–242 (2012).
- 526 40. Eklöf, A. *et al.* The dimensionality of ecological networks. *Ecol. Lett.* **16**, 577–583
527 (2013).
- 528 41. Sargent, R. D., Goodwillie, C., Kalisz, S. & Ree, R. H. Phylogenetic evidence for a
529 flower size and number trade-off. *Am. J. Bot.* **94**, 2059–2062 (2007).
- 530 42. Kettle, C. J. *et al.* Ecological Implications of a Flower Size/Number Trade-Off in
531 Tropical Forest Trees. *PLoS One* **6**, e16111 (2011).
- 532 43. Healy, K., Ezard, T. H. G., Jones, O. R., Salguero-Gómez, R. & Buckley, Y. M. Animal
533 life history is shaped by the pace of life and the distribution of age-specific mortality
534 and reproduction. *Nat. Ecol. Evol.* **3**, 1217–1224 (2019).
- 535 44. Hegland, S. J. & Totland, Ø. Relationships between species' floral traits and pollina-
536 tor visitation in a temperate grassland. *Oecologia* **145**, 586–594 (2005).
- 537 45. Kaiser-Bunbury, C. N., Vázquez, D. P., Stang, M. & Ghazoul, J. Determinants of the
538 microstructure of plantPollinator networks. *Ecology* **95**, 3314–3324 (2014).
- 539 46. Palmer, T. M., Stanton, M. L. & Young, T. P. Competition and coexistence: Exploring
540 mechanisms that restrict and maintain diversity within mutualist guilds. *Am. Nat.* **162**,

541 S63–S79 (2003).

542 47. Phillips, R. D., Peakall, R., van der Niet, T. & Johnson, S. D. Niche perspectives on
543 plantPollinator interactions. *Trends Plant Sci.* **25**, 779–793 (2020).

544 48. Vázquez, D. P. *et al.* Species abundance and asymmetric interaction strength in
545 ecological networks. *Oikos* **116**, 1120–1127 (2007).

546 49. Encinas-Viso, F., Revilla, T. A. & Etienne, R. S. Phenology drives mutualistic network
547 structure and diversity. *Ecol. Lett.* **15**, 198–208 (2012).

548 50. Gervasi, D. D. L. & Schiestl, F. P. Real-time divergent evolution in plants driven by
549 pollinators. *Nat. Commun.* **8**, 14691 (2017).

550 51. Mackin, C. R., Peña, J. F., Blanco, M. A., Balfour, N. J. & Castellanos, M. C. Rapid
551 evolution of a floral trait following acquisition of novel pollinators. *J. Ecol.* **109**, 2234–
552 2246 (2021).

553 52. Poisot, T. *et al.* Global knowledge gaps in species interaction networks data. *J.*
554 *Biogeogr.* **48**, 1552–1563 (2021).

555 53. Fortuna, M. A. *et al.* Nestedness versus modularity in ecological networks: Two
556 sides of the same coin? *J. Anim. Ecol.* **79**, 811–817 (2010).

557 54. Carvalheiro, L. G. *et al.* The potential for indirect effects between co-flowering plants
558 via shared pollinators depends on resource abundance, accessibility and relatedness.
559 *Ecol. Lett.* **17**, 1389–1399 (2014).

560 55. Poisot, T. *et al.* Mangal making ecological network analysis simple. *Ecography* **39**,
561 384–390 (2016).

562 56. Chamberlain, S. *et al.* Taxize: Taxonomic information from around the web. r

- 563 package version 0.9.99 (2020). at <https://cran.R-project.org/package=taxize>.
- 564 57. Smith, S. A. & Brown, J. W. Constructing a broadly inclusive seed plant phylogeny.
565 *Am. J. Bot.* **105**, 302–314 (2018).
- 566 58. Jin, Y. & Qian, H. V. PhyloMaker: An R package that can generate very large
567 phylogenies for vascular plants. *Ecography* **42**, 1353–1359 (2019).
- 568 59. Stekhoven, D. J. & Bühlmann, P. MissForestNon-parametric missing value imputa-
569 tion for mixed-type data. *Bioinformatics* **28**, 112–118 (2012).
- 570 60. Penone, C. *et al.* Imputation of missing data in life-history trait datasets: Which
571 approach performs the best? *Methods Ecol. Evol.* **5**, 961–970 (2014).
- 572 61. Chamberlain, S. *et al.* Package ‘PVR’. r package version 0.3 (2018). at [https://cran.R-](https://cran.R-project.org/package=PVR)
573 [project.org/package=PVR](https://cran.R-project.org/package=PVR).
- 574 62. Diniz-Filho, J. A. F. *et al.* On the selection of phylogenetic eigenvectors for ecological
575 analyses. *Ecography* **35**, 239–249 (2012).
- 576 63. Legendre, P. & Legendre, L. *Numerical ecology*. (Elsevier, Amsterdam, 2012).
- 577 64. Revell, L. J. Phytools: An R package for phylogenetic comparative biology (and
578 other things). *Methods Ecol. Evol.* **3**, 217–223 (2012).
- 579 65. Abdi, H. & Williams, L. J. Principal component analysis. *WIREs Comp. Stats.* **2**,
580 433–459 (2010).
- 581 66. Ballantyne, G., Baldock, K. C. R. & Willmer, P. G. Constructing more informative
582 plantPollinator networks: Visitation and pollen deposition networks in a heathland
583 plant community. *Proc. Royal Soc. B* **282**, 20151130 (2015).
- 584 67. Vázquez, D. P., Morris, W. F. & Jordano, P. Interaction frequency as a surrogate for

- 585 the total effect of animal mutualists on plants. *Ecol. Lett.* **8**, 1088–1094 (2005).
- 586 68. Vázquez, D. P. *et al.* The strength of plantPollinator interactions. *Ecology* **93**, 719–725
587 (2012).
- 588 69. Hung, K.-L. J., Kingston, J. M., Albrecht, M., Holway, D. A. & Kohn, J. R. The
589 worldwide importance of honey bees as pollinators in natural habitats. *Proc. Royal Soc.*
590 *B* **285**, 20172140 (2018).
- 591 70. Bürkner, P.-C. Brms: An R package for Bayesian multilevel models using Stan. *J.*
592 *Stat. Softw.* **80**, 1–28 (2017).
- 593 71. Gabry, J., Simpson, D., Vehtari, A., Betancourt, M. & Gelman, A. Visualization in
594 Bayesian workflow. *J. R. Stat. Soc. Ser. A Stat. Soc.* **182**, 389–402 (2019).
- 595 72. Dormann, C. F., Gruber, B. & Fründ, J. Introducing the bipartite package: Analysing
596 ecological networks. *interaction* **1**, (2008).
- 597 73. Blüthgen, N., Menzel, F. & Blüthgen, N. Measuring specialization in species interac-
598 tion networks. *BMC Ecol.* **6**, 9 (2006).
- 599 74. Therneau, T., Atkinson, B., Ripley, B. & Ripley, M. B. Package ‘rpart’. r package
600 version 4.1-15 (2015). at <https://cran.R-project.org/package=rpart>.
- 601 75. Milborrow, S. Rpart. plot: Plot rpart Models. An Enhanced Version of plot. rpart. r
602 package version 3.0.9 (2015). at <https://cran.R-project.org/package=rpart.plot>.