

Trade-offs among plant reproductive traits determine interactions with floral visitors

Jose B. Lanuza^{1,2*}, Romina Rader¹, Jamie Stavert³, Liam K. Kendall⁴,
Manu E. Saunders¹ and Ignasi Bartomeus²

Plant life strategies are constrained by cost-benefit trade-offs that determine plant form and function. Despite the recent advances in the understanding of plant trade-offs with vegetative and physiological traits, little is known about the plant reproductive economics and how they constrain plant life strategies and shape interactions with floral visitors. Here, we investigated plant reproductive trade-offs and their association with floral visitors with a dataset of 1506 plant species with 17 reproductive and 3 vegetative traits from 28 plant-pollinator network studies across 18 countries. Then, we tested whether a species reproductive strategy predicted plant-pollinator interactions (presence-absence and visitation rate) and if the interaction strategy (its role in a pollination network) was predicted by individual traits. We found that 51.8% of trait variation was explained by two independent axes that encompassed plant form and function. Specifically, the first axis indicated the presence of a trade-off between flower number and flower size (PC1, 26.72%) and the second axis indicated a trade-off for the level of pollinator dependency (PC2, 25.08%). The reproductive trade-offs explained some of the qualitative interaction with pollinators (presence-absence) but none of the variability of visitation rates. However, noteworthy differences across pollinator guilds were found in both cases. Finally, we found that specific traits with special relevance of floral rewards were key in the understanding of the species network role. Our results highlight the need to consider plant reproductive trade-offs to improve our understanding of plant life strategies and plant-pollinator interactions at broader spatial scales.

¹ School of Environmental and Rural Science, University of New England, Armidale, New South Wales 2350, Australia. ² Estación Biológica de Doñana (EBD-CSIC), E-41092 Seville, Spain. ³ Department of Conservation | Te Papa Atawhai, Auckland, New Zealand. ⁴ Centre for Environmental and Climate Science, Lund University, Sölvegatan 37, S-223 62 Lund, Sweden. * e-mail: barragansljose@gmail.com

There is an astonishing diversity of floral structures and plant reproductive strategies among flowering plants^{1,2}, which have long been of interest to pollination biologists in terms of their relevance to plant-pollinator interactions. However, most studies that have explored plant reproductive biology (e.g., mating and compatibility systems) and floral trait (e.g., flower size or nectar provision) variation have concentrated on

the individual or community level and thus, broader macroecological patterns remain poorly investigated^{3–7}. Indeed, studies depicting species' life history strategies generally focus on vegetative traits and rarely consider reproductive biology traits⁸. As a consequence, a unified framework that explores the compromises among floral traits and their relevance to plant life strategies is currently lacking¹⁰. At the same time, there is growing interest in the determinants of plant-pollinator interactions via trait-based approaches¹¹ and trait-matching analyses¹². However, floral traits have been overlooked beyond highly specialised plant-pollinator systems^{10,13} and the role of plant reproductive biology remains little explored in plant-pollinator interactions (but see references^{14,15}).

With the recent availability of large trait databases, plant ecological strategies are increasingly being examined^{16,17}, and are facilitating the identification of global patterns and constraints of plant form and function^{8,18,19}. However, the main focus has been on vegetative traits such as leaf²⁰ or wood²¹ trade-offs with little or no attention given to reproductive biology and floral traits²², also critical to plant form and function. For instance, short lived versus perennial species tend to have low versus high levels of outcrossing, respectively^{6,23}. Further, outcrossing levels are positively correlated with flower size²⁴. In addition, the presence of costly rewards (e.g., pollen or nectar) and showy flowers or floral displays can only be understood through consideration of plant species' reliance upon animal pollination (pollinator dependence) and their role in attracting pollinators²⁵. Hence, exploring plant life strategies with reproductive biology and floral trade-offs, in conjunction with their pollinator dependence, is necessary for a balanced understanding of plant economics.

Several studies have identified links between plant traits and plant-pollinator network properties^{3,26,27}. Moreover, plant traits can also define species' network roles (e.g., specialists vs generalists). For example, species that occupy trait space extremes are more likely to exhibit higher specialization and be more reliant on trait-matching^{28,29}. This morphological matching between plant and floral visitors can determine plant-

pollinator interactions, and thus shape their interaction network structure^{30,31}. Despite the increasing knowledge of the relevance of traits on the species network roles, little is known about how plant reproductive and floral traits determine plant species' network roles at a macroecological scale.

Here, we explore the potential trade-offs among plant floral and reproductive biology traits and how these influence the structure of plant-pollinator networks. First, we identify the major axes of trait variation and trade-offs that determine plant form and function. Second, we investigate how plant species' position in trait-space influences their interaction patterns with different guilds of floral visitors. Finally, we investigate how the main axes of trait variation and individual traits influence plant species roles within networks.

RESULTS

Plant strategies. The phylogenetically informed principal component analysis (pPCA) captured by the first two and three axes 51.8% and 70.97% of trait variation, respectively (Fig. 1 and Supplementary Fig. S5) and had a phylogenetic correlation (λ) of 0.76. The first principal component (PC1) represented 26.72% of the trait variation and indicated a trade-off between flower number and flower size. We refer to this axis as the 'flower number - flower size trade-off', as already described in previous studies^{32,33}. Hence, one end of the spectrum comprised species with high investment in flower number and plant height but small flower size, short style length and low ovule number. The other end of this spectrum comprised species that were short in height and invested in large flowers, long styles, many ovules, but few flowers. The main contributing traits to PC1 were plant height, flower number, ovule number and flower size (loadings $> |0.5|$; Supplementary Table S3) but style length also contributed moderately on PC1 (loading = -0.33). The second principal component (PC2) represented 25.05% of the trait variation and indicated a trade-off between low and high pollinator dependence. We refer to

102 this axis as the ‘pollinator dependence trade-off’. The main driver of trait variation on
103 PC2 was autonomous selfing (loading = 0.85) but the other traits (except ovule number)
104 also made moderate contributions (loadings from 0.27 to 0.4; Supplementary Table S3).
105 We found that high pollinator dependence was associated with larger and a higher
106 number of flowers, greater plant height and longer styles. In contrast, species with high
107 levels of autonomous selfing tended to have fewer and smaller flowers, had shorter
108 styles and were shorter in height. Further, PC3 explained a considerable amount of trait
109 variability (19.17%) and the main contributors to this axis were style length (loading
110 = -0.66) and the degree of autonomous selfing (loading = -0.51). The remaining traits,
111 apart from ovule number, were moderately correlated to changes on PC3 (loadings
from -0.23 to -0.46; Supplementary Table S3). Thus, because style length was correlated
with all traits on PC3 and was the main driver of trait variation, we refer to this axis
as the ‘style length trade-off’. Further, the pPCA with the subset of species that had
nectar and pollen quantity data showed that nectar quantity (microlitres of nectar per
flower) was positively associated with flower size, style length and ovule number (PC1,
23.40%); and pollen quantity (pollen grains per flower) was positively correlated with
flower number and plant height and negatively associated with autonomous selfing
(PC2, 21.67%; Supplementary Fig. S6). This pPCA explained similar variance with the
first two principal components (45.07%) and similar associations of traits despite some
variability in the loadings (Supplementary Table S4).



****Fig. 1 | Plant life-history strategies.**** Phylogenetically informed principal component analysis (pPCA) of 1,236 plant species from 28 plant-pollinator network studies. The solid arrows indicate the direction of the different quantitative traits (flower number, plant height, style length, flower size, ovule number and level of autonomous selfing) across the two main axes of trait variation. The length of the arrows indicate the weight of the variables on each principal component and the dashed lines show the opposed direction of trait variation. The icons at both ends of arrows and dashed lines illustrate the extreme form of the trait continuum.

112 We found that most categorical traits were statistically associated with the first two axes
 113 of trait variation (Fig. 2 and Supplementary Table S2). Flower symmetry, which was
 114 only associated with PC2 (Sum of squares = 8.51, F-value = 14.72, $P < 0.01$), and nectar
 115 provision, which was independent of PC1 and PC2 (PC1: Sum of squares = 0.37, F-value

116 = 0.29 , $P = 0.59$; PC2: Sum of squares = 0.83, F-value = 1.43, $P = 0.23$) showed lack of
117 statistical association. In addition, we found (with a Tukey test) statistical differences
118 between the different levels of categorical traits in the trait space (Supplementary Fig.
119 S7). Regarding self compatibility, we found larger differences on PC2 (i.e., species
120 with unisexual flowers that were self incompatible were statistically differentiated from
121 species with partial or full self compatibility; Supplementary Fig. S7a and Fig. S7b). Life
122 forms differed statistically across both axes of trait variation and followed a gradient
123 of larger life forms (trees and shrubs) with higher pollinator dependence to smaller
124 ones (herbs) with lower pollinator dependence (Supplementary Fig. S7c and Fig. S7d).
125 Consequently, lifespan also followed this gradient but perennial and short lived species
126 only differed statistically on PC2 (Supplementary Fig. S7e and Fig. S7f). Species with
127 unisexual flowers (monoecious and dioecious) were clustered on both extremes of
128 the first two principal components and had the highest pollinator dependence and
129 highest number of flowers (Supplementary Fig. S7g and Fig. S7h). Moreover, we
130 found that the campanulate and capitulum flower shapes were differentiated from tube,
131 papilionaceous, open and brush shapes in the trait space. The former morphologies
132 had larger flowers and greater pollinator dependence, while the latter had higher
133 flower number and greater autonomous selfing (Supplementary Fig. S7i and Fig. S7j).
134 Regarding flower symmetry, zygomorphic flowers were associated with lower levels of
135 pollinator dependence, whereas actinomorphic flowers had higher levels of pollinator
136 dependence (Supplementary Fig. S7k and Fig. S7l).




Fig. 2 | Location of the different qualitative traits on the trait space. The panel is composed by the traits that showed statistical association with the first two axes of trait variation: compatibility system (a), life form (b), lifespan (c), breeding system (d), flower shape (e) and flower symmetry (f).

Phylogenetic signal of traits. We found a strong phylogenetic signal ($P < 0.01$) in all quantitative traits (Supplementary Table S5). The traits that showed the highest phylogenetic signal were ovule number ($\lambda = 1$), pollen grains per flower ($\lambda = 1$) and plant height ($\lambda = 0.96$), followed by flower length ($\lambda = 0.75$), flower width ($\lambda = 0.73$), number of flowers per plant ($\lambda = 0.69$) and nectar concentration ($\lambda = 0.65$). The traits that showed a moderate phylogenetic signal were inflorescence width ($\lambda = 0.57$), style length ($\lambda = 0.49$) and autonomous selfing ($\lambda = 0.34$). Finally, microliters of nectar per flower showed the lowest phylogenetic signal of all traits ($\lambda = 0.14$).

Visitation patterns. The main axes of trait variation explained little of the overall visitation rates ($conditionalR2 = 0.31$; $marginalR2 = 0.06$) but showed relevant trends when we explored the interaction with the different floral visitor guilds (Fig. 3). All floral visitor guilds visited plant species with higher pollinator dependence more frequently (PC2; Fig. 3b and Fig. 3e). In addition, all guilds other than Syrphids and Lepidoptera (i.e., all Hymenoptera, non-Syrphidae-Diptera and Coleoptera) showed greater visitation rates on species with small numerous flowers (PC1; Fig. 3a and Fig. 3d). On the style length trade-off (PC3), Anthophila-Hymenoptera, Lepidoptera and Non-Anthophila-Hymenoptera showed greater visitation rates on plant species with larger styles and higher levels of selfing; while Syrphidae, Non-Syrphidae-Diptera and Coleoptera showed higher visitation rates on species with shorter styles and lower selfing (Fig. 3c and Fig. 3f). Furthermore, when running an additional model that separates the most represented families of Anthophila-Hymenoptera (bees; $marginalR2 = 0.30$; $conditionalR2 = 0.03$) showed that the family Apidae was the main driver of the observed patterns (Supplementary Fig. S8).




Fig. 3 | Interaction (yes/no) and visitation rates across the three main axes of trait variation per floral visitor guild. Fitted posterior estimates of the presence/absence of interaction (a, b and c) and number of visits (d, e and f) made by the different floral visitors guilds in relation to PC1, PC2 and PC3. PC1 represents the flower number - flower size trade-off, PC2 represents the pollinator dependence trade-off and PC3, the style length trade-off. For visualization purposes, due to large differences between the visitation rates of bees and the rest of guilds, the number of visits was log-transformed (Y-axis of lower panel).

Plant species functional roles. The variance of the different plant species-level network metrics was poorly explained by the three main axes of trait variation (Supplementary Fig. S9; interaction frequency ~ PCs, *conditionalR2* = 0.11, *marginalR2* = 0.02; normalized degree ~ PCs, *conditionalR2* = 0.24, *marginalR2* = 0.02; and, specialization ~ PCs, *conditionalR2* = 0.37, *marginalR2* = 0.03). Overall, the most notable trends were found on PC1 and PC3 for interaction frequency and specialization. On the flower number - flower size trade-off (PC1), interaction frequency was higher for plant species with more flowers but was lower for plant species with larger flowers. On PC1, specialization showed the opposite trend. On the style length trade-off (PC3), interaction frequency was lower for plants with shorter styles and lower autonomous selfing and higher for species with longer styles and higher autonomous selfing. Again, specialization showed the opposite trend to interaction frequency.

When we further investigate which combination of traits drive plant network roles, we show that the regression tree for visitation frequency was best explained by plant height, nectar concentration and style length (Fig. 4a). Specifically, species taller than 3.9 m had the highest interaction frequency, while species that were shorter than 3.9 m and had a nectar concentration lower than 16% had the lowest interaction frequency. Normalized degree was best explained by nectar concentration, pollen grains per flower, plant height, flower width and autonomous selfing (Fig. 4b). Species with a nectar concentration over 49% had the highest levels of normalized degree, whereas species with nectar concentration lower than 49%, more than 21,000 pollen grains per flower and height less than 0.78 m had the lowest normalized degree. Finally, specialization was best explained by plant height, ovule number, pollen grains per flower and autonomous selfing (Fig. 4c). Overall, plant species with the highest specialization were shorter than 1.3 m, had more than 14,000 pollen grains per flower and autonomously self-pollinated less than 11% of their fruits. In contrast, species taller or equal than 5.1 m and with lower than 14 ovules per flower had the lowest specialization values.




Fig. 4 | Contribution of traits in plant's network roles. Regression tree analysis of interaction frequency (log-transformed), normalized degree and specialization for the subset of species with quantitative data for pollen and nectar traits. The superior value inside the node indicates the mean value of the different species-level metric and the lower value, the percentage of species that are considered in each node. Thus, the top node has the mean value of the named trait for the 100% of species. Each node has a yes/no question and when the condition is fulfilled, the branch turns to the 'yes' direction and when not, to the 'no' direction. This rationale is followed in all the regression trees as indicated in the first branch division of the topmost node of each tree.

DISCUSSION

Here, we show that plant species exhibit clear trade-offs in their floral, reproductive biology and vegetative traits. These trade-offs are differentiated on three main axes of trait variation: (i) flower number - flower size, (ii) pollinator dependence and (iii) style length. Although these axes of trait variation did not explain overall visitation rates, we found that plant life strategies were clearly associated with different floral visitor guilds. Interestingly, pollen and nectar related traits were better than all other traits for characterizing plant species functional roles within their interaction networks.

Over 50% of plant trait variation was captured by the flower number - flower size and pollinator dependence trade-offs. Variation in these two axes of trait variation appears to be associated with the commonly described 'fast-slow continuum' in plant⁸ and animal³⁴ life-history strategies as indicated by the different floral and reproductive biology traits associated with plant height, life form and lifespan. The 'slow' part of the continuum (i.e., tall trees and shrubs) had plant species with many flowers, few ovules, higher pollinator dependence, greater occurrence of self-incompatibility and more complex breeding systems (e.g., monoecious and dioecious species). In contrast, species that employed the 'fast' strategy (i.e., short herbs), had fewer flowers, more ovules, greater occurrence of self compatibility and lower pollinator dependence. Further, on the first two axes of trait variation, we found additional support for the positive association already described in a previous study between outcrossing rate and floral display²⁴. This trend was further confirmed when looking at the specific correlation between floral display and autonomous selfing for our dataset (see Supplementary Fig. S10).

Despite the low predictive power of the main axes of trait variation on overall visitation patterns, we found commonalities among and within floral visitors guilds across these axes that indicate the influence of life-history strategies on floral visitation rate. For

example, all guilds showed an increasing trend of visitation on plant species with higher pollinator dependence, which were also associated with larger floral displays and greater pollen quantities (Fig. 1 and Supplementary Fig. S6). This trend is consistent with previous plant-pollinator studies that show higher visitation rates on species that make greater reproductive investment³⁵⁻³⁷. In regard to the flower number - flower size and style length trade-offs, different guilds showed contrasting visitation rates, which could be associated with different pollination syndromes at a macroecological scale. For instance, the guilds of bees and syrphids were clearly associated with opposing life-strategies on PC1 and PC3 (Fig. 3) indicating a possible niche partitioning among these two guilds^{38,39}. However, these plant-pollinator associations do not account for some of the traits (i.e., pollen and nectar quantity) that had greater relevance on the species network roles (Fig. 4) because of insufficient data availability and lack of pollinator efficiency measurements for more complete descriptions of the pollination syndromes¹³. In any case, it is worth noting that this general pattern emerges in field-measured plant-pollinator networks, where other local factors such as species relative abundances, will surely explain part of the observed variability^{12,40,41}.

To conclude, we provide the first description of plant floral and reproductive biology trade-offs using a large global dataset of plant traits and interactions with floral visitors. This allowed us to identify the major reproductive strategies of flowering plants and how plant life strategies determine their interaction with floral visitor guilds. Although the percentage of explained variation found with the first two axes (~50%) is lower than other studies that have evaluated life histories with vegetative traits^{18,19}, this percentage is consistent with the largest and most recent study that has characterized life strategies with both vegetative and reproductive (i.e., semelparity and iteroparity) traits⁸. Future work needs to integrate these floral and reproductive compromises with the already described vegetative and physiological trade-offs¹⁸ to create a more comprehensive spectrum of trait variation¹⁰. Further, the varying level of phylogenetic signal among traits deserves further attention to understand evolutionary changes on mating and flower morphology in response to pollinators^{42,43}. Finally, a more complete

description of the macroecological patterns of plant-pollinator interactions including unrepresented areas of the world⁴⁴ and traits that have been largely ignored (e.g., floral rewards and plant mating system) are needed to progress in our understanding of the global patterns in plant-pollinator interactions.

MATERIALS AND METHODS

Plant-pollinator network studies. We selected 28 studies from 18 different countries that constituted a total of 64 plant-pollinator networks. These studies recorded plant-pollinator interactions in natural systems and were selected so that we had broad geographical representation. Although these studies differ in sampling effort and methodology, all studies provided information about plant-pollinator interactions (weighted and non-weighted), which we used to build a database of plant species that are likely to be animal pollinated. Many of these networks are freely available either as published studies^{3,45} or available in online archives (e.g., ‘The Web of Life’⁴⁵ and ‘Mangal’⁴⁶). In total, our network dataset (see Supplementary Table S1) constituted 60 weighted (interaction frequency) and 4 unweighted (presence/absence of the interaction) networks, each sampled at a unique location and year, as well as eight meta-webs where interactions were pooled across several locations and multiple years.

Taxonomy of plants and pollinators. All species names, genera, families and orders were retrieved and standardized from the taxonomy data sources NCBI (<https://www.ncbi.nlm.nih.gov/taxonomy>) for plants and ITIS (<https://www.itis.gov/>) for pollinators, using the R package *taxize*⁴⁷. We filled the ‘not found’ searches manually using <http://www.theplantlist.org/> and <http://www.mobot.org/> for plants and <http://www.catalogueoflife.org/> for floral visitors.

Functional traits. We selected 20 different functional traits based on their relevance to plant reproduction and data availability (Table 1). These included twelve quantitative

and eight categorical traits belonging to three broader trait groupings (13 floral, 4 reproductive biology and 3 vegetative, Supplementary Information). For each plant species, we undertook an extensive literature and online search across a wide range of resources (plant databases, online floras, books, journals and images). From a total of 30,120 cells (20 columns \times 1,506 species) we were able to fill 24,341 cells (80.8% of the dataset, see Supplementary Fig. S1 for missing values information for each trait).

Phylogenetic Distance. We calculated the phylogenetic distance between different plant species using the function *get_tree* from the package *rtrees* (<https://github.com/daijiang/rtrees>), which downloads phylogenetic distances from the extended R implementation of the Open Tree of Life^{48,49}.

Data Imputation. Trait missing values were imputed with the function *missForest*⁵⁰ which allows imputation of data sets with continuous and categorical variables. We accounted for the phylogenetic distance among species on the imputation process by including the eigenvectors of a principal component analysis of the phylogenetic distance (PCoA) which has been shown to improve the performance of *missForest*⁵¹. To extract the eigenvectors, we used the function *PVRdecomp* from the package *PVR*⁵² based on a previous conceptual framework that considers phylogenetic eigenvectors⁵³. Although the variable of autonomous selfing had a high percentage of missing values (68%), we were able to solve this by back transforming the qualitative column of autonomous selfing to numerical. The categories of ‘none’, ‘low’, ‘medium’ and ‘high’ were converted to representative percentages of each category 0%, 13%, 50.5% and 88% respectively. This reduced the percentage of missing values for this column from 68% to 35% and allowed the imputation of this variable. However, we were unable to include nectar and pollen traits on the imputation process because of the high percentage of missing values (Supplementary Fig. S1). Hence, the imputed dataset had 1,506 species, seven categorical and eight numerical variables and 5.79% of missing values. Further, we conducted an additional imputation on the filtered raw data by filled cells of either

Table 1 | Quantitative and categorical traits used in this study.

Quantitative traits		Categorical traits		
Type	Traits	Type	Traits	Categories
Vegetative	Plant height (m)	Vegetative	Lifepan	Short-lived Perennial
Floral	Flower width (mm)	Vegetative	Life form	Herb Shrub Tree
Floral	Flower length (mm)	Floral	Flower shape	Brush Campanulate Capitulum Open Papilionaceous Tube
Floral	Inflorescence width (mm)	Floral	Flower symmetry	Actinomorphic Zygomorphic
Floral	Style length (mm)	Floral	Nectar	Presence Absence
Floral	Ovules per flower	Reproductive biology	Autonomous selfing	None Low Medium High
Floral	Flowers per plant	Reproductive biology	Compatibility system	Self-incomp. Part. self-comp. Self-comp.
Floral	Nectar (μ l)	Reproductive biology	Breeding system	Hermaphrodite Monoecious Dioecious
Floral	Nectar (mg)			
Floral	Nectar concentration (%)			
Floral	Pollen grains per flower			
Reproductive biology	Autonomous selfing (fruit set)			

pollen grains per flower or microlitres of nectar. This subset comprised 755 species, 8.01% missing values and all traits but milligrams of nectar (~50% of missing values) were included in the imputation process.

Plant strategies. We explored the trade-offs between different quantitative plant functional traits with a phylogenetically informed Principal Component Analysis (pPCA). We did not include the quantitative variables of flower length and inflorescence width because they were highly and moderately correlated to flower width respectively (Pearson's correlation = 0.72, $P < 0.01$ and Pearson's correlation = 0.36, $P < 0.01$ respectively), and thus we avoided overemphasizing flower size on the spectrum of trait variation. Although qualitative traits were not included in the dimensionality reduction analysis, we also investigated the association of the different qualitative traits with the main axes of trait variation. Prior to the analyses, we excluded outliers and standardized the data. Due to the high sensitivity of dimensionality reduction to outliers, we excluded values within the 2.5th–97.5th percentile range⁵⁴, and thus our final dataset had 1,236 species. Then, we log transformed the variables to reduce the influence of outliers and z-transformed ($X = 0$, $SD = 1$) so that all variables were within the same numerical range. We performed the pPCA using the function *phyl.pca* from the package *phytools*⁵⁵ with the method lambda (λ) that calculates the phylogenetic correlation between 0 (phylogenetic independence) and 1 (shared evolutionary history) and we implemented the mode covariance because values for each variables were on the same scale following transformation⁵⁶. Moreover, to corroborate that our imputation of missing values did not affect our results, we conducted a pPCA on the full dataset without missing values (see Supplementary Fig. S2). We found little difference between the explained variance with the imputed dataset (51.08%) and the dataset without missing values (52.87%). In addition, the loadings on each principal component had a similar contribution and correlation patterns, with the exception of plant height which showed slight variations between the imputed and non-imputed dataset. Finally, we conducted an additional phylogenetic informed principal component analysis for the subset of species with pollen and nectar quantity. For this, we included all quantitative traits considered in

the main pPCA plus pollen grains and microlitres of nectar per flower.

Phylogenetic signal of traits. We calculated the phylogenetic signal of the different quantitative traits on the imputed dataset with the full set of species ($N = 1,506$) with the package *phytools* version 0.7-70⁵⁵ and we used Pagel's λ as a measurement of the phylogenetic signal. However, for pollen and nectar traits, phylogenetic signal was calculated only on the subset of species that had quantitative information for these traits ($N = 755$).

Network analyses. Analyses were conducted on the subset of 60 weighted networks sampled in a unique flowering season and site, which included 556 plant and 1,126 pollinator species. These networks were analysed in their qualitative (presence-absence) and quantitative (interaction frequency) form. First, we analysed the binary version of these weighted networks with presence-absence information that assumes equal weight across interactions. Second, we analysed the untransformed weighted networks with interaction frequency that accounts for the intensity of the interaction. Although floral visitors are not always pollinators and interaction frequency does not consider each pollinator species efficiency⁵⁷, interaction frequency can provide valuable information of the contribution of floral visitors to pollination^{58,59}. In total, our network dataset (excluding meta-webs and non-weighted networks) included 2,256 interactions of Anthophila-Hymenoptera (i.e., bees) with plants, 1,768 non-Syrphidae-Diptera interactions, 845 Syrphidae interactions, 437 Lepidoptera interactions, 432 Coleoptera interactions and 362 non-Anthophila-Hymenoptera interactions. Sampling methods varied across networks but this was accounted for in analyses by considering them in the random effects of the modelling process. All analyses were conducted in R version 4.0.3.

Visitation patterns. We used Bayesian modelling (see below for details) to explore the effect of floral visitor groups and the main axes of trait variation (pPCA with imputed dataset) on both qualitative (presence/absence) and quantitative (visitation rate)

floral interactions per plant species. For this, we divided floral visitors into six main guilds that differ in life form, behaviour and are likely to play a similar ecological role: (i) Hymenoptera-Anthophila (bees), (ii) Hymenoptera-non-Anthophila (non-bee Hymenoptera), (iii) Syrphidae-Diptera, (iv) non-Syrphidae-Diptera, (v) Lepidoptera and (vi) Coleoptera. Moreover, because Hymenoptera-Anthophila was the most represented group with 2,256 records and had the highest frequency of visits of all groups, we also explored the presence-absence of interaction and visitation rate of the main Hymenoptera-Anthophila families (Andrenidae, Apidae, Colletidae, Halictidae and Megachilidae) on the trait space. In addition, we found that *Apis mellifera* was the floral visitor with the largest proportion of records counted (7.55% of the total). This finding is consistent with previous research showing that *A. mellifera* was the most frequent floral visitor in a similar dataset of 80 plant-pollinator networks in natural ecosystems⁶⁰. Hence, to control for the effect of *A. mellifera* on the observed visitation patterns of bees, we conducted an analogous analysis with presence-absence of interaction and visitation rate excluding *A. mellifera*. We found that *A. mellifera*, was partly driving some of the observed trends on PC1 (Supplementary Fig. S3). However, we did not detect major differences on PC2 and PC3.

We implemented Bayesian generalized linear mixed models using the R package *brms*⁶¹. We modelled the frequency of visits as a function of the main axes of plant trait variation and their interactions with floral visitor functional groups (Visits ~ PC1 x FGs + PC2 x FGs + PC3 x FGs). Because we were interested in possible differences in the visitation patterns among floral visitors groups to plants with different strategies, we included interactions between the main axes of trait variation (PC1, PC2 and PC3) and the floral visitor guilds. In this model, we added a nested random effect of networks nested within the study system to capture the variation in networks among studies and within networks. Moreover, we included the phylogenetic covariance matrix as a random factor due to the possible shared evolutionary histories of species and therefore lack of independence across them. We specified this model with a zero inflated negative binomial distribution and weakly informative priors from the *brms* function. We run

this model for 3,000 iterations and with previous 1,000 warm up iterations. We set delta (Δ) to 0.99 to avoid divergent transitions and visualized the posterior predictive checks with the function *pp_check* using the *bayesplot* package⁶².

Plant species functional roles. We investigated whether different quantitative traits determined plant species functional roles using Bayesian modelling and regression trees. For this, we selected simple and complementary species-level network metrics commonly applied in bipartite network studies⁶³ with a straightforward ecological interpretation relevant to our research goals. The different plant species-level metrics were: (i) sum of visits per plant species; (ii) normalized degree, calculated as the number of links per plant species divided by the total possible number of partners; and (iii) specialization⁶⁴, which measures the deviation of an expected random choice of the available interaction partners and ranges between 0 (maximum generalization) and 1 (maximum specialization). Normalized degree and specialization were calculated with the *specieslevel* function from the R package *bipartite*⁶³.

First, we modelled the distinct plant species metrics (sum of visits, normalized degree and plant specialization) as a function of the three main axes of trait variation (plant species level metric \sim PC1 + PC2 + PC3). For each response variable (i.e., each plant species level metric), we used different distribution families (zero inflated negative binomial for the sum of visits, weibull for normalized degree and zero one inflated beta for specialization). Finally, we used the same random factors, model settings and conducted the same posterior predictive checks for each model as detailed above in the 'visitation patterns section'.

Second, to better understand these complex trait relationships, we used regression trees. Regression trees are recursive algorithms which can detect complex relationships among predictors and allow identification of the relevance of specific trait combinations on species functional roles. We focused exclusively on quantitative traits because almost all categorical traits were statistically associated with the first two axes of trait variation

(Supplementary Table S2). We conducted this analysis using the *rpart* package⁶⁵ version 4.1-15 with method ‘*anova*’ with a minimum of 50 observations per terminal node and we used the *rpart.plot* package⁶⁶ version 3.0.9 to plot the regression trees. We considered the species level indices as response variables (interaction frequency, normalized degree and specialization) and we performed one regression tree per metric using the different quantitative traits as predictors. We calculated two regression trees per plant species-level metric, one for the full set of species and another for the subset of species for which we had pollen and nectar traits. We focused on regression trees that included floral rewards because they consistently showed pollen and nectar traits as being the best for explaining the different species-level metrics (see Supplementary Fig. S4).

References

1. Barrett, S. C. H. The evolution of plant sexual diversity. *Nat. Rev. Genet.* **3**, 274–284 (2002).
2. Schiestl, F. P. & Johnson, S. D. Pollinator-mediated evolution of floral signals. *Trends Ecol. Evol.* **28**, 307–315 (2013).
3. Carvalheiro, L. G. *et al.* The potential for indirect effects between co-flowering plants via shared pollinators depends on resource abundance, accessibility and relatedness. *Ecol. Lett.* **17**, 1389–1399 (2014).
4. Baude, M. *et al.* Historical nectar assessment reveals the fall and rise of floral resources in Britain. *Nature* **530**, 85–88 (2016).
5. Munoz, F., Violle, C. & Cheptou, P.-O. CSR ecological strategies and plant mating systems: Outcrossing increases with competitiveness but stress-tolerance is related to

- 429 mixed mating. *Oikos* **125**, 1296–1303 (2016).
- 430 6. Moeller, D. A. *et al.* Global biogeography of mating system variation in seed plants.
431 *Ecol. Lett.* **20**, 375–384 (2017).
- 432 7. Grossenbacher, D. L. *et al.* Self-compatibility is over-represented on islands. *New*
433 *Phytol.* **215**, 469–478 (2017).
- 434 8. Salguero-Gómez, R. *et al.* Fast-slow continuum and reproductive strategies structure
435 plant life-history variation worldwide. *Proc. Natl. Acad. Sci. U.S.A.* **113**, 230–235 (2016).
- 436 9. Rüger, N. *et al.* Beyond the fastSlow continuum: Demographic dimensions structur-
437 ing a tropical tree community. *Ecol. Lett.* **21**, 1075–1084 (2018).
- 438 10. Roddy, A. B. *et al.* Towards the flower economics spectrum. *New Phytol.* **229**, 665–672
439 (2021).
- 440 11. Fenster, C. B., Armbruster, W. S., Wilson, P., Dudash, M. R. & Thomson, J. D.
441 Pollination Syndromes and Floral Specialization. *Annu. Rev. Ecol. Evol. Syst.* **35**,
442 375–403 (2004).
- 443 12. Bartomeus, I. *et al.* A common framework for identifying linkage rules across
444 different types of interactions. *Funct. Ecol.* **30**, 1894–1903 (2016).
- 445 13. Dellinger, A. S. Pollination syndromes in the 21st century: Where do we stand and
446 where may we go? *New Phytol.* **228**, 1193–1213 (2020).
- 447 14. Tur, C., Castro-Urgal, R. & Traveset, A. Linking Plant Specialization to Dependence
448 in Interactions for Seed Set in Pollination Networks. *PLoS One* **8**, e78294 (2013).
- 449 15. Devaux, C., Lepers, C. & Porcher, E. Constraints imposed by pollinator behaviour
450 on the ecology and evolution of plant mating systems. *J. Evol. Biol.* **27**, 1413–1430 (2014).
- 451 16. Kattge, J. *et al.* TRY a global database of plant traits. *Glob. Chang. Biol.* **17**, 2905–2935

452 (2011).

453 17. Salguero-Gómez, R. *et al.* The compadre Plant Matrix Database: An open online
454 repository for plant demography. *J. Ecol.* **103**, 202–218 (2015).

455 18. Díaz, S. *et al.* The global spectrum of plant form and function. *Nature* **529**, 167–171
456 (2016).

457 19. Carmona, C. P. *et al.* Erosion of global functional diversity across the tree of life. *Sci.*
458 *Adv.* **7**, eabf2675 (2021).

459 20. Wright, I. J. *et al.* The worldwide leaf economics spectrum. *Nature* **428**, 821–827
460 (2004).

461 21. Chave, J. *et al.* Towards a worldwide wood economics spectrum. *Ecol. Lett.* **12**,
462 351–366 (2009).

463 22. E-Vojtkó, A., Bello, F. de, Durka, W., Kühn, I. & Götzenberger, L. The neglected
464 importance of floral traits in trait-based plant community assembly. *J. Veg. Sci.* **31**,
465 529–539 (2020).

466 23. Barrett, S. C. H. Mating strategies in flowering plants: The outcrossing-selfing
467 paradigm and beyond. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **358**, 991–1004 (2003).

468 24. Goodwillie, C. *et al.* Correlated evolution of mating system and floral display traits
469 in flowering plants and its implications for the distribution of mating system variation.
470 *New Phytol.* **185**, 311–321 (2010).

471 25. Ollerton, J., Winfree, R. & Tarrant, S. How many flowering plants are pollinated by
472 animals? *Oikos* **120**, 321–326 (2011).

473 26. Lázaro, A., Hegland, S. J. & Totland, Ø. The relationships between floral traits and
474 specificity of pollination systems in three Scandinavian plant communities. *Oecologia*

- 475 157, 249–257 (2008).
- 476 27. Bartomeus, I. Understanding Linkage Rules in Plant-Pollinator Networks by Using
477 Hierarchical Models That Incorporate Pollinator Detectability and Plant Traits. *PLoS*
478 *One* **8**, e69200 (2013).
- 479 28. Junker, R. R. *et al.* Specialization on traits as basis for the niche-breadth of flower
480 visitors and as structuring mechanism of ecological networks. *Funct. Ecol.* **27**, 329–341
481 (2013).
- 482 29. Coux, C., Rader, R., Bartomeus, I. & Tylianakis, J. M. Linking species functional
483 roles to their network roles. *Ecol. Lett.* **19**, 762–770 (2016).
- 484 30. Stang, M., Klinkhamer, P. G. L., Waser, N. M., Stang, I. & van der Meijden, E.
485 Size-specific interaction patterns and size matching in a plantPollinator interaction web.
486 *Ann. Bot.* **103**, 1459–1469 (2009).
- 487 31. Ibanez, S. Optimizing size thresholds in a plant-pollinator interaction web: Towards
488 a mechanistic understanding of ecological networks. *Oecologia* **170**, 233–242 (2012).
- 489 32. Sargent, R. D., Goodwillie, C., Kalisz, S. & Ree, R. H. Phylogenetic evidence for a
490 flower size and number trade-off. *Am. J. Bot.* **94**, 2059–2062 (2007).
- 491 33. Kettle, C. J. *et al.* Ecological Implications of a Flower Size/Number Trade-Off in
492 Tropical Forest Trees. *PLoS One* **6**, e16111 (2011).
- 493 34. Healy, K., Ezard, T. H. G., Jones, O. R., Salguero-Gómez, R. & Buckley, Y. M. Animal
494 life history is shaped by the pace of life and the distribution of age-specific mortality
495 and reproduction. *Nat. Ecol. Evol* **3**, 1217–1224 (2019).
- 496 35. Hegland, S. J. & Totland, Ø. Relationships between species' floral traits and pollina-

- 497 tor visitation in a temperate grassland. *Oecologia* **145**, 586–594 (2005).
- 498 36. Lázaro, A., Jakobsson, A. & Totland, Ø. How do pollinator visitation rate and seed
499 set relate to species' floral traits and community context? *Oecologia* **173**, 881–893 (2013).
- 500 37. Kaiser-Bunbury, C. N., Vázquez, D. P., Stang, M. & Ghazoul, J. Determinants of the
501 microstructure of plantPollinator networks. *Ecology* **95**, 3314–3324 (2014).
- 502 38. Palmer, T. M., Stanton, M. L. & Young, T. P. Competition and coexistence: Exploring
503 mechanisms that restrict and maintain diversity within mutualist guilds. *Am. Nat.* **162**,
504 S63–S79 (2003).
- 505 39. Phillips, R. D., Peakall, R., van der Niet, T. & Johnson, S. D. Niche perspectives on
506 plantPollinator interactions. *Trends Plant Sci.* **25**, 779–793 (2020).
- 507 40. Vázquez, D. P. *et al.* Species abundance and asymmetric interaction strength in
508 ecological networks. *Oikos* **116**, 1120–1127 (2007).
- 509 41. Encinas-Viso, F., Revilla, T. A. & Etienne, R. S. Phenology drives mutualistic network
510 structure and diversity. *Ecol. Lett.* **15**, 198–208 (2012).
- 511 42. Gervasi, D. D. L. & Schiestl, F. P. Real-time divergent evolution in plants driven by
512 pollinators. *Nat. Commun.* **8**, 14691 (2017).
- 513 43. Mackin, C. R., Peña, J. F., Blanco, M. A., Balfour, N. J. & Castellanos, M. C. Rapid
514 evolution of a floral trait following acquisition of novel pollinators. *J. Ecol.* **109**, 2234–
515 2246 (2021).
- 516 44. Poisot, T. *et al.* Global knowledge gaps in species interaction networks data. *J.*
517 *Biogeogr.* **48**, 1552–1563 (2021).
- 518 45. Fortuna, M. A. *et al.* Nestedness versus modularity in ecological networks: Two

- sides of the same coin? *J. Anim. Ecol.* **79**, 811–817 (2010).
46. Poisot, T. *et al.* Mangal making ecological network analysis simple. *Ecography* **39**, 384–390 (2016).
47. Chamberlain, S. *et al.* Taxize: Taxonomic information from around the web. r package version 0.9.99 (2020). at <https://cran.R-project.org/package=taxize>.
48. Smith, S. A. & Brown, J. W. Constructing a broadly inclusive seed plant phylogeny. *Am. J. Bot.* **105**, 302–314 (2018).
49. Jin, Y. & Qian, H. V. PhyloMaker: An R package that can generate very large phylogenies for vascular plants. *Ecography* **42**, 1353–1359 (2019).
50. Stekhoven, D. J. & Bühlmann, P. MissForestNon-parametric missing value imputation for mixed-type data. *Bioinformatics* **28**, 112–118 (2012).
51. Penone, C. *et al.* Imputation of missing data in life-history trait datasets: Which approach performs the best? *Methods Ecol. Evol.* **5**, 961–970 (2014).
52. Chamberlain, S. *et al.* Package ‘PVR’. r package version 0.3 (2018). at <https://cran.R-project.org/package=PVR>.
53. Diniz-Filho, J. A. F. *et al.* On the selection of phylogenetic eigenvectors for ecological analyses. *Ecography* **35**, 239–249 (2012).
54. Legendre, P. & Legendre, L. *Numerical ecology*. (Elsevier, Amsterdam, 2012).
55. Revell, L. J. Phytools: An R package for phylogenetic comparative biology (and other things). *Methods Ecol. Evol.* **3**, 217–223 (2012).
56. Abdi, H. & Williams, L. J. Principal component analysis. *WIREs Comp. Stats.* **2**,

540 433–459 (2010).

541 57. Ballantyne, G., Baldock, K. C. R. & Willmer, P. G. Constructing more informative
542 plantPollinator networks: Visitation and pollen deposition networks in a heathland
543 plant community. *Proc. Royal Soc. B* **282**, 20151130 (2015).

544 58. Vázquez, D. P., Morris, W. F. & Jordano, P. Interaction frequency as a surrogate for
545 the total effect of animal mutualists on plants. *Ecol. Lett.* **8**, 1088–1094 (2005).

546 59. Vázquez, D. P. *et al.* The strength of plantPollinator interactions. *Ecology* **93**, 719–725
547 (2012).

548 60. Hung, K.-L. J., Kingston, J. M., Albrecht, M., Holway, D. A. & Kohn, J. R. The
549 worldwide importance of honey bees as pollinators in natural habitats. *Proc. Royal Soc.*
550 *B* **285**, 20172140 (2018).

551 61. Bürkner, P.-C. Brms: An R package for Bayesian multilevel models using Stan. *J.*
552 *Stat. Softw.* **80**, 1–28 (2017).

553 62. Gabry, J., Simpson, D., Vehtari, A., Betancourt, M. & Gelman, A. Visualization in
554 Bayesian workflow. *J. R. Stat. Soc. Ser. A Stat. Soc.* **182**, 389–402 (2019).

555 63. Dormann, C. F., Gruber, B. & Fründ, J. Introducing the bipartite package: Analysing
556 ecological networks. *interaction* **1**, (2008).

557 64. Blüthgen, N., Menzel, F. & Blüthgen, N. Measuring specialization in species interac-
558 tion networks. *BMC Ecol.* **6**, 9 (2006).

559 65. Therneau, T., Atkinson, B., Ripley, B. & Ripley, M. B. Package ‘rpart’. r package
560 version 4.1-15 (2015). at <https://cran.R-project.org/package=rpart>.

561 66. Milborrow, S. Rpart. plot: Plot rpart Models. An Enhanced Version of plot. rpart. r
562 package version 3.0.9 (2015). at <https://cran.R-project.org/package=rpart.plot>.