An explicit defining quartic in 265 variables

10 November 2026

Abstract

Under the specified modular-realization, elliptic, pole-realization and prime-pattern inputs, we specify an integer quartic in 265 total variables defining the integers in the rationals. The quadratic contact-height estimate with exponent 7/3 becomes 7/6 on two disjoint five-contact blocks. Three comparison scales allow the use of the effective height theorem for four elliptic abscissas. A common elliptic chart uses four field coordinates, and all comparison ratios use the plain formal parameter. Four factors encode sixteen local tests; shared fourth powers distinguish their valuation classes. Inverse weights allow the numerator fourth powers to enter without separate outputs. The complete quadratic system has 237 rows; the expanded quartic has exactly 3485 monomials and content one. The fixed parameter $N = 14521$ is sufficient. The resulting coefficient height is $149^{58086} + 1$, with 419333 binary digits. All coefficient and numerical threshold assertions have independent exact certificates. No additional *abc* conjecture is used.

1 The family and its arithmetic scope

Write h for the absolute logarithmic Weil height; for a reduced rational number r/s, with $s > 0$, it is $h(r/s) = \log \max(|r|, s)$. For an integer polynomial F, write $H(F)$ for the largest absolute coefficient. All variable counts include the distinguished input X. For a fixed integer $N \geq 1$ let $A_N = 149^{2N+1}$. The complete tables

`quadratic-family265-v19.json`
`polynomial-family265-v19.json`

specify 237 primitive integer quadratic polynomials and their sum of squares

$$P_N(X, Y_1, \ldots, Y_{264}) = \sum_{j=1}^{237} Q_{N,j}^2.$$

The parameter A_N is a coefficient, not a variable.

Theorem 1.1. *Assume the arithmetic premises stated below. For every integer $N \geq 14521$ and every $x \in \mathbb{Q}$,*

$$x \in \mathbb{Z} \quad \Longleftrightarrow \quad \exists \mathbf{y} \in \mathbb{Q}^{264} \ P_N(x, \mathbf{y}) = 0.$$

For every $N \geq 1$, the polynomial P_N has degree exactly four, content one, and 3485 nonzero coordinate monomials. For $N \geq 53$,

$$H(P_N) = 149^{4N+2} + 1.$$

In particular $A_{14521} = 149^{29043}$ gives a defining quartic of coefficient height $149^{58086} + 1$, whose binary length is 419333. The coefficient tables specify this polynomial completely.

Set

$$K=\mathbb{Q}(\rho),\quad \rho^2=2,\quad D=(3-\rho)^2=11-6\rho,\qquad E:y^2=x^3-Dx,$$
$$P=2(1+\rho,2\rho)=\left(\frac{57-28\rho}{8},\frac{-660+417\rho}{32}\right).$$

We retain the following arithmetic premises, with the precise domains and hypotheses recorded in [1, 2, 3]:

1. $2E(K)=\mathbb{Z}P$ and $h(x([n]P))\geq n^2-6$ for every nonzero integer n, together with the specified point charts and group domains.

2. The modular-realization input of [7, Definition 6.1], with its normalized coefficient character. The multiplicity theorem of [8, Theorem 4.1] proves

$$h(u)\leq HM_5(u)^{7/3},\qquad H=2^{221234}. \tag{1}$$

   Here $M_5(u)$ is the odd contact radical outside 76176 for $\infty,0,1,2401,-343/32$, and u avoids the four finite contacts. Explicitly, if $u=r/s$ is reduced with $s>0$, then $M_5(u)$ is the product of primes $p>76176$ having a positive odd valuation on at least one of

$$s,\quad r,\quad r-s,\quad r-2401s,\quad 32r+343s.$$

   The estimate is a consequence of the stated modular input, rather than an additional *abc* assumption.

3. The pole obstruction outside $2,3,17$, and its simultaneous realization for positive inputs less than 17 with square reduced denominator and the stated local squareclasses and valuation bounds at those three primes. Its normalized quadratic equations are retained.

4. The simultaneous prime-pattern, conjugate reduction-order, finite congruence, and real approximation assertions used by the source. The weighted selection and fixed denominator dilation are justified from the cited analytic estimates in [10].

The effective sum-height theorem used below is proved in [6]; it adds no arithmetic premise. Contact amplification is an elementary consequence of (1). Exact coefficient verification does not prove the retained arithmetic premises.

2 Two contact blocks and three effective comparison scales

Put

$$\Theta=\{0,-1/2,-1,1/2399,-32/407\},\qquad \Theta_2=\Theta\cup(2-\Theta),\qquad F_2(t)=\prod_{\theta\in\Theta_2}(t-\theta).$$

The ten contacts are distinct, omit one, and are symmetric about one. Let Σ consist of the primes at most 76176 and the prime 2029147. Outside this set, all contact differences and all numbers $1-\theta$ are units. Define $M_2^\circ(t)$ as the product of primes $p\notin\Sigma$ for which $v_p(t-\theta)$ is positive and odd for some $\theta\in\Theta_2$. Exact factorizations in the independent certificate verify this exceptional set. In particular

$$F_2(1)=-\frac{82429855236}{7878870169},\qquad M_2^\circ(1)=1.$$

Set $E_2=2029147$ and $H_2=2^{221284}$. Then

$$h(t)\leq H_2M_2^\circ(t)^{7/6}\qquad (t\notin\Theta_2). \tag{2}$$

Indeed, apply (1) to $2 + 1/t$ and $2 + 1/(2 - t)$. The inverse projective matrices have maximum absolute row sums three and seven, respectively. If M_+, M_- are the two block radicals with Σ omitted, the original radicals are at most $E_2 M_+$ and $E_2 M_-$. Outside Σ these radicals are coprime, with $M_+ M_- = M_2^\circ(t)$. Multiplying the two inequalities

$$h(t) - \log 7 \le H E_2^{7/3} M_\pm^{7/3}$$

when the left side is positive gives the exponent $7/6$. If it is nonpositive, the asserted bound is immediate. Finally $E_2 < 2^{21}$ and $\log 7 < 2$ give

$$\log 7 + H E_2^{7/3} < 2 + 2^{221283} < H_2.$$

The retained root points are

$$Q_1 = [n]P, \quad Q_2 = [a]P, \quad Q_3 = [t]P, \quad Q_4 = [b]P, \qquad B = \max(|n|, |a|, |t|, |b|),$$

with nonzero integer labels. Set

$$\mathcal{S} = \sum_{i=1}^{4} x(Q_i) = S_0 + S_1 \rho.$$

Each abscissa is a positive square at both real embeddings, so $S_0 > 12$ and $S_0 > \sqrt{2}|S_1|$. The effective abscissa sum-height theorem [6] gives $B^2 \le 2h(\mathcal{S}) + 80$. Put $U = S_0$ and $V = S_0 + S_1$, so $\mathcal{S} = (1 - \rho)U + \rho V$. The product of the two archimedean coefficient sums is seven. The finite-place coefficients are integral, and therefore $h(\mathcal{S}) \le h(U) + h(V) + \frac{1}{2}\log 7$. Since $\log 7 < 2$, this proves

$$B^2 \le 2h(S_0) + 2h(S_0 + S_1) + 82. \tag{3}$$

Use the three positive rational square scales

$$a_0 = 1, \qquad a_1 = S_0^2, \qquad a_2 = (S_0 + S_1)^2.$$

For each scale put $t_* = a_* \lambda$ and $\delta = t_* - 1$. On the nonzero domain use the balanced contact and the local weight

$$R = \frac{F_2(t_*)}{\delta^{10}}, \qquad q = \frac{\delta^{10}}{F_2(t_*)} - 1 = R^{-1} - 1, \qquad \gamma = \frac{408}{409}. \tag{4}$$

The contact circuit stores $r = \delta^2$ and its powers r^2, r^3. Its actual quadratic normalization is

$$F_2(t_*)(1 + q) = r^2 r^3. \tag{5}$$

Thus $R^{-1} = 1 + q$ when $\delta \ne 0$, and no additional coordinate for this reciprocal is needed. No contact value can occur: $F_2(t_*) = 0$ would force $r = 0$ and $t_* = 1$, contradicting $F_2(1) \ne 0$. The reciprocal-weight equation below imposes $q \ne 0$. The remaining exceptional stratum $\delta = 0$ has $q = -1$ and $M_2^\circ(t_*) = 1$; no value of R is used on that stratum. Every witness constructed in completeness has a selected contact prime with $v_p(q) = -1$, and therefore satisfies $q \ne 0$.

For $j = 0, 1, 2$, write $t_j = a_j \lambda$ and let $M_j = M_2^\circ(t_j)$. Suppose that the kth power of each M_j divides a nonzero integer J. Since $t_0 \ne 0$,

$$2h(S_0) \le h(t_1) + h(t_0), \qquad 2h(S_0 + S_1) \le h(t_2) + h(t_0).$$

Consequently (2) and (3) give

$$B^2 \le 4H_2 |J|^{7/(6k)} + 82 \le 5H_2 |J|^{7/(6k)}. \tag{6}$$

The three radicals need not be pairwise coprime. No ineffective constant occurs in this inequality.

3 A quadratic chart for the common reference point

Write $a_P = x(P)$, $b_P = y(P)$, and set

$$A_P = a_P^2 + D, \qquad B_P = a_P^2 - D, \qquad C_P = D^2 + 6a_P^2 D - 3a_P^4.$$

The constants a_P, b_P, D, A_P, C_P are nonzero. The first four nonvanishing assertions follow from the displayed coordinates and $D = (3-\rho)^2 > 0$ at both real embeddings; exact arithmetic gives

$$C_P = -\frac{94897379}{4096} + \frac{4195821}{256}\rho, \qquad \mathrm{Nm}(C_P) = -\frac{8203357319351}{16777216} \neq 0.$$

Retain four field coordinates S_H, T_H, f, F_0. Put

$$\begin{aligned} L_f &= 2b_P D(a_P T_H - b_P), \\ Q_f &= a_P A_P S_H T_H + b_P A_P S_H - b_P B_P T_H + 4a_P^2 D, \\ L_g &= 2b_P D(a_P S_H + b_P), \\ Q_g &= -a_P A_P S_H T_H - b_P B_P S_H + b_P A_P T_H - 4a_P^2 D. \end{aligned}$$

Impose the three quadratic equations

$$\begin{aligned} L_f f &= Q_f, \\ L_g F_0 &= Q_g, \\ D^2 f F_0 + a_P D(f + F_0) + a_P S_H T_H + b_P(S_H - T_H) + D &= 0. \end{aligned} \tag{7}$$

Lemma 3.1. *These equations describe exactly the points $H \in E(K)$ for which H and $H - P$ are finite with nonzero abscissas, through*

$$S_H = \frac{y(H)}{x(H)}, \qquad T_H = \frac{y(H-P)}{x(H-P)}, \qquad f = \frac{1}{x(H)}, \qquad F_0 = \frac{1}{x(H-P)}.$$

In particular $fF_0 \neq 0$ at every solution. The statement includes all finite exceptional cases of the displayed equations.

Proof. For the algebraic calculation abbreviate $a = a_P$, $b = b_P$, $S = S_H$, $T = T_H$ and $g = F_0$. Thus $b^2 = a(a^2 - D)$. Define

$$\Phi = aS^2T^2 - (a^2 - D)(S^2 + T^2) + 2(a^2 + D)ST + 4aD$$

and the polynomial expressions

$$\begin{aligned} 2bDf_0 &= -(a^2 + D)S + (a^2 - D)T - bS^2 - aS^2T, \\ 2bDg_0 &= (a^2 + D)T - (a^2 - D)S - bT^2 + aST^2. \end{aligned}$$

Direct expansion gives

$$\begin{aligned} Df_0^2 + S^2 f_0 - 1 &= \frac{a^2S^2 - b^2}{4ab^2D}\Phi, \\ Dg_0^2 + T^2 g_0 - 1 &= \frac{a^2T^2 - b^2}{4ab^2D}\Phi, \\ L_f f_0 - Q_f = -a\Phi, &\qquad L_g g_0 - Q_g = a\Phi. \end{aligned} \tag{8}$$

Let $r_f = L_f f - Q_f$, $r_g = L_g g - Q_g$, and $R = D^2 fg + aD(f + g) + aST + b(S - T) + D$. The further exact identity

$$\begin{aligned} L_f L_g R = D^2 r_f r_g &+ (aDL_g + D^2 Q_g) r_f \\ &+ (aDL_f + D^2 Q_f) r_g - aD^2 C_P \Phi \end{aligned} \tag{9}$$

shows that every solution has $\Phi = 0$, including the cases in which one leading coefficient vanishes. Also

$$(2bD)^2 R(f_0, g_0) = D^2\big((a^2 - D) - aST - b(S - T)\big)\Phi.$$

If $L_f L_g \neq 0$, these identities give $f = f_0, g = g_0$. If $L_f = 0$, then $T = b/a$ and $\Phi = 0$ forces

$$S = \frac{a^4 - 6Da^2 + D^2}{2b(a^2 + D)}, \qquad L_g = -\frac{aDC_P}{a^2 + D} \neq 0.$$

Thus $g = g_0 = 1/a$. The coefficient of f in R is $D(a^2 + D)/a \neq 0$, so $f = f_0$. The case $L_g = 0$ is symmetric. In particular both leading coefficients cannot vanish simultaneously; at $S = -b/a, T = b/a$ one has $Q_f = C_P \neq 0$.

Equation (8) now proves $fg \neq 0$ and gives the points $H = (f^{-1}, S/f)$ and $H' = (g^{-1}, T/g)$ on E. Two more identities prove the translation:

$$a(f_0 T + g_0 S) + b(g_0 - f_0) = S + T,$$

$$f_0 g_0 (aS + b)^2 - a(1 - af_0)^2 = -\frac{(aS + b)^2\big(aST + b(S - T) + D\big)}{4b^2 D^2}\Phi.$$

If $x(H) \neq a$, the second identity gives the abscissa of $H - P$ from the chord slope $(x(H)S + b)/(x(H) - a)$. The first states that $H, -H', -P$ are collinear, and therefore gives the required ordinate. If $x(H) = a$, then $S = \pm b/a$. At $S = b/a$ the formula gives $f = -a/D \neq 1/a$, which is impossible. At $S = -b/a$ it gives $H = -P$ and

$$T = -\frac{a^4 - 6Da^2 + D^2}{2b(a^2 + D)}, \qquad g = \frac{4b^2}{(a^2 + D)^2},$$

which are the coordinates of $H' = -2P$ in the stated form. Thus $H' = H - P$ in all cases.

Conversely, the chord-and-tangent formulas for any allowed H give $\Phi = 0$ and the expressions f_0, g_0 above; the same explicit doubling computation covers $H = -P$. The displayed identities then prove all three equations. This proves both directions. □

Write $P = 2R_0$ and $H = [m]R_0 + T_0$, with $T_0 \in E[2](K)$, and put $h = m/2$. This notation is an elliptic index; it does not define multiplication of a point by a nonintegral rational number.

4 The scalar certificate and the point domains

Put $\beta = 51X^2 + 2504$, let $S = Q_2 = [a]P$, and write $C_S = S - P = [a - 1]P$. The scalar normalizer is

$$\begin{aligned} \ell &= 149, \qquad A_N = \ell^{2N+1}, \\ r_s &= \beta(\beta - 1)(\beta - 2401)(32\beta + 343)(2\beta - 1), \\ R_s &= \frac{A_N r_s}{1 + \ell^2 r_s}. \end{aligned} \tag{10}$$

All five factors of r_s are positive, since $\beta \geq 2504$, so $R_s > 0$. The retained square-abscissa chart for S is

$$x_S = h_S^2, \qquad j_S^2 = x_S^2 - D, \qquad S = (x_S, h_S j_S).$$

It has $h_S x_S j_S \neq 0$. The fixed-subtraction coordinate $g_S \in K$ satisfies

$$8(x_P x_S + D)g_S + (39 - 36\rho)h_S + (8 - 14\rho)j_S = 0.$$

The rational component of g_S has a retained inverse witness. At $S = P$ the coordinate g_S is zero, so this guard gives $a \neq 1$. All nonzero points of $\mathbb{Z}P$ are nontorsion. Thus C_S is a nonzero

nontorsion point, and the usual fixed-subtraction formulas have no remaining zero denominator. Explicitly, define

$$N_S = (-660 + 417\rho)h_S + (-228 + 112\rho)j_S,$$
$$D_S = (-704 + 384\rho)h_S + (660 - 417\rho)j_S + (6888 - 5040\rho)g_S.$$

Then $D_S \neq 0$ and $z(C_S) = N_S/D_S$, where $z = -x/y$. These identities are obtained from the chord-and-tangent formulas; they are the same fixed-subtraction identities independently checked in the exact function field by the supplied certificate.

Use the common chart of Lemma 3.1, so $H = (f^{-1}, S_H/f)$ and $H - P = (F_0^{-1}, T_H/F_0)$. Before any logarithm is used, the scalar ratio coordinate $Z = Z_0 + \rho Z_1$ is determined by

$$ZD_S = -N_S S_H.$$

When H is nontorsion this is $Z = z(C_S)/z(H)$. Set $F = \mathrm{Nm}_{K/\mathbb{Q}}(f) \neq 0$. The reduced scalar norm equations are

$$A = \frac{Z_0^2 + R_s Z_1^2}{R_s^4}, \qquad W_s = \frac{A^2 + R_s F^2}{R_s^4}, \qquad b_s = R_s(R_s + 9792)W_s^2. \tag{11}$$

The actual quadratic source stores $R_s^2, R_s^4, Z_1^2, F^2, A, W_s$ and a scaled square of W_s; thus these equations are quadratic after clearing. The normalized scalar pole has exactly the ordinary pole input b_s in (11). To verify its scaling literally, let M be the coefficient of the stored W_s-square coordinate in source row491, let K be the coefficient of $e\bar{b}$ in row499, and let L be the coefficient of $\bar{b}p^2$ in row494. The exact coefficient identities are

$$K = 82479248M, \qquad 3L = 17K.$$

The rational substitutions

$$b_s = -K\bar{b}, \quad e' = -e/K, \quad p' = p/K, \quad u' = u/K, \quad v' = -v, \quad w' = w/K$$

convert those rows to the retained normalized pole equations. The independent checker verifies these identities. Since $R_s > 0$ and $F \neq 0$, one has $W_s > 0$ and $b_s > 0$ at every rational source solution.

Lemma 4.1. *Let $p > 76176$ or $p = 149$, and suppose $\nu = v_p(R_s) > 0$ is odd. Then*

$$v_p(F) > 5\nu/4, \qquad v_p(Z_0) > 23\nu/8, \qquad v_p(Z_1) > 19\nu/8.$$

Once the point interpretations hold, these imply

$$v_p(a - 1) > 19\nu/8.$$

Proof. The number $9792 = 2^6 \cdot 3^2 \cdot 17$ is a unit at p. The valuation of the nonzero pole input is $\nu + 2v_p(W_s)$, so the pole obstruction makes it positive and $v_p(W_s) > -\nu/2$. The two nonzero terms of a norm $u^2 + R_s v^2$ have opposite valuation parity. A zero term can be omitted. Hence the second norm in (11) gives

$$v_p(A) > 7\nu/4, \qquad v_p(F) > 5\nu/4.$$

The first norm gives the two asserted bounds on Z_0, Z_1.

At some place above p the value $f(H)$ has positive valuation, because the valuation of its norm is positive. Thus H is in the formal group there. The displayed bounds give $v(Z) > 19\nu/8 > 0$, so $v(z(C_S)) > 0$. On $y^2 = x^3 - Dx$, at a good odd prime a point with positive z-valuation is either formal or reduces to $(0, 0)$. Indeed $z^2 = x/(x^2 - D)$, and D is a unit. Translation by

$(0,0)$ sends z to $-z$ and exchanges these two cases. Choose $C' = C_S + \epsilon(0,0)$ to be formal. The integral identity

$$[2m]C' = [4(a-1)]H$$

kills both two-torsion terms. The formal logarithm is an isometry, so its ratio has valuation $v_p((a-1)/h) = v(Z)$. Since $h = m/2$ is p-integral, $v_p(a-1) \geq v(Z) > 19\nu/8$. □

The integer $\ell = 149$ is prime. For the five contact roots $0, 1, 2401, -343/32, 1/2$, the residues of $(\theta - 2504)/51$ are respectively $59, 97, 109, 108, 78$ modulo 149. Their 74th powers are all 148, so they are nonsquares. Trial division through 12 certifies primality. The preceding lemma applies at 149: the curve has good reduction there, 9792 is a unit, and the pole obstruction excludes only $2, 3, 17$. The generic contact cutoff is unnecessary at this separately verified prime. Thus r_s is an ℓ-adic unit when X is integral at ℓ. If $v_\ell(X) < 0$, then $v_\ell(r_s) = 10v_\ell(X)$. It follows that

$$v_\ell(R_s) = \begin{cases} 2N+1, & v_\ell(X) \geq 0, \\ 2N-1, & v_\ell(X) < 0. \end{cases} \tag{12}$$

This is always positive and odd. These finite facts are included in the independent exact certificate.

Lemma 4.1, using only its norm inequalities at this stage, gives $v_\ell(F) > 0$. Every finite allowed point in

$$E[2](K) \ \cup \ (P + E[2](K)) \ \cup \ (-P + E[2](K))$$

has reciprocal-abscissa norm an ℓ-adic unit. To check this, use the two-torsion translations and

$$\mathrm{Nm}(x_P) = 41^2/64, \quad \mathrm{Nm}(x_P - (3-\rho)) = 17^2/64, \quad \mathrm{Nm}(x_P + (3-\rho)) = 63^2/64.$$

The remaining torsion abscissas have norm seven. Thus none of these points can equal H. In particular $h \notin \{-1, 0, 1\}$, H and $H - P$ are nontorsion, and all required denominators and formal-logarithm interpretations are now justified. This exclusion precedes every use of a comparison logarithm.

Let $c = |a - 1| > 0$. Outside the fixed cutoff, every prime of $M_5(\beta)$ is a positive odd contact of r_s and R_s. The contact at infinity contributes none, since a negative valuation of $\beta = 51X^2 + 2504$ is even there. Lemma 4.1 therefore implies $M_5(\beta)^3 \mid c$. The prime $\ell = 149$ is below the defining cutoff of M_5, so it is automatically absent from $M_5(\beta)$. The forced-prime estimate is consequently independent of these radical factors. We obtain

$$\ell^{L_N} M_5(\beta)^3 \mid c, \qquad L_N = \left\lfloor \frac{19(2N-1)}{8} \right\rfloor + 1. \tag{13}$$

Together with (1), this gives

$$h(\beta) \leq H \left(\frac{c}{\ell^{L_N}} \right)^{7/9} \leq H \ell^{-L_N} c, \qquad c \geq \ell^{L_N}. \tag{14}$$

The last inequality uses $c/\ell^{L_N} \geq 1$. These scalar estimates are independent of the comparison logarithms.

5 Plain parameter ratios and the local tests

For $f = 1/x$ and $z = -x/y$, extend their regular values at the identity by zero. The retained point relations are

$$U_0 = H, \quad V_0 = H - P, \quad U_i = [u_i]P, \quad V_i = U_i - Q_i \qquad (1 \leq i \leq 4).$$

The charts permit $U_i = O$ or $V_i = O$ for $i > 0$. For each numerator pair retain its quadratic pair-chart coordinates $\mathcal{V}_i, \mathcal{Q}_i, \mathcal{W}_i, \mathcal{I}_i, \mathcal{K}_i$ as specified in [4]. Their exact relations give

$$z(U_i) = -\frac{\mathcal{V}_i}{\mathcal{I}_i}, \qquad z(V_i) = -\frac{\mathcal{Q}_i}{\mathcal{K}_i}, \qquad \mathcal{I}_i\mathcal{K}_i \neq 0,$$

including the identity branches with zero numerator. The common-point chart stores both negative reciprocal parameters $S_H = -1/z(H)$ and $T_H = -1/z(H - P)$. Thus the quadratic equations

$$L_i\mathcal{I}_i = \mathcal{V}_i S_H, \qquad L'_{i,0}\mathcal{K}_i = \mathcal{Q}_i T_H, \qquad L'_i = \sigma L'_{i,0}$$

mean precisely

$$L_i = \frac{z(U_i)}{z(H)}, \qquad L'_i = \sigma\Big(\frac{z(V_i)}{z(H - P)}\Big).$$

No numerator-parameter coordinate or logarithm correction is required. Put

$$D_H = f(H), \quad A_i = f(U_i), \quad B_i = \sigma f(V_i), \quad C_i = L_i - L'_i, \quad I = L_1 - \beta,$$

$$G = L_3^2 - L_2(L_2 - 2)L_3L_4 + (L_2 - 2)^2L_4^2 - 1.$$

The required orders are two for D_H, three for C_2, C_3, C_4, G, one for each of C_1, I, and one for the nine A_i, B_i. The final input comparison uses order one.

6 Four factors and extraction at one place

At a contact prime $p \notin \Sigma$, δ is a unit and $\nu = v_p(R) = v_p(F_2(t_*))$ is positive and odd, and $v_p(q) = -\nu$. This follows from the unique contact factor with positive odd valuation and does not require any condition on the valuation of the scale. All valuations of K are unramified and normalized by $v(p) = 1$. For a point on E, every finite value of $v(f)$ is even: if $v(x) < 0$ then $2v(y) = 3v(x)$, while if $v(x) > 0$ then $2v(y) = v(x)$; the remaining case is valuation zero. This also holds for conjugated reciprocal abscissas. The common-point lemma identifies $F_0 = f(H - P)$ on every solution, including the exceptional leading-coefficient cases. Its even valuation is therefore available before the exclusion of torsion classes and before any formal-parameter quotient is used.

Use

$$\begin{aligned}
T_1 &= 1 + q^9C_2^4 + q^3C_1^4 + q^2D_H^2 + q^4B_1^4,\\
T_2 &= 1 + q^9G^4 + q^3A_3^2 + q^2B_0^2 + q^4B_2^4,\\
T_3 &= 1 + q^9C_3^4 + q^3A_2^2 + q^2I^4 + q^4B_3^4,\\
T_4 &= 1 + q^9C_4^4 + q^3A_1^2 + q^2A_4^2 + q^4B_4^4, \qquad W = T_1T_2T_3T_4.
\end{aligned}$$

Lemma 6.1. *At a contact prime, every factor is nonzero and has valuation at most zero. A factor of valuation zero is congruent to one in the residue field.*

Proof. First suppose every displayed test in a factor is nonzero. The first nonconstant term in every factor has valuation $4v(C) - 9\nu$, where C is the indicated deep comparison. Its residue modulo four is $-\nu$. The second term has either valuation $4v(C_1) - 3\nu$ or $2v(A_i) - 3\nu$. The even valuation of every reciprocal abscissa therefore makes its residue ν modulo four. The third term has either valuation $2v(f) - 2\nu$, with $v(f)$ even, or $4v(I) - 2\nu$. Its residue modulo four is two. Finally the fourth term has valuation $4v(B_i) - 4\nu$, which has residue four modulo eight because $v(B_i)$ is even.

The classes $-\nu, \nu, 2$ modulo four are pairwise distinct and nonzero because ν is odd. The fourth class differs from them and from the constant term's valuation zero. Thus the finite term

valuations are pairwise distinct. Zero tests simply remove terms. The minimum is attained exactly once, which proves nonvanishing and valuation at most zero. If the minimum is zero, all nonconstant terms have positive valuation, so the residue is one. □

Proposition 6.2. *At each contact prime, the retained pole system and the cleared trace equations imply $\nu > 0$ and give all sixteen required test orders at one place above that prime.*

Proof. Write $W = w_0 + w_1\rho$. The literal cleared trace equations are

$$w_0^2 - 2w_0\tau - 2w_1^2 = 0, \qquad (1+q)\mathfrak{b} = \gamma^2\tau^2.$$

By Lemma 6.1, $W \neq 0$. The first equation therefore gives $w_0 \neq 0$ and $\tau = 1/\operatorname{Tr}(W^{-1}) \neq 0$; indeed $w_0 = 0$ would give $w_1 = 0$. Hence $\mathfrak{b} = \gamma^2 R/\operatorname{Tr}(W^{-1})^2$. The factor γ is a unit at every consumed contact prime. If $\ell = \max_{v|p} v(W) \leq 0$, the trace inequality gives $v_p(\mathfrak{b}) \leq \nu + 2\ell$. This valuation is odd; the retained pole obstruction forces it positive. Consequently $\nu > 0$ and $\ell > -\nu/2$. At a place attaining ℓ, the factor valuations are nonpositive and add to ℓ. Each factor valuation is therefore at least ℓ, and each of its term valuations is strictly greater than $-\nu/2$.

The resulting lower bounds, before integer rounding, are

test	strict lower bound	integer order
C_2, C_3, C_4, G	$17\nu/8$	3
C_1	$5\nu/8$	1
I	$3\nu/8$	1
D_H	$3\nu/4$	2
A_1, A_2, A_3	$5\nu/4$	1
A_4, B_0	$3\nu/4$	1
B_1, B_2, B_3, B_4	$7\nu/8$	1

Here ν is a positive odd integer. The order two for D_H uses its even valuation. In fact each of the nine reciprocal-abscissa tests has order at least two; order one suffices for its subsequent formal-group use. Zero tests satisfy all the stated inequalities. □

7 Integrality before the formal congruences

At a contact place selected by Proposition 6.2, the points H, U_i are formal in the primary completion, and $H - P, V_i$ are formal in the conjugate completion. The reciprocal-abscissa tests have positive even valuation, so every nonidentity point has formal depth at least one. In the parameter $z = -x/y$, write

$$\log_E(Q) = z(Q)\mathcal{A}(Q), \qquad \mathcal{A}(Q) \equiv 1 \pmod{p^{4v(z(Q))}}.$$

The value of the unit is taken to be one at the identity. This is the constant-term formal logarithm estimate proved in [9].

The integral group identities

$$[m]U_i = [2u_i]H, \qquad [m-2]V_i = [2(u_i - n_i)](H - P)$$

kill the torsion in $H = [m]R_0 + T_0$. Their full logarithmic ratios are u_i/h and $(u_i - n_i)/(h-1)$, where $(n_1, n_2, n_3, n_4) = (n, a, t, b)$. At least one of $h, h-1$ is a p-adic unit, so at least one full ratio is integral. The formal units preserve the valuation of its plain parameter ratio. The positive-order test C_i transfers integrality to the other plain ratio, and then to the other full ratio.

Only after this transfer may the relative errors of order four be used as absolute congruences. They give

$$L_i \equiv u_i/h \pmod{p^4}, \qquad L_i' \equiv (u_i - n_i)/(h-1) \pmod{p^4}.$$

Since $n_i = h(u_i/h) - (h-1)((u_i - n_i)/(h-1))$, the high cross tests give $L_i \equiv n_i \pmod{p^3}$ for $i > 1$. The first cross test gives $L_1 \equiv n \pmod{p}$, while I gives $L_1 \equiv \beta \pmod{p}$. These arguments include identity numerator points, half-integral h, and inert primes, where conjugation preserves the unique valuation.

8 Soundness at the explicit parameter

Let $N \geq 14521$. Put $L = L_N$ and $c_0 = 149^L$; then $c \geq c_0$ and $B \geq c - 1$. At $N = 14521$ one has $L = 68973$. Exact integer comparison establishes

$$\begin{aligned} (149^{68973} - 1)^4 &> (14 \cdot 2^{221284})^9, \\ 149^{68973} &> 40 \cdot 2^{221234}, \\ 149^{68973} &> 40 \cdot 265546. \end{aligned} \tag{15}$$

These inequalities remain true at every larger N. In particular $c > 8$ and (14) gives $h(\beta) < c/40$.

Put $\Delta = t^2 - a(a-2)tb + (a-2)^2b^2 - 1$. The local comparison proves that each M_j^3 divides Δ. If $\Delta \neq 0$, then $|\Delta| \leq 14B^4$ and (6), with $k = 3$, gives

$$B^2 \leq 5H_2(14B^4)^{7/18} < 14H_2B^{14/9}.$$

The last strict inequality follows from $5^{18} < 14^{11}$. It implies $B < (14H_2)^{9/4}$, contradicting $B \geq c - 1$ and (15). Thus $\Delta = 0$. A stronger form of the integral Lucas descent gives

$$\log B \geq (c-5)\log(c-2) \qquad (c \geq 8). \tag{16}$$

Here is the complete elementary argument. Put $A_0 = |a|$ and $w = \operatorname{sgn}(a)(a-2)b$. Then $t^2 - A_0tw + w^2 = 1$. For $\eta = (A_0 + \sqrt{A_0^2 - 4})/2$, every positive norm-one unit of $\mathbb{Z}[\eta]$ is a power of η. Indeed multiply a unit by a power of η to place it in $[1, \eta)$; its coefficient of η is $(v - v^{-1})/(\eta - \eta^{-1})$, an integer in $[0, 1)$, so it is zero and $v = 1$. Let $U_0 = 0$, $U_1 = 1$, and $U_{j+1} = A_0U_j - U_{j-1}$ for $j \geq 1$. The unit identities show that $|w| = U_j$ and $|t| \in \{U_{j-1}, U_{j+1}\}$ for some $j \geq 1$. Modulo $A_0 - 2$ one has $U_j \equiv j$; modulo $A_0 + 2$ one has $U_j \equiv (-1)^{j-1}j$. Since w is a nonzero multiple of $|a-2|$, it follows that $j \geq A_0 - 2$ if $a > 0$ and $j \geq A_0 + 2$ if $a < 0$. Induction gives $U_j \geq (A_0 - 1)^{j-1}$ for $j \geq 1$. Therefore $|t| \geq (A_0 - 1)^{A_0 - 4}$. Finally $A_0 \geq c - 1$, which proves (16).

If $n - \beta \neq 0$, let J be its nonzero reduced numerator. The local input comparisons make all three radicals divide J, while $|J| \leq 2B\exp(h(\beta))$. Equation (6) with $k = 1$ yields

$$\log B \leq \frac{6}{5}\log(5H_2) + \frac{7}{5}\log 2 + \frac{7}{5}h(\beta) < 265546 + \frac{7}{5}h(\beta) < \frac{3c}{50}.$$

Here $\log 2 < 1$, $\log 5 < 3\log 2$, and (15) justify every numerical bound. For $c \geq 8$, the Lucas bound (16) is at least $c - 4 \geq c/2$, a contradiction. Hence $n = \beta \in \mathbb{Z}$. If $X = u/v$ is reduced with $v > 0$, integrality of $51X^2 + 2504$ implies $v^2 \mid 51$; squarefreeness of 51 gives $v = 1$. This proves soundness for every $N \geq 14521$.

9 Simultaneous completeness

Fix an integer input and $N \geq 1$. Choose a real neighborhood of a nonzero reference point $[h_*]P$, with $64 \mid h_*$, on which both values of $|z(H)|$ are bounded below. Then choose a sufficiently

small real neighborhood of the identity for C_S. It gives $|Z_0|, |Z_1| \le 1$. For nonzero $H \in \mathbb{Z}P$ the positive-square abscissa bound also gives $|\operatorname{Nm}(f(H))| \le 1/7$. For $R_s \ge 1$, equations (11) imply

$$A \le 2R_s^{-3}, \qquad W_s \le 4R_s^{-10} + \frac{1}{49}R_s^{-3} < 5R_s^{-3}.$$

Therefore

$$0 < b_s \le 25\frac{R_s + 9792}{R_s^5} < 17 \qquad (R_s \ge 10).$$

The displayed rational function is decreasing on $(0, \infty)$, and its value at ten is less than 17. Since $r_s \ge 1$,

$$R_s \ge \frac{149^{2N+1}}{1 + 149^2} \ge \frac{149^3}{1 + 149^2} > 10.$$

Thus the required real preparation holds for every $N \ge 1$ with explicit margins.

The finite preparation uses no new arithmetic theorem. For integer input, r_s is a positive integer and

$$R_s(R_s + 9792) = \frac{A_N r_s(A_N r_s + 9792(1 + \ell^2 r_s))}{(1 + \ell^2 r_s)^2}.$$

The numerator is coprime to the displayed denominator, so its reduced denominator is square. At the three preparation primes the elementary input congruences give

$$v_2(R_s) \in \{2, 3\}, \qquad v_3(R_s) = 1, \qquad v_{17}(R_s) = 0.$$

In particular $r_s \equiv 15 \pmod{17}$ and $1 + 149^2 r_s \equiv 3 \pmod{17}$, so the normalizer denominator is also a unit at 17. Thus $R_s(R_s + 9792) = R_s^2(1 + 9792/R_s)$ is a local square at all three primes. Choose the shared index h_0 sufficiently divisible at the finite support of this rational number, and then choose the index $c = a - 1$ still more divisible. The points $H = [h_0]P$ and $C_S = [c]P$ are then formal at the required finite places, and all components of $f(H)$ and Z have the needed arbitrarily large valuations. Make W_s integral at every prime where the displayed numerator has odd valuation, and deep enough at $2, 3, 17$ to supply the pole bounds. At every other prime the valuation of b_s is even. These conditions give a square reduced denominator and all pole preparation hypotheses. The source's progression and compact-real-group argument supplies the chosen real neighborhoods at the same time, together with the retained nonzero chart guards. It also permits c to be arbitrarily large and to satisfy the retained divisibility conditions for the Lucas construction.

After S is fixed, continuity preserves this finite scalar preparation and its guards for all h in one fixed congruence class modulo a sufficiently divisible integer L. Retain the source's integral Lucas construction, choose the four root labels, and fix the three positive scales and compact neighborhoods for all comparison tests before selecting contact primes. The additional contact primes avoid the finite scalar preparation support. Hence these later congruences do not conflict with any scalar congruence.

Choose a fixed positive integer D_0 clearing all contact coefficients $D_0^2\theta/a_*$ and sufficiently divisible at the prepared primes, including Σ, and the prime divisors of the scales. In the prime pattern write

$$\lambda = \frac{u}{D_0^2 w^2}.$$

The zero contact makes u one of the selected primes, outside that finite set. Thus $v_p(\lambda) \le -2v_p(D_0)$ there. Make this upper bound low enough that $t_* = a_*\lambda$ has valuation below zero and all nonzero contacts. Then $v_p(\delta) = v_p(t_*)$, and

$$R = \prod_{\theta \in \Theta_2} \left(1 + \frac{1 - \theta}{\delta}\right) \longrightarrow 1 \tag{17}$$

in every prepared completion. The weight $q = R^{-1} - 1$ tends to zero. Increasing the fixed denominator divisibility makes its valuation uniformly arbitrarily large. Thus all weighted test terms are small at every prepared place, and $\mathrm{Tr}(W^{-1})$ is arbitrarily close to two.

The factors in (17) are arbitrarily close to one. Requiring congruence to one modulo eight at 2, and modulo p at an odd prepared prime p, makes their product a local square. The pole input is $\gamma^2 R/\,\mathrm{Tr}(W^{-1})^2$, so it is also a square at $2, 3, 17$. The fixed factor has valuations

$$v_2(\gamma) = 3, \qquad v_3(\gamma) = v_{17}(\gamma) = 1.$$

Since the reciprocal trace is close to $1/2$, the pole input has valuations $4, 2, 2$ at those three primes. These satisfy all retained preparation inequalities.

This preparation is valid for any fixed nonzero rational scales. The prime-pattern input in [10, 1] applies to arbitrary fixed distinct integer coefficients. After clearing $D_0^2\theta/a_*$ and removing repeated coefficients, it applies here without a square restriction on a_*. All denominator conditions are fixed before prime selection.

Select the contact primes with the inherited opposite-order separation. Take $H = [h]P$, $U_i = [hn_i]P$ and $V_i = [(h-1)n_i]P$. The common-index congruences preserve the scalar progression and make H, U_i formal at one place and $H - P, V_i$ formal at its conjugate. Only natural depth one is needed on either branch; the source's stronger primary preparation is also permitted. Both full logarithmic ratios are exactly n_i. Plain-parameter errors therefore have order at least four. All C_i, and I since $n_1 = \beta$, have valuation at least four, and the Lucas equation for the chosen labels gives $v(G) \geq 4$. Every reciprocal-abscissa test has valuation at least two. At the selected place $v_p(q) = -1$, so every nonconstant term of every packed factor has positive valuation. Thus $W \equiv 1$ there. At the opposite place, each factor has valuation at most zero, with residue one when its valuation is zero. The trace of W^{-1} is a unit with residue one or two. The comparison pole input has valuation one at every selected contact prime.

The exact prime-pattern identity is

$$R = \frac{a_*^{10} \prod_{\theta \in \Theta_2} (u - (\theta/a_*) D_0^2 w^2)}{(D_0^2 w^2)^{10} \delta^{10}}.$$

Every factor outside the selected prime numerators is a rational square, including a_*^{10} even when a_* is not itself a square. Thus at every prime outside those numerators the valuation is even. Multiplication by γ^2 and division by the squared trace preserve this parity. This argument includes primes dividing δ; no unit assertion about δ is made away from contacts. Equation (17) and the preparatory divisibility give the required local squareclasses and valuation bounds at $2, 3, 17$.

For the real preparation, write $r = \delta^2$ and let $\alpha_1, \ldots, \alpha_5 > 0$ be the five squared distances of the contact pairs from one. Then

$$R = \prod_{j=1}^{5} (1 - \alpha_j/r), \qquad q \sim \frac{\alpha_1 + \cdots + \alpha_5}{r} > 0 \quad \text{as } r \longrightarrow +\infty.$$

On the fixed compact real neighborhoods the tests remain bounded, so $W \to 1$ and $\mathrm{Tr}(W^{-1}) \to 2$. Consequently

$$\mathfrak{b} \longrightarrow \frac{\gamma^2}{4} = \frac{41616}{167281} \in (0, 17).$$

All factor values are positive at both real embeddings for sufficiently large λ, because $q > 0$ and all test powers are even. The simultaneous selection supplies the required real size after all fixed data have been chosen. The pole-realization input supplies its witnesses, and the exact power identities fill the quadratic circuit. This proves completeness for every integer input and every $N \geq 1$.

10 Exact quadratic specification

Retain three scalar inverse weights s_1, s_2, s_4 and a fifth-power weight w_5 for each detector, with the quadratic equations

$$qs_1 = 1, \qquad s_2 = s_1^2, \qquad s_4 = s_2^2, \qquad w_5 s_4 = q. \tag{18}$$

They give $q \neq 0$, $s_j = q^{-j}$ for $j = 1, 2, 4$, and $w_5 = q^5$. The factor coordinates satisfy the quadratic field equations

$$\begin{aligned} s_4 T_1 &= s_4 + w_5 C_2^4 + s_1 C_1^4 + s_2 D_H^2 + (B_1^2)^2, \\ s_4 T_2 &= s_4 + w_5 G^4 + s_1 A_3^2 + s_2 B_0^2 + (B_2^2)^2, \\ s_4 T_3 &= s_4 + w_5 C_3^4 + s_1 A_2^2 + s_2 I^4 + (B_3^2)^2, \\ s_4 T_4 &= s_4 + w_5 C_4^4 + s_1 A_1^2 + s_2 A_4^2 + (B_4^2)^2. \end{aligned} \tag{19}$$

Each displayed comparison square or fourth power is a stored field value. Squaring a stored B_i^2 is quadratic in its two rational components. Multiplication by q^4 proves exact equivalence with the four factor definitions above. There is no additional zero stratum: $s_4 \neq 0$ follows from $qs_1 = 1$, and all fourth powers are uniquely determined by their quadratic definitions. Two intermediate field products and the final field value W compute $T_1 T_2 T_3 T_4$.

The four contact outputs are

$$r = (t_* - 1)^2, \qquad r_2 = r^2, \qquad r_3 = r r_2, \qquad F_2(t_*) = P_5(r) = \prod_{\theta \in \Theta} (r - (\theta - 1)^2).$$

Every integer from zero through five is a sum of two elements of $\{0, 1, 2, 3\}$, so the last expression is quadratic in the retained outputs. Its normalization is $F_2(t_*)(1 + q) = r_2 r_3$. The scalar normalizer is the quadratic equation $R_s + 22201 r_s R_s - A_N r_s = 0$. The scalar chain is $R_2 = R_s^2$, $R_4 = R_2^2$. The two scalar norm rows use R_4, as in (11).

The six field fourth powers $C_1^4, C_2^4, C_3^4, C_4^4, I^4, G^4$ are stored once and shared by all three detectors. The four required B_i^4 values occur directly as squares of stored field coordinates in (19). All remaining point and scalar equations are fixed by the literal coefficient tables. The two nontrivial scale coordinates are quadratic outputs. In the stable identifiers of the coefficient table,

$$S_0 = x_{82} + x_{84} + 2x_{86} + x_{88}, \qquad S_1 = 2(x_{83} + x_{85} + x_{87} + x_{89}).$$

The factors two account for the source's half-component convention; they are verified directly from the four square-abscissa equations. The third detector has the same equations as the first, with its argument replaced by $(S_0 + S_1)^2 \lambda$. The exact count is

point, scalar, contact, ratio, and weight coordinates	217
six shared fourth powers	12
four factors and intermediate products for three detectors	36
total coordinates, including the input	265
point, scalar, contact, ratio, and weight quadratic rows	183
shared fourth-power rows	12
factor and product rows	42
total quadratic rows	237

The complete tables fix every coordinate, equation, coefficient, and normalization; the input is coordinate one. For the common point, the six primitive rational component equations are replaced by the fixed rational row combinations in the certificate, whose determinant

is $-1/362963666515968$. The producer clears denominators and makes each resulting row primitive before squaring. The row change is nonsingular and preserves the common zero set. The independent checkers import no producer. The common-point certificate checks twelve general identities, including the master elimination identity without division, the curve equations and the translation law on every exceptional stratum. The coefficient checker reconstructs all six common-point rows, the direct ratio equations, and the full sum of squares, and compares every remaining row with its pinned coefficient source. The contact certificate recovers P_5 by Newton finite differences at six exact arguments, checks all five roots, and independently verifies $P_5((t-1)^2) = F_2(t)$ as a degree-ten polynomial identity. It reconstructs every modified quadratic equation, compares the other equations with their pinned coefficient source, and checks the contact exceptional set and both height matrices. It also verifies the scalar pole scaling, the forced prime and its nonsquare tests, and the scalar extraction bounds. The contact normalization and its remaining $t_* = 1$ stratum are verified directly. The current coefficient checker checks all reciprocal-weight equations and reconstructs every cleared factor from its field formula. It compares 196 unmodified rows with their pinned source, reconstructs thirty changed factor and product rows, eight new shared fourth-power rows, and three fifth-power weight rows, and checks the entire expanded sum of squares. The residue certificate verifies the distinct valuation classes modulo eight and the rational extraction bounds for all integer valuations. The projection proof uses these literal equations and the preceding arithmetic argument. It expands the entire sum of squares independently by unordered pairs of terms. The supplementary local certificate exhausts all odd residues of ν modulo eight and all possible valuation residues of every test. These complete residue classes verify the noncancellation assertion for all integer valuations, together with the exact extraction bounds and sufficient completeness depths. The infinite arithmetic arguments are the proofs above.

The expanded quartic has 3486 terms before powers of A_N belonging to the same coordinate monomial are collected, and 3485 monomials after collection. Its unique coefficient involving A_N^2 is $A_N^2 + 1$, at the square of the actual scalar contact-product coordinate. Every other coefficient is a nonzero constant or a nonzero constant multiple of A_N. Consequently no support cancellation occurs for positive A_N. The largest constant coefficient C_0 has 1520 bits, the largest linear multiplier is 44402, and exact comparison gives

$$C_0 > 44402^2, \qquad 149^{210} + 1 < C_0 < 149^{214} + 1.$$

Thus $H(P_N) = \max(C_0, A_N^2 + 1)$ for every $N \geq 1$ and the asserted exact height holds for $N \geq 53$. The distinct sufficient defining threshold $N \geq 14521$ follows from the arithmetic proof above. At that parameter the exact height is $149^{58086} + 1$, with 419333 binary digits. The producer `build_quartic265_v19.py` and independent checker `check_quartic265_v19.py` reproduce all these assertions.

References

[1] Project repository, *Arithmetic dependencies of the defining-polynomial constructions*, `arithmetic_dependencies_20261108_v8.tex`.

[2] Project repository, *A family of quartics in 297 variables*, `quartic297_20261109_v12.tex`, especially Sections 2–6.

[3] Project repository, *Common local weights and quartics in 297 and 342 variables*, `shared_local_weights_20261109_v12.tex`.

[4] Project repository, *An explicit quartic in 322 variables*, `quartic322_20261109_v13.tex`.

[5] Project repository, *Height Bounds from Disjoint Sets of Rational Contacts*, `contact_height_amplification_20261109_v13.tex`.

[6] Project repository, *Effective heights of sums of elliptic abscissas*, `effective_elliptic_sums_20261108_v10.tex`.

[7] Project repository, *Evaluation of Differentials and Heights of Modular Jacobians*, `evaluation_of_differentials_20261109_v17.tex`, Definition 6.1 and Theorem 6.2.

[8] Project repository, *Multiplicity of modular factors and quadratic contact-height bounds*, `multiplicity_and_contact_heights_20261110_v19.tex`, Theorem 4.1.

[9] Project repository, *Uniform Rational Approximations to an Elliptic Logarithm*, `elliptic_logarithms_20261108_v8.tex`.

[10] Project repository, *Prime selection with fixed congruences and separated reduction orders*, `prime_selection_20261109_v12.tex`.