

## BACKGROUND

10-30% of non-hospitalized cases of SARS-CoV-2 infection are estimated to suffer from some Long Covid symptoms [1,2].

Up to 50% of individuals with other Long Covid symptoms reported also having problems with memory, cognition or concentration [3].

There are mixed results in regards to the protective effect of vaccination against Long Covid symptoms [4–6].

## INTRODUCTION

Those who have experienced SARS-CoV-2 were found to have:

- Lower scores on cognitive tasks, particularly those requiring reasoning, planning and problem solving [7].
- Lower scores in reasoning and verbal domains but not in the short-term memory [8].
- Lower performance on memory tasks, verbal item memory task in particular [9]. This is in contrast to usual pattern of memory deterioration found in aging [10].

The current study roughly follows the design of Guo et al., (2022) [9], allowing us to replicate their previous results with two important additions:

- (1) Inclusion of information regarding vaccination status.
- (2) Inclusion of additional memory tasks, allowing us to disentangle stimulus effects (pictorial vs. verbal) from processing effect (item vs. associative retrieval).

### Demographics

Age  
Sex  
Education  
Country  
Ethnicity  
Occupation

### Health

Height  
Weight  
Medical conditions  
Lifestyle

### Covid


Covid status  
Vaccination status  
Vaccination details  
Covid symptoms  
Long Covid symptoms  
Timing

### Cognitive tasks

Nonverbal associative memory task  
Verbal associative memory task  
Nonverbal item memory task  
Verbal item memory task  
Digit span memory task  
Category Fluency task  
Word/Syntax Understanding task  
Wisconsin Card Sorting Task

## METHODS

- Data collection ongoing since February 2022.
- Data collection on Gorilla.sc consisting of a series of questionnaires and cognitive tasks.
- Participants recruitment mainly through social media, Long Covid support groups and word of mouth and to lesser extent through Prolific.co and Addenbrooke's hospital Long Covid clinic.



## GUO ET AL. (2022) [9]

- Significant negative influence of Covid-19 status on memory performance.
- Larger trend for the verbal item recognition test and remained even after controlling for age, sex, country, education level.
- Smaller trend in the same direction was found for the nonverbal associative memory test.
- No significant difference between the groups in Executive Functions performance.

## ANALYSIS

Analysis	Predictions	Between subjects	Within subjects	Dependent measures
Analysis 1 and 2: Replication of Guo et al. (2022)	Worse accuracy and slower RTs for C+	C+ vs.. C-		Verbal item memory accuracy / RTs
		C+ vs. C-		Nonverbal associative memory accuracy / RTs
Analysis 3 and 4: Comparison of mnemonic effect	Worse accuracy and slower RTs for C+ group in verbal item than nonverbal associative task	C+ vs. C-	Verbal item vs. nonverbal associative	Accuracy/RTs
Analysis 5: Disentangling memory effects	Worse memory in C+, possible effect of memory type or stimulus type or both	C+ vs. C-	Item vs. associative verbal vs. nonverbal	Accuracy/RTs
Analysis 6: Effect of vaccination status	Better accuracy and RTs for C+ and vaccinated than C+ and non-vaccinated, no difference between C- and vaccinated x C- and nonvaccinated	C+ vs. C-		Accuracy and RTs for all cognitive tasks separately
		Vaccinated vs. Nonvaccinated		

## OUTSTANDING QUESTIONS

- Is it necessary to correct for multiple comparisons when the comparisons are preregistered? What would be a good method of correcting?
- How to deal with potential outliers in the data and how to preregister it?
- What other analyses may be interesting?

## REFERENCES

[1] Bull-Otterston L, Baca S, Saydah S, et al. Post-COVID Conditions Among Adult COVID-19 Survivors Aged 18–64 and ≥65 Years – United States, March 2020–November 2021. *Morb Mortal Wkly Rep* 2022; 71: 713–717.

[2] Ceban F, Ling S, Lui LMW, et al. Fatigue and cognitive impairment in Post-COVID-19 Syndrome: A systematic review and meta-analysis. *Brain Behav Immun* 2022; 101: 93–135.

[3] Dennis A, Cuthbertson DJ, Wootton D, et al. Multi-organ impairment and long COVID: a 1-year prospective, longitudinal cohort study. *J R Soc Med* 2023; 01410768231154703.

[4] Mizrahi B, Sudry T, Flaks-Manov N, et al. Long covid outcomes at one year after mild SARS-CoV-2 infection: nationwide cohort study. *BMJ* 2023; e072529.

[5] Al-Aly Z, Bowe B, Xie Y. Long COVID after breakthrough SARS-CoV-2 infection. *Nat Med* 2022; 28: 1461–1467.

[6] Ayoubkhani D, Bosworth ML, King S, et al. Risk of Long Covid in people infected with SARS-CoV-2 after two doses of a COVID-19 vaccine: community-based, matched cohort study. 2022; 2022.02.23.22271388.

[7] Hampshire A, Trender W, Chamberlain SR, et al. Cognitive deficits in people who have recovered from COVID-19. *EclinicalMedicine* 2021; 39: 101044.

[8] Wild CJ, Norton L, Menon DK, et al. Disentangling the cognitive, physical, and mental health sequelae of COVID-19. *Cell Rep Med* 2022; 3: 100750.

[9] Guo P, Benito Ballesteros A, Yeung SP, et al. COVCOG 2: Cognitive and Memory Deficits in Long COVID: A Second Publication From the COVID and Cognition Study. *Front Aging Neurosci*; 14. Epub ahead of print 2022. DOI: 10.3389/fnagi.2022.804937.

[10] Cohn M, Emrich S, Moscovitch M. Age-Related Deficits in Associative Memory: The Influence of Impaired Strategic Retrieval. *Psychol Aging* 2008; 23: 93–103.