

From Λ CDM to EDE

Lecture 1: Theory

Laura Herold

CosmoVerse@Corfu May 13 - 18, 2024

Overview

Today:

- **Lecture 1:** Theory from Λ CDM to EDE (1:30h)
- **Hands-on session 1:** From theory to predictions (1h)

Tomorrow:

- **Lecture 2:** Observation – Can EDE solve the Hubble tension? (1h)
- **Hands-on session 2:** Let's analyse EDE with cosmic microwave background and supernovae (1:30h)

Outline: From Λ CDM to EDE

1. Introduction: basic equations in a homogeneous & isotropic universe
(Friedmann equations, matter content in the universe, defining distances)
2. Hubble tension: How to solve the tension? How does CMB constrain H_0 ?
3. Early Dark Energy: Idea behind EDE, background dynamics of EDE

Outline: From Λ CDM to EDE

1. Introduction: basic equations in a homogeneous & isotropic universe
(Friedmann equations, matter content in the universe, defining distances)
2. Hubble tension: How to solve the tension? How does CMB constrain H_0 ?
3. Early Dark Energy: Idea behind EDE, background dynamics of EDE




Please ask questions any time!

Introduction

Short “crash course” to fix the notation

Natural units: $c = 1$

Resources



General Relativity

- Let's imagine it was 100 years ago:
 - We didn't know about dark matter (DM)
 - We didn't know about dark energy (DE)
 - But a few years ago, Albert Einstein had published the theory of General Relativity (GR)

1916.

Nº 7.

ANNALEN DER PHYSIK.
VIERTE FOLGE. BAND 49.

1. *Die Grundlage
der allgemeinen Relativitätstheorie;
von A. Einstein.*

Die im nachfolgenden dargelegte Theorie bildet die denkbar weitgehendste Verallgemeinerung der heute allgemein als „Relativitätstheorie“ bezeichneten Theorie; die letztere nennt

General Relativity

- We will not go into details here but only sketch the rough idea

For more about
GR, see Matteo
Martinelli's lecture

General Relativity

For more about GR, see Matteo Martinelli's lecture

- We will not go into details here but only sketch the rough idea
- Einstein Equations:

$$R^{\mu\nu} - \frac{R}{2}g^{\mu\nu} + \Lambda g^{\mu\nu} = 8\pi G T^{\mu\nu},$$

Ricci curvature tensor metric cosmological constant energy-momentum tensor

The diagram shows the Einstein field equations: $R^{\mu\nu} - \frac{R}{2}g^{\mu\nu} + \Lambda g^{\mu\nu} = 8\pi G T^{\mu\nu}$. Four green arrows point to the terms from left to right: a horizontal arrow points to the Ricci curvature tensor $R^{\mu\nu}$; another horizontal arrow points to the metric tensor $g^{\mu\nu}$; a vertical arrow points to the cosmological constant $\Lambda g^{\mu\nu}$; and a diagonal arrow points to the energy-momentum tensor $8\pi G T^{\mu\nu}$.

“Matter tells space how to curve, space tells matter how to move”
(Misner++ 1973)

General Relativity

- Solution for Einstein Equations for the universe as a whole?
- Cosmological Principle:

“On sufficiently large scales, the properties of the universe are the same for all observers.”

or

*“The universe is spatially **homogeneous and isotropic** on large scales.”*


General Relativity

- Solution for Einstein Equations for the universe as a whole?
- Cosmological Principle:


“On sufficiently large scales, the properties of the universe are the same for all observers.”

or

*“The universe is spatially **homogeneous** and **isotropic** on large scales.”*





“the same in every point”




“the same in every direction”


Isotropic or homogeneous?



Isotropic or homogeneous?




homogenous but not isotropic




isotropic but not homogenous

Isotropic or homogeneous?



homogenous but not isotropic



isotropic but not homogenous

Friedmann Equations

- Friedmann 1922, Robertson 1935, Walker 1937: For a spatially homogeneous and isotropic universe, the metric simplifies to

$$ds^2 = -dt^2 + a^2(t) \left[\frac{dr^2}{1-kr^2} + r^2 d\Omega^2 \right]$$

spacetime line element


Friedmann Equations

- Friedmann 1922, Robertson 1935, Walker 1937: For a spatially homogeneous and isotropic universe, the metric simplifies to

$$ds^2 = -dt^2 + a^2(t) \left[\frac{dr^2}{1-kr^2} + r^2 d\Omega^2 \right]$$

spacetime line element

scale factor



$$a(t_{\text{today}}) = a_0 = 1$$

Figure credit: Bianchi, Rovelli, Kolb

Friedmann Equations

- Friedmann 1922, Robertson 1935, Walker 1937: For a spatially homogeneous and isotropic universe, the metric simplifies to

$$ds^2 = -dt^2 + a^2(t) \left[\frac{dr^2}{1 - kr^2} + r^2 d\Omega^2 \right]$$

spacetime line element scale factor curvature parameter

$$k = \begin{cases} -1 & \text{hyperbolic} \\ 0 & \text{flat} \\ +1 & \text{spherical} \end{cases}$$

Friedmann Equations

- A perfect fluid is a fluid, which can be completely characterised by its (energy) density and pressure
- The energy momentum tensor of a perfect fluid is

$$T_{\mu\nu}^{\text{pf}} = (\rho + p) u_\mu u_\nu + p g_{\mu\nu}$$


energy density pressure

- $u_\mu = \dot{x}_\mu$ is the 4-velocity of the observer
- Inserting this in the Einstein Equations, yields the Friedmann Equations

Friedmann Equations

- Inserting the FLRW-metric and the energy momentum tensor of the perfect fluid into the Einstein equations, yields the **Friedmann equations**:

$$\left(\frac{\dot{a}}{a}\right)^2 + \frac{k}{a^2} = \frac{8\pi G}{3}\rho + \frac{\Lambda}{3} \quad (\text{i})$$

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3}(\rho + 3p) + \frac{\Lambda}{3} \quad (\text{ii})$$

Friedmann Equations

- Inserting the FLRW-metric and the energy momentum tensor of the perfect fluid into the Einstein equations, yields the **Friedmann equations**:

$$\left(\frac{\dot{a}}{a}\right)^2 + \frac{k}{a^2} = \frac{8\pi G}{3}\rho + \frac{\Lambda}{3} \quad (\text{i})$$

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3}(\rho + 3p) + \frac{\Lambda}{3} \quad (\text{ii})$$

- Inserting (ii) into the temporal derivative of (i), yields the **continuity equation**:

$$\dot{\rho} = -3(\rho + p)\frac{\dot{a}}{a} \quad (\text{iii})$$

Friedmann Equations

$$\left(\frac{\dot{a}}{a}\right)^2 + \frac{k}{a^2} = \frac{8\pi G}{3}\rho + \frac{\Lambda}{3} \quad (\text{i})$$

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3}(\rho + 3p) + \frac{\Lambda}{3} \quad (\text{ii})$$

$$\dot{\rho} = -3(\rho + p)\frac{\dot{a}}{a} \quad (\text{iii})$$

- Where $\rho = \rho_{\text{tot}}$ is the energy density of the universe.

- If we define $\rho_k = -\frac{3}{8\pi G}\frac{k}{a^2}$ and $\rho_\Lambda = \frac{\Lambda}{8\pi G}$, one can rewrite (i) as

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3}\rho_{\text{tot}} \quad (\text{i})$$

$$\Lambda\text{CDM model} \quad \rho_{\text{tot}} = \rho_r + \rho_m + \rho_\Lambda + \rho_k$$

Λ CDM model

$$\rho_{\text{tot}} = \rho_r + \rho_m + \rho_\Lambda + \cancel{\rho_k}$$

Λ CDM model

$$\rho_{\text{tot}} = \rho_r + \rho_m + \rho_\Lambda$$




Figure credit: Florian Wolz

Λ CDM model

- Equation of state:

$$\rho_{\text{tot}} = \rho_r + \rho_m + \rho_\Lambda$$

$$p = w \cdot \rho$$




Figure credit: Florian Wolz

Λ CDM model

$$\rho_{\text{tot}} = \rho_r + \rho_m + \rho_\Lambda$$

- Equation of state: $p = w \cdot \rho$
- EOS-parameter w for different matter species:
 - Dark matter and baryonic matter: $w = 0$




Figure credit: Florian Wolz

Λ CDM model

$$\rho_{\text{tot}} = \rho_r + \rho_m + \rho_\Lambda$$

- Equation of state: $p = w \cdot \rho$
- EOS-parameter w for different matter species:
 - Dark matter and baryonic matter: $w = 0$
 - Radiation: $w = \frac{1}{3}$




Figure credit: Florian Wolz

Λ CDM model

$$\rho_{\text{tot}} = \rho_r + \rho_m + \rho_\Lambda$$

- Equation of state: $p = w \cdot \rho$
- EOS-parameter w for different matter species:
 - Dark matter and baryonic matter: $w = 0$
 - Radiation: $w = \frac{1}{3}$
 - Dark energy: $w = -1$




Figure credit: Florian Wolz

Friedmann Equations

- Equation of state: $p = w \cdot \rho$
- Inserting the E.O.S into the continuity equation $\dot{\rho} = -3(\rho + p)\frac{\dot{a}}{a}$ yields:
$$\dot{\rho} = -3\rho(1 + w)\frac{\dot{a}}{a}$$

Friedmann Equations

- Equation of state:


$$p = w \cdot \rho$$

- Inserting the E.O.S into the continuity equation $\dot{\rho} = -3(\rho + p)\frac{\dot{a}}{a}$ yields:

$$\dot{\rho} = -3\rho(1 + w)\frac{\dot{a}}{a}$$

... (your exercise later)

$$\frac{\rho(a)}{\rho_0} = \left(\frac{a}{a_0}\right)^{-3(1+w)}$$



The subscript 0 refers to the current time “today”

The Hubble parameter

- The first Friedmann equation describes the rate of expansion as a function of the energy content ρ of the universe:

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3} \rho_{\text{tot}} \quad (\text{i})$$

The Hubble parameter

- The first Friedmann equation describes the rate of expansion as a function of the energy content ρ of the universe:

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3} \rho_{\text{tot}} \quad (\text{i})$$

- The rate of expansion of the universe is called the **Hubble parameter**:

$$H(t) = \frac{\dot{a}}{a}$$

The Hubble parameter

- The first Friedmann equation describes the rate of expansion as a function of the energy content ρ of the universe:

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3} \rho_{\text{tot}} \quad (\text{i})$$

- The rate of expansion of the universe is called the **Hubble parameter**:

$$H(t) = \frac{\dot{a}}{a}$$

- The Hubble parameter **today** is called the **Hubble constant**:

$$H_0 = H(t = t_0)$$


today

The Hubble parameter

- The first Friedmann equation describes the rate of expansion as a function of the energy content ρ of the universe:

$$H^2(t) = \frac{8\pi G}{3} \rho_{\text{tot}} \quad (\text{i})$$

- The rate of expansion of the universe is called the **Hubble parameter**:

$$H(t) = \frac{\dot{a}}{a}$$

- The Hubble parameter **today** is called the **Hubble constant**:

$$H_0 = H(t = t_0)$$


today

The Hubble parameter

$$H^2(t) = \frac{8\pi G_N}{3} \rho_{\text{tot}}(t)$$

The Hubble parameter

$$H^2(t) = \frac{8\pi G_N}{3} \rho_{\text{tot}}(t)$$

- Defining $\rho_{\text{crit}} = \frac{3H_0^2}{8\pi G_N}$
- 
- density of flat universe

The Hubble parameter

$$H^2(t) = \frac{8\pi G_N}{3} \rho_{\text{tot}}(t)$$

- Defining $\rho_{\text{crit}} = \frac{3H_0^2}{8\pi G_N}$

$$\frac{H^2(a)}{H_0^2} = \frac{\rho_{\text{tot}}(a)}{\rho_{\text{crit}}}$$

The Hubble parameter

$$H^2(t) = \frac{8\pi G_N}{3} \rho_{\text{tot}}(t)$$

- Defining $\rho_{\text{crit}} = \frac{3H_0^2}{8\pi G_N}$ and the fractional energy densities $\Omega_I = \frac{\rho_I}{\rho_{\text{crit}}}$:

$$\frac{H^2(a)}{H_0^2} = \frac{\rho_{\text{tot}}(a)}{\rho_{\text{crit}}}$$

The Hubble parameter

$$H^2(t) = \frac{8\pi G_N}{3} \rho_{\text{tot}}(t)$$

- Defining $\rho_{\text{crit}} = \frac{3H_0^2}{8\pi G_N}$ and the fractional energy densities $\Omega_I = \frac{\rho_I}{\rho_{\text{crit}}}$:

$$\frac{H^2(a)}{H_0^2} = \frac{\rho_{\text{tot}}(a)}{\rho_{\text{crit}}} = \Omega_{\text{tot}}(a)$$

The Hubble parameter

$$H^2(t) = \frac{8\pi G_N}{3} \rho_{\text{tot}}(t)$$

- Defining $\rho_{\text{crit}} = \frac{3H_0^2}{8\pi G_N}$ and the fractional energy densities $\Omega_I = \frac{\rho_I}{\rho_{\text{crit}}}$:

$$\frac{H^2(a)}{H_0^2} = \frac{\rho_{\text{tot}}(a)}{\rho_{\text{crit}}} = \Omega_{\text{tot}}(a) = \Omega_{r,0} \left(\frac{a_0}{a}\right)^4 + \Omega_{m,0} \left(\frac{a_0}{a}\right)^3 + \Omega_{k,0} \left(\frac{a_0}{a}\right)^2 + \Omega_{\Lambda,0}$$

The Hubble parameter

$$H^2(t) = \frac{8\pi G_N}{3} \rho_{\text{tot}}(t)$$

- Defining $\rho_{\text{crit}} = \frac{3H_0^2}{8\pi G_N}$ and the fractional energy densities $\Omega_I = \frac{\rho_I}{\rho_{\text{crit}}}$:

$$\frac{H^2(a)}{H_0^2} = \frac{\rho_{\text{tot}}(a)}{\rho_{\text{crit}}} = \Omega_{\text{tot}}(a) = \Omega_{r,0} \left(\frac{a_0}{a}\right)^4 + \Omega_{m,0} \left(\frac{a_0}{a}\right)^3 + \Omega_{k,0} \left(\frac{a_0}{a}\right)^2 + \Omega_{\Lambda,0}$$

Defining the redshift $1 + z = \frac{a_0}{a}$, one can rewrite this in the famous form:

The Hubble parameter

$$H^2(t) = \frac{8\pi G_N}{3} \rho_{\text{tot}}(t)$$

- Defining $\rho_{\text{crit}} = \frac{3H_0^2}{8\pi G_N}$ and the fractional energy densities $\Omega_I = \frac{\rho_I}{\rho_{\text{crit}}}$:

$$\frac{H^2(a)}{H_0^2} = \frac{\rho_{\text{tot}}(a)}{\rho_{\text{crit}}} = \Omega_{\text{tot}}(a) = \Omega_{r,0} \left(\frac{a_0}{a}\right)^4 + \Omega_{m,0} \left(\frac{a_0}{a}\right)^3 + \Omega_{k,0} \left(\frac{a_0}{a}\right)^2 + \Omega_{\Lambda,0}$$

Defining the redshift $1 + z = \frac{a_0}{a}$, one can rewrite this in the famous form:

$$\boxed{\frac{H^2(a)}{H_0^2} = \Omega_{r,0} (1+z)^4 + \Omega_{m,0} (1+z)^3 + \Omega_{k,0} (1+z)^2 + \Omega_{\Lambda,0}}$$

The Hubble parameter

$$H^2(t) = \frac{8\pi G_N}{3} \rho_{\text{tot}}(t)$$

- Defining $\rho_{\text{crit}} = \frac{3H_0^2}{8\pi G_N}$ and the fractional energy densities $\Omega_I = \frac{\rho_I}{\rho_{\text{crit}}}$:

$$\frac{H^2(a)}{H_0^2} = \frac{\rho_{\text{tot}}(a)}{\rho_{\text{crit}}} = \Omega_{\text{tot}}(a) = \Omega_{r,0} \left(\frac{a_0}{a}\right)^4 + \Omega_{m,0} \left(\frac{a_0}{a}\right)^3 + \Omega_{k,0} \left(\frac{a_0}{a}\right)^2 + \Omega_{\Lambda,0}$$

Defining the redshift $1 + z = \frac{a_0}{a}$, one can rewrite this in the famous form:

$$\boxed{\frac{H^2(a)}{H_0^2} = \Omega_{r,0} (1+z)^4 + \Omega_{m,0} (1+z)^3 + \Omega_{k,0} (1+z)^2 + \Omega_{\Lambda,0}}$$

$$\Omega_{I,0} = \Omega_I(t_0) = \Omega_I$$

Distances in an expanding universe

Distances in an expanding universe

- In an expanding universe, consider two coordinate systems:
 - a comoving system that expands with the universe
 - a fixed coordinate system, in which objects drift apart with the expansion

Distances in an expanding universe

- In an expanding universe, consider two coordinate systems:
 - a comoving system that expands with the universe
 - a fixed coordinate system, in which objects drift apart with the expansion
- A **physical distance** r_p is related to the **comoving distance** χ via: $r_p(t) = a(t)\chi(t)$

Distances in an expanding universe

- In an expanding universe, consider two coordinate systems:
 - a comoving system that expands with the universe
 - a fixed coordinate system, in which objects drift apart with the expansion
- A **physical distance** r_p is related to the **comoving distance** χ via: $r_p(t) = a(t)\chi(t)$
- To find an expression for χ , consider a photon travelling an infinitesimal distance r_p :

$$1 = c = \frac{dr_p}{dt}$$

Distances in an expanding universe

- In an expanding universe, consider two coordinate systems:
 - a comoving system that expands with the universe
 - a fixed coordinate system, in which objects drift apart with the expansion
- A **physical distance** r_p is related to the **comoving distance** χ via: $r_p(t) = a(t)\chi(t)$
- To find an expression for χ , consider a photon travelling an infinitesimal distance r_p :

$$1 = c = \frac{dr_p}{dt} = a \frac{d\chi}{dt}$$

Distances in an expanding universe

- In an expanding universe, consider two coordinate systems:
 - a comoving system that expands with the universe
 - a fixed coordinate system, in which objects drift apart with the expansion
- A **physical distance** r_p is related to the **comoving distance** χ via: $r_p(t) = a(t)\chi(t)$
- To find an expression for χ , consider a photon travelling an infinitesimal distance r_p :

$$1 = c = \frac{dr_p}{dt} = a \frac{d\chi}{dt} \implies \chi(t) = \int_{t_1}^{t_2} \frac{dt'}{a(t')}$$

Distances in an expanding universe

- In an expanding universe, consider two coordinate systems:
 - a comoving system that expands with the universe
 - a fixed coordinate system, in which objects drift apart with the expansion
- A **physical distance** r_p is related to the **comoving distance** χ via: $r_p(t) = a(t)\chi(t)$
- To find an expression for χ , consider a photon travelling an infinitesimal distance r_p :

$$1 = c = \frac{dr_p}{dt} = a \frac{d\chi}{dt} \implies \boxed{\chi(t) = \int_{t_1}^{t_2} \frac{dt'}{a(t')} = \int_{z_1}^{z_2} \frac{dz'}{H(z')}}$$

Distances in an expanding universe

- The angular diameter distance D_A is defined such that it holds

$$\theta = \frac{s}{D_A}$$

angle

physical size of the object

ang. diam. distance

Distances in an expanding universe


- The angular diameter distance D_A is defined such that it holds

$$\theta = \frac{s}{D_A}$$

angle

physical size of the object

ang. diam. distance



Distances in an expanding universe

- The angular diameter distance D_A is defined such that it holds

$$\theta = \frac{s}{D_A}$$

$$D_A(t) = a(t)\chi(t) = \frac{1}{1+z(t)} \int_0^{z(t)} \frac{dz'}{H(z')}$$

*To derive an equation for D_A , note that the proper size, s , of the object can also be expressed as $s = \chi(t)\theta \cdot a(t)$, where $\chi(t)\theta$ corresponds to the comoving size of the object.

Distances in an expanding universe


- The **luminosity distance** D_L is defined such that it holds

$$F = \frac{L}{4\pi D_L^2}$$

Distances in an expanding universe

- The **luminosity distance** D_L is defined such that it holds

$$F = \frac{L}{4\pi D_L^2}$$



Distances in an expanding universe

- The **luminosity distance** D_L is defined such that it holds

$$F = \frac{L}{4\pi D_L^2}$$


$$D_L(t) = \frac{\chi(t)}{a(t)} = [1 + z(t)] \int_0^{z(t)} \frac{c \, dz'}{H(z')}$$

*Since the Universe is expanding, it holds that $F = \frac{L a^2}{4\pi \chi^2(a)}$, where the additional factor of a^2 comes from the fact that the expansion of the Universe leads to a dilution of photons ($\propto a$) and to an increase in wavelength ($\propto a$).

How to solve the Hubble tension?


The Hubble tension

- **Indirect measurements:** cosmic microwave background (CMB), baryon acoustic oscillations (BAO), galaxy clustering
- **Direct measurements:** distance ladder (Cepheids, TRGB, SNe, ...), gravitational lensing, ...




The Hubble tension

- **Indirect measurements:** cosmic microwave background (CMB), baryon acoustic oscillations (BAO), galaxy clustering → precise but model dependent
- **Direct measurements:** distance ladder (Cepheids, TRGB, SNe, ...), gravitational lensing, ...




The Hubble tension

- **Indirect measurements:** cosmic microwave background (CMB), baryon acoustic oscillations (BAO), galaxy clustering → precise but model dependent
- **Direct measurements:** distance ladder (Cepheids, TRGB, SNe, ...), gravitational lensing, ... → less precise due to astrophysical modelling but independent of cosmological model



The Hubble tension


- General strategy to solve the H_0 tension:
Assume direct measurements are correct
and **change cosmological model** in order
to *infer* a higher H_0
- Goal: “get CMB- H_0 to ~ 73 km/s/Mpc”



Expansion rate H_0 [km/s/Mpc]


How does the CMB constrain H_0 ?

- In order to understand how we can solve the Hubble tension, we need to understand how the CMB constrains H_0
- The CMB provides the most important cosmological probe when it comes to constraining the parameters of the cosmological model with high accuracy
- However, it is an indirect probe of H_0 : **it depends on the cosmological model** that is assumed → the CMB constrains the universe **at early times** and **predicts H_0 today**




Credit: WMAP Collaboration

Sky in optical



Credit: Gaia collaboration


Sky in microwave



70 GHz *Planck* map


Sky in microwave

$\sim 2.7K$




70 GHz *Planck* map

Sky in microwave



70 GHz Planck map

Sky in microwave



-300

μK

300


Credit: *Planck* Collaboration

The CMB

- How to analyse the CMB map with temperature fluctuations?
 - 1. Decompose the temperature fluctuations into a set of waves with various wavelength
 - 2. Plot the strength of each wavelength: **Power spectrum**
- In 1 dimension: decomposing into waves = Fourier transform
- On the unit sphere: decomposing into waves = **Spherical harmonics** decomposition

The CMB

- Decompose temperature fluctuations: $\Delta T(\theta, \phi) = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} a_{\ell m} Y_{\ell m}(\theta, \phi)$





coefficients

spherical harmonics

The CMB


- Decompose temperature fluctuations: $\Delta T(\theta, \phi) = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} a_{\ell m} Y_{\ell m}(\theta, \phi)$
- The $a_{\ell m}$ contain the same information as the map itself



spherical harmonics

The CMB


- Decompose temperature fluctuations: $\Delta T(\theta, \phi) = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} a_{\ell m} Y_{\ell m}(\theta, \phi)$



- The $a_{\ell m}$ contain the same information as the map itself
- ℓ defines the multipole, the higher ℓ the smaller the scale

spherical harmonics

The CMB




- Decompose temperature fluctuations: $\Delta T(\theta, \phi) = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} a_{\ell m} Y_{\ell m}(\theta, \phi)$
- The $a_{\ell m}$ contain the same information as the map itself
- ℓ defines the multipole, the higher ℓ the smaller the scale
- The power spectrum is then given by:

$$C_{\ell} = \frac{1}{2\ell + 1} \sum_{m=-\ell}^{\ell} a_{\ell,m} a_{\ell,m}^*$$

spherical harmonics

The CMB

Planck Collaboration




$$D_\ell = \frac{\ell(\ell + 1)}{2\pi} C_\ell$$

*You will need this in the
hands-on session later

The CMB

How to analyse data like this?




- To predict the CMB power spectrum: one needs to solve full fluid-equations (the Boltzmann equations) for all components of the universe
- This can be done with cosmological Boltzmann solvers like **CAMB** (Lewis&Bridle 2002) and **CLASS** (Blas, Lesgourgues, Tram 2011)
- These Boltzmann solvers take as input the Λ CDM parameters and can be sampled within MCMC samplers

*You will need this in the hands-on session later

The CMB


Λ CDM model: 6 parameters



*You will need this in the
hands-on session later

The CMB

Λ CDM model: 6 parameters




- $\omega_{\text{cdm}} = h^2 \Omega_{\text{cdm}}$:
- $\omega_b = h^2 \Omega_b$:
- $h = H_0 / (100 \text{ km/s/Mpc})$:
- τ_{reio} :
- A_s :
- n_s :

*You will need this in the hands-on session later

The CMB

Λ CDM model: 6 parameters




- $\omega_{\text{cdm}} = h^2 \Omega_{\text{cdm}}$: physical energy density in CDM
- $\omega_b = h^2 \Omega_b$:
- $h = H_0 / (100 \text{ km/s/Mpc})$:
- τ_{reio} :
- A_s :
- n_s :

*You will need this in the hands-on session later

The CMB

Λ CDM model: 6 parameters




- $\omega_{\text{cdm}} = h^2 \Omega_{\text{cdm}}$: physical energy density in CDM
- $\omega_b = h^2 \Omega_b$: physical energy density in baryons
- $h = H_0 / (100 \text{ km/s/Mpc})$:
- τ_{reio} :
- A_s :
- n_s :

*You will need this in the hands-on session later

The CMB

Λ CDM model: 6 parameters




- $\omega_{\text{cdm}} = h^2 \Omega_{\text{cdm}}$: physical energy density in CDM
- $\omega_b = h^2 \Omega_b$: physical energy density in baryons
- $h = H_0 / (100 \text{ km/s/Mpc})$: dimensionless Hubble constant
- τ_{reio} :
- A_s :
- n_s :

*You will need this in the hands-on session later

The CMB

Λ CDM model: 6 parameters




- $\omega_{\text{cdm}} = h^2 \Omega_{\text{cdm}}$: physical energy density in CDM
- $\omega_b = h^2 \Omega_b$: physical energy density in baryons
- $h = H_0 / (100 \text{ km/s/Mpc})$: dimensionless Hubble constant
- τ_{reio} : optical depth to reionization
- A_s :
- n_s :

*You will need this in the hands-on session later

The CMB

Λ CDM model: 6 parameters




- $\omega_{\text{cdm}} = h^2 \Omega_{\text{cdm}}$: physical energy density in CDM
- $\omega_b = h^2 \Omega_b$: physical energy density in baryons
- $h = H_0/(100 \text{ km/s/Mpc})$: dimensionless Hubble constant
- τ_{reio} : optical depth to reionization
- A_s : amplitude of the primordial power spectrum
- n_s :

*You will need this in the hands-on session later

The CMB

Λ CDM model: 6 parameters




- $\omega_{\text{cdm}} = h^2 \Omega_{\text{cdm}}$: physical energy density in CDM
- $\omega_b = h^2 \Omega_b$: physical energy density in baryons
- $h = H_0/(100 \text{ km/s/Mpc})$: dimensionless Hubble constant
- τ_{reio} : optical depth to reionization
- A_s : amplitude of the primordial power spectrum
- n_s : tilt of the primordial power spectrum

*You will need this in the hands-on session later

The CMB

Planck Collaboration



$$D_\ell = \frac{\ell(\ell + 1)}{2\pi} C_\ell$$

*You will need this in the hands-on session later

The baryon acoustic oscillations (BAO)

- The early universe was a hot plasma of baryons and photons

The baryon acoustic oscillations (BAO)

- The early universe was a hot plasma of baryons and photons
- Small density fluctuations sourced pressure-density waves
 - Gravity as driving force
 - Photon pressure as restoring force

The baryon acoustic oscillations (BAO)

- The early universe was a hot plasma of baryons and photons
- Small density fluctuations sourced pressure-density waves
 - Gravity as driving force
 - Photon pressure as restoring force
- Like rain drops on the water surface many waves overlap



Credit: Mabel Amber

The baryon acoustic oscillations (BAO)

- The early universe was a hot plasma of baryons and photons
- Small density fluctuations sourced pressure-density waves
 - Gravity as driving force
 - Photon pressure as restoring force
- Like rain drops on the water surface many waves overlap
- Once electrons and protons recombine, photons can travel freely → the density waves freeze



Credit: Mabel Amber

The baryon acoustic oscillations (BAO)

- Let's compute the (comoving) distance that the sound waves travelled: the sound horizon

The baryon acoustic oscillations (BAO)

- Let's compute the (comoving) distance that the sound waves travelled: the sound horizon
- The sound waves travel from the big bang to the time of recombination t^* (when protons and electrons combine into atoms): $r_s = \int_0^{t^*} c_s(t) dt$

The baryon acoustic oscillations (BAO)

- Let's compute the (comoving) distance that the sound waves travelled: the sound horizon
- The sound waves travel from the big bang to the time of recombination t^*

(when protons and electrons combine into atoms): $r_s = \int_0^{t^*} \frac{c_s(t)}{a(t)} dt$

- While the sound waves travel, the universe expands. This slows down the sound waves in the comoving coordinate frame

The baryon acoustic oscillations (BAO)


- Comoving sound horizon:

$$r_s = \int_0^{t^*} \frac{c_s(t)}{a(t)} dt$$

The baryon acoustic oscillations (BAO)

- Comoving sound horizon:

$$r_s = \int_0^{t^*} \frac{c_s(t)}{a(t)} dt = \int_0^{a^*} \frac{c_s(a)}{\dot{a}} da$$



$$\frac{da}{dt} = \dot{a}$$

The baryon acoustic oscillations (BAO)

- Comoving sound horizon:

$$r_s = \int_0^{t^*} \frac{c_s(t)}{a(t)} dt = \int_0^{a^*} \frac{c_s(a)}{\dot{a}} da = - \int_{\infty}^{z^*} \frac{c_s(z)a^2}{\dot{a}} dz$$



$$\frac{dz}{da} = \frac{d(1/a)}{da} = -\frac{1}{a^2}$$

The baryon acoustic oscillations (BAO)

- Comoving sound horizon:

$$r_s = \int_0^{t^*} \frac{c_s(t)}{a(t)} dt = \int_0^{a^*} \frac{c_s(a)}{\dot{a}a} da = - \int_{\infty}^{z^*} \frac{c_s(z)a^2}{\dot{a}a} dz = \int_{z^*}^{\infty} \frac{c_s(z)}{H(z)} dz$$

The baryon acoustic oscillations (BAO)

- Comoving sound horizon:

$$r_s = \int_0^{t^*} \frac{c_s(t)}{a(t)} dt = \int_0^{a^*} \frac{c_s(a)}{\dot{a}} da = - \int_{\infty}^{z^*} \frac{c_s(z)a^2}{\dot{a}} dz = \int_{z^*}^{\infty} \frac{c_s(z)}{H(z)} dz$$


- The sound speed can be expressed as (e.g. Dodelson&Schmidt, 2020):

$$c_s(z) = \frac{1}{\sqrt{3(1 + R(z))}}$$

- With R being the baryon-photon ratio:

$$R(z) = \frac{3\omega_b}{4\omega_\gamma} \frac{1}{1+z}$$


The baryon acoustic oscillations (BAO)



- What we observe is the *angular size of the sound horizon* θ_s

$$r_s^{\text{phys}}(z^*) = a(z^*) r_s(z^*)$$

The baryon acoustic oscillations (BAO)




- What we observe is the *angular size of the sound horizon* θ_s
- In the small-angle approximation:

$$\theta_s = \frac{r_s^{\text{phys}}(z^*)}{D_A(z^*)}$$

$$r_s^{\text{phys}}(z^*) = a(z^*) r_s(z^*)$$

The baryon acoustic oscillations (BAO)



- What we observe is the *angular size of the sound horizon* θ_s
- In the small-angle approximation:
- where the angular diameter distance is (see slide earlier):

$$D_A(t) = \int_0^{z(t)} \frac{dz'}{H(z')}$$

$$r_s^{\text{phys}}(z^*) = a(z^*) r_s(z^*)$$

How does the CMB constrain H_0 ?

- We know that the CMB directly constrains

$$\theta_s = \frac{r_s^{\text{phys}}(z^*)}{D_A(z^*)} = \frac{\int_{z^*}^{\infty} c_s(z) dz / H(z)}{\int_0^{z^*} dz / H(z)}$$

How does the CMB constrain H_0 ?

- We know that the CMB directly constrains

$$\theta_s = \frac{r_s^{\text{phys}}(z^*)}{D_A(z^*)} = \frac{\int_{z^*}^{\infty} c_s(z) dz / H(z)}{\int_0^{z^*} dz / H(z)}$$

- Cosmology enters mainly over the Hubble parameter here:

$$H(z) = H_0 \sqrt{\Omega_r(1+z)^4 + \Omega_m(1+z)^3 + \Omega_\Lambda}$$

How does the CMB constrain H_0 ?

- We know that the CMB directly constrains

$$\theta_s = \frac{r_s^{\text{phys}}(z^*)}{D_A(z^*)} = \frac{\int_{z^*}^{\infty} c_s(z) dz / H(z)}{\int_0^{z^*} dz / H(z)}$$

- Cosmology enters mainly over the Hubble parameter here:


$$H(z) = H_0 \sqrt{\Omega_r(1+z)^4 + \Omega_m(1+z)^3 + \Omega_\Lambda}$$

- If we get Ω_r , Ω_m , Ω_Λ from somewhere else, this becomes an implicit equation for H_0 – How does the CMB constrain Ω_r , Ω_m , Ω_Λ ?


How does the CMB constrain Ω_r ?

- The radiation density Ω_r is precisely measured by the CMB temperature

$$T = 2.725 \pm 0.002 \text{ K} \text{ (COBE - FIRAS measurement)}$$



Credit: Berkeley Center for Cosmological Physics



How does the CMB constrain Ω_r ?

- The radiation density Ω_r is precisely measured by the CMB temperature

$$T = 2.725 \pm 0.002 \text{ K} \text{ (COBE - FIRAS measurement)}$$

- For a black-body spectrum, the energy density in radiation is then given by (e.g. Weinberg 2008, Ch. 2.1):

$$\rho_{0,\text{CMB}} = \int_0^{\infty} h\nu \cdot n(\nu) d\nu = \frac{8\pi^5 k_B^4}{15h^3 c^3} T^4 = 4.64 \cdot 10^{-34} \text{ g/cm}^3$$

How does the CMB constrain Ω_r ?

- The radiation density Ω_r is precisely measured by the CMB temperature

$$T = 2.725 \pm 0.002 \text{ K} \text{ (COBE - FIRAS measurement)}$$

- For a black-body spectrum, the energy density in radiation is then given by (e.g. Weinberg 2008, Ch. 2.1):

$$\rho_{0,\text{CMB}} = \int_0^{\infty} h\nu \cdot n(\nu) d\nu = \frac{8\pi^5 k_B^4}{15h^3 c^3} T^4 = 4.64 \cdot 10^{-34} \text{ g/cm}^3$$

- Taking into account the earlier time of decoupling of neutrinos, one finds:

$$\rho_\nu = 0.4 \rho_{0,\text{CMB}}$$

- Since the CMB and neutrinos are by far the dominant contribution:

$$\Omega_r = 1.4 \rho_{0,\text{CMB}} / \rho_{\text{crit}} = 4.15 \cdot 10^{-5} h^{-2}$$

For more about neutrinos, see Olga Mena's lecture

How does the CMB constrain Ω_m ?

- It does not constrain Ω_m , but $\omega_m = h^2\Omega_m$, where $\omega_m = \omega_{\text{cdm}} + \omega_b$

How does the CMB constrain Ω_m ?

- It does not constrain Ω_m , but $\omega_m = h^2 \Omega_m$, where $\omega_m = \omega_{\text{cdm}} + \omega_b$
- ω_m is constrained by the height of the acoustic peaks:
 - a smaller ω_m leads to a later time of matter-radiation equality
 - this leads to a stronger decay of the gravitational potentials at recombination → **early integrated Sachs-Wolfe effect (eISW)**
 - Since the eISW adds in phase with the BAO, this leads to a boost of all peaks, particularly the first peak.

How does the CMB constrain Ω_Λ ?

- The sum of the energy densities has to satisfy:

$$\Omega_r + \Omega_m + \Omega_\Lambda = \Omega_k$$

- For a flat universe:

$$\Omega_r + \Omega_m + \Omega_\Lambda = 1$$

- Hence, if we know Ω_r and $\omega_m = h^2\Omega_m$, we can compute

$$\Omega_\Lambda = 1 - \Omega_r - \omega_m/h^2$$

How does the CMB constrain H_0 ?

How does the CMB constrain H_0 ?

- We know that the CMB directly constrains

$$\theta_s = \frac{r_s^{\text{phys}}(z^*)}{D_A(z^*)} = \frac{\int_{z^*}^{\infty} c_s(z) dz / H(z)}{\int_0^{z^*} dz / H(z)}$$

- Cosmology enters mainly over the Hubble parameter here:

$$H(z) = H_0 \sqrt{\Omega_r(1+z)^4 + \Omega_m(1+z)^3 + \Omega_\Lambda}$$

How does the CMB constrain H_0 ?

- We know that the CMB directly constrains

$$\theta_s = \frac{r_s^{\text{phys}}(z^*)}{D_A(z^*)} = \frac{\int_{z^*}^{\infty} c_s(z) dz / H(z)}{\int_0^{z^*} dz / H(z)}$$

- Cosmology enters mainly over the Hubble parameter here:

$$H(z) = H_0 \sqrt{\Omega_r(1+z)^4 + \Omega_m(1+z)^3 + \Omega_\Lambda}$$

- Since we can constrain Ω_r and ω_m independently, this becomes an implicit equation for h

$$\frac{H^2(z)}{(100 \text{ km/s/Mpc})^2} = h^2 \Omega_r (1+z)^4 + \omega_m (1+z)^3 + (h^2 - \omega_m - h^2 \Omega_r)$$

How does the CMB constrain H_0 ?

- We know that the CMB directly constrains

$$\theta_s = \frac{r_s^{\text{phys}}(z^*)}{D_A(z^*)} = \frac{\int_{z^*}^{\infty} c_s(z) dz / H(z)}{\int_0^{z^*} dz / H(z)}$$

* $c_s(z)$ and z^* also depend on cosmology but we neglect that here

- Cosmology enters mainly over the Hubble parameter here:


$$H(z) = H_0 \sqrt{\Omega_r(1+z)^4 + \Omega_m(1+z)^3 + \Omega_\Lambda}$$

- Since we can constrain Ω_r and ω_m independently, this becomes an implicit equation for h

$$\frac{H^2(z)}{(100 \text{ km/s/Mpc})^2} = h^2 \Omega_r (1+z)^4 + \omega_m (1+z)^3 + (h^2 - \omega_m - h^2 \Omega_r)$$

Solutions to the Hubble tension


Solutions to the Hubble tension



$$\theta_s = \frac{r_s(z^*)}{D_A(z^*)} = \frac{\int_{z^*}^{\infty} c_s(z) dz / H(z)}{\int_0^{z^*} dz / H(z)}$$

- θ_s is measured precisely by CMB
 $\rightarrow \theta_s$ fixed
- Two options to solve the Hubble tension:

Solutions to the Hubble tension




$$\theta_s = \frac{r_s(z^*)}{D_A(z^*)} = \frac{\int_{z^*}^{\infty} c_s(z) dz / H(z)}{\int_0^{z^*} dz / H(z)}$$

- θ_s is measured precisely by CMB
→ θ_s fixed
- Two options to solve the Hubble tension:

Early-time solutions
modify r_s

Solutions to the Hubble tension



- θ_s is measured precisely by CMB
→ θ_s fixed
- Two options to solve the Hubble tension:

Early-time solutions
modify r_s

Late-time solutions
modify D_A

Late-time solutions

- Modify $D_A(z^*) = \int_0^{z^*} \frac{dz}{H(z)}$ by modifying the expansion rate between today (0) and recombination z^* (since $\Omega_r \approx 0$ at times after z^*):

$$H(z) = H_0 \sqrt{\Omega_m(1+z)^3 + 1 - \Omega_m}$$

Late-time solutions

- Modify $D_A(z^*) = \int_0^{z^*} \frac{dz}{H(z)}$ by modifying the expansion rate between today (0) and recombination z^* (since $\Omega_r \approx 0$ at times after z^*):

$$H(z) = H_0 \sqrt{\Omega_m(1+z)^3 + 1 - \Omega_m}$$

- However, $H(z)$ is well constrained by galaxy BAO data and supernova data
→ It is challenging to increase H_0 enough
- But there are some promising approaches (see other lectures)

Early-time solutions

- Modify $r_s = \int_{z^*}^{\infty} \frac{c_s(z) dz}{H(z)} \rightarrow$ 3 options:
 - ▶ modify z^*
 - ▶ modify $c_s(z)$
 - ▶ modify $H(z)$

Early-time solutions

- Modify $r_s = \int_{z^*}^{\infty} \frac{c_s(z) dz}{H(z)} \rightarrow 3 \text{ options:}$

► **modify z^***

- modify $c_s(z)$
- modify $H(z)$



E.g. by modifying the mass of the electron m_e :

- This shifts the energy levels of the atoms,
- and changes the ionization energy,
- which changes the time of recombination z^*

Early-time solutions

- Modify $r_s = \int_{z^*}^{\infty} \frac{c_s(z) dz}{H(z)} \rightarrow$ 3 options:
 - ▶ modify z^*
 - ▶ **modify $c_s(z)$**
 - ▶ modify $H(z)$

Early-time solutions

- Modify $r_s = \int_{z^*}^{\infty} \frac{c_s(z) dz}{H(z)} \rightarrow 3 \text{ options:}$
 - ▶ modify z^*
 - ▶ modify $c_s(z)$
 - ▶ **modify $H(z)$** →


E.g. by introducing an energy density, which boosts the expansion rate before recombination z^* (at early times $\Omega_\Lambda \approx 0$):

$$H(z) = H_0 \sqrt{\Omega_r(1+z)^4 + \Omega_m(1+z)^3 + \Omega_q}$$

This leads to a smaller r_s .

The most successful of such ideas is **EDE**


Idea behind Early Dark Energy



Angular scale of sound horizon θ_s
measured with 0.03% precision by *Planck*.

$$\theta_s = \frac{r_s(z^*)}{D_A(z^*)} = \frac{\int_{z^*}^{\infty} c_s(z) dz / H(z)}{\int_0^{z^*} dz / H(z)}$$

Idea behind Early Dark Energy




Angular scale of sound horizon θ_s
measured with 0.03% precision by *Planck*.

Introducing an additional component
before recombination reduces r_s .

$$\theta_s = \frac{r_s(z^*)}{D_A(z^*)} = \frac{\int_{z^*}^{\infty} c_s(z) dz / H(z)}{\int_0^{z^*} dz / H(z)}$$

Idea behind Early Dark Energy



$$\theta_s = \frac{r_s(z^*)}{D_A(z^*)} = \frac{\int_{z^*}^{\infty} c_s(z) dz / H(z)}{\int_0^{z^*} dz / H(z)}$$

Angular scale of sound horizon θ_s
measured with 0.03% precision by *Planck*.


Introducing an additional component
before recombination reduces r_s .



θ_s fixed

Angular diameter distance
 D_A decreases.

Idea behind Early Dark Energy



$$\theta_s = \frac{r_s(z^*)}{D_A(z^*)} = \frac{\int_{z^*}^{\infty} c_s(z) dz / H(z)}{\int_0^{z^*} dz / H(z)}$$

Angular scale of sound horizon θ_s
measured with 0.03% precision by *Planck*.

Introducing an additional component
before recombination reduces r_s .

θ_s fixed

Angular diameter distance
 D_A decreases.

$H(z) = H_0 \sqrt{\Omega_m(z) + \Omega_r(z) + \Omega_\Lambda}$

H_0 increases.


Short history of Early Dark Energy

- EDE has already been studied before the Hubble tension emerged in the context of quintessence models (Doran++ 2001, Wetterich++ 2004, Doran&Robbers 2006, Kamionkowski++ 2014)
- The axion-like EDE was proposed as a solution to the Hubble tension (Karwal&Kamionkowski 2016, Poulin++ 2018, 2019)
- There are many versions of the EDE model, but we will focus on the (most commonly studied) axion-like EDE model
- The axion-like EDE model is modelled as a **scalar field**
- Here: Only background equations

Scalar fields in an expanding spacetime

Intuition

- A scalar field $\phi(t, \vec{x})$ assigns each point in spacetime a single number
- Examples
 - Temperature, density, pressure,... as a function of position and time
 - Potential fields like the gravitational potential, electric potential
 - The inflaton driving inflation is most commonly a scalar field




Scalar field

- The action of a scalar field minimally coupled to the metric is:

$$\mathcal{S}_\phi = \int d^4x \sqrt{|\det(g)|} \left[-\frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - V(\phi) \right]$$

integral over whole spacetime determinant of metric kinetic term potential term



Very similar to inflation: see William Giare's lecture

Scalar field

- The action of a scalar field minimally coupled to the metric is:

$$\mathcal{S}_\phi = \int d^4x \sqrt{|\det(g)|} \left[-\frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - V(\phi) \right]$$

integral over whole spacetime determinant of metric kinetic term potential term

- One can compute the energy momentum tensor of the scalar field via:

$$T_{\mu\nu} = \partial_\mu \phi \partial_\nu \phi - g_{\mu\nu} \left(\frac{1}{2} g^{\alpha\beta} \partial_\alpha \phi \partial_\beta \phi - V(\phi) \right)$$

Scalar field

- This gives for the energy-momentum tensor:

$$T_{\mu\nu} = \partial_\mu \phi \partial_\nu \phi - g_{\mu\nu} \left(\frac{1}{2} g^{\alpha\beta} \partial_\alpha \phi \partial_\beta \phi - V(\phi) \right) = \dot{\phi}^2 \delta_\mu^0 \delta_\nu^0 + \left[\frac{\dot{\phi}^2}{2} - V(\phi) \right] g_{\mu\nu}$$

Scalar field

- This gives for the energy-momentum tensor:

$$T_{\mu\nu} = \partial_\mu \phi \partial_\nu \phi - g_{\mu\nu} \left(\frac{1}{2} g^{\alpha\beta} \partial_\alpha \phi \partial_\beta \phi - V(\phi) \right) = \dot{\phi}^2 \delta_\mu^0 \delta_\nu^0 + \left[\frac{\dot{\phi}^2}{2} - V(\phi) \right] g_{\mu\nu}$$

- By comparing $T_{\mu\nu}$ to the energy momentum tensor of the perfect fluid:
 $T_{\mu\nu} = (\rho + p)\delta_\mu^0 \delta_\nu^0 + pg_{\mu\nu}$, one can read off the energy density and pressure of the scalar field:

$$\rho_\phi = \frac{\dot{\phi}^2}{2} + V(\phi), \quad p_\phi = \frac{\dot{\phi}^2}{2} - V(\phi).$$

Scalar field

- Inserting $\rho_\phi = \frac{\dot{\phi}^2}{2} + V(\phi)$, $p_\phi = \frac{\dot{\phi}^2}{2} - V(\phi)$ into the first Friedmann equation and the continuity equation yields:


Scalar field

- Inserting $\rho_\phi = \frac{\dot{\phi}^2}{2} + V(\phi)$, $p_\phi = \frac{\dot{\phi}^2}{2} - V(\phi)$ into the first Friedmann equation and the continuity equation yields:

$$H^2 = \frac{8\pi G_N}{3} \left(\frac{\dot{\phi}^2}{2} + V(\phi) \right)$$
$$\ddot{\phi} + 3H\dot{\phi} + \frac{dV}{d\phi} = 0$$



“Hubble
friction”



Potential
term

Scalar field

- Inserting $\rho_\phi = \frac{\dot{\phi}^2}{2} + V(\phi)$, $p_\phi = \frac{\dot{\phi}^2}{2} - V(\phi)$ into the first Friedmann equation and the continuity equation yields:

$$\boxed{H^2 = \frac{8\pi G_N}{3} \left(\frac{\dot{\phi}^2}{2} + V(\phi) \right)}$$
$$\ddot{\phi} + 3H\dot{\phi} + \frac{dV}{d\phi} = 0$$

- The second equation is the **Klein-Gordon equation** for a scalar field in an expanding space, where the second term ($3H\dot{\phi}$) is the Hubble-drag term and the third term ($\frac{1}{2}dV/d\phi$) is the potential-gradient term

Early Dark Energy

Early Dark Energy

- EDE is a scalar field ϕ in a potential $V(\phi)$, which boosts the expansion rate before recombination

Early Dark Energy

- EDE is a scalar field ϕ in a potential $V(\phi)$, which boosts the expansion rate before recombination
- Potential motivated by “axion” (we take it as a phenomenological potential):

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$

Early Dark Energy

- EDE is a scalar field ϕ in a potential $V(\phi)$, which boosts the expansion rate before recombination
- Potential motivated by “axion” (we take it as a phenomenological potential):

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$

normalization of
the potential:

$$V_0 = m^2 f^2$$

m: mass

f: decay
constant

Early Dark Energy

- EDE is a scalar field ϕ in a potential $V(\phi)$, which boosts the expansion rate before recombination
- Potential motivated by “axion” (we take it as a phenomenological potential):

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$

normalization of
the potential:

$$V_0 = m^2 f^2$$

m: mass

f: decay
constant



Early Dark Energy

- EDE is a scalar field ϕ in a potential $V(\phi)$, which boosts the expansion rate before recombination
- Potential motivated by “axion” (we take it as a phenomenological potential):

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$



index n :

$n = 1$ “standard” axion; however: doesn’t decay quickly enough

$n = 3$: is preferred value by the data (Poulin++ 2020)
→ decays fast enough (commonly fixed in EDE data analysis)

Early Dark Energy

- EDE is a scalar field ϕ in a potential $V(\phi)$, which boosts the expansion rate before recombination
- Potential motivated by “axion” (we take it as a phenomenological potential):


$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$

Free parameters of the model:

- m mass ($V_0 = m^2 f^2$)
- f “decay constant”
- $\theta_i = \phi_i/f$ initial value of the field
- ($n = 3$)


Early Dark Energy

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$



Early Dark Energy

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$




Disclaimer: the following will include many “hand-wavy” approximations.

Analytical computations are too complicated otherwise and numerics is needed.

Early Dark Energy

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$




At early times:

- Dynamics governed by Klein-Gordon equation:

$$\ddot{\phi} + 3H\dot{\phi} + \cancel{\frac{dV}{d\phi}} = 0$$

Early Dark Energy

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$



At early times:

- Dynamics governed by Klein-Gordon equation:


$$\ddot{\phi} + 3H\dot{\phi} + \cancel{\frac{dV}{d\phi}} = 0$$

- ϕ starts high up in the potential, where the potential is very flat:

$$\frac{dV}{d\phi} \approx 0$$

Early Dark Energy

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$



At early times:

- Dynamics governed by Klein-Gordon equation:

$$\ddot{\phi} + 3H\dot{\phi} + \cancel{\frac{dV}{d\phi}} = 0$$

- ϕ starts high up in the potential, where the potential is very flat:

$$\frac{dV}{d\phi} \approx 0$$

- Moreover, at early times the expansion rate H is large; hence “Hubble friction” dominates:


$$3H\dot{\phi} \gg \frac{dV}{d\phi}$$

Early Dark Energy

At early times:

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$

$$\ddot{\phi} + 3H\dot{\phi} = 0$$




Early Dark Energy

At early times:

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$

$$\ddot{\phi} + 3H\dot{\phi} = 0$$



- Define $\psi = \dot{\phi}$:


$$\dot{\psi} = -3H\psi$$

Early Dark Energy

At early times:

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$

$$\ddot{\phi} + 3H\dot{\phi} = 0$$




- Define $\psi = \dot{\phi}$:
$$\dot{\psi} = -3H\psi$$
- For simplicity at first order:
$$H \approx \text{const.}$$
- Then $\dot{\phi} = \psi \sim e^{-3Ht}$

Early Dark Energy

At early times:

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$

$$\ddot{\phi} + 3H\dot{\phi} = 0$$



- Define $\psi = \dot{\phi}$:
$$\dot{\psi} = -3H\psi$$
- For simplicity at first order:
$$H \approx \text{const.}$$
- Then $\dot{\phi} = \psi \sim e^{-3Ht} \approx 0$


H large at early times

Early Dark Energy

At early times:


- At early times: $\dot{\phi} \approx 0$

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$



Early Dark Energy

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$




At early times:

- At early times: $\dot{\phi} \approx 0$
- This means for the energy density & pressure

$$\rho_\phi = \frac{\dot{\phi}^2}{2} + V(\phi), \quad p_\phi = \frac{\dot{\phi}^2}{2} - V(\phi).$$

Early Dark Energy

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$



At early times:


- At early times: $\dot{\phi} \approx 0$
- This means for the energy density & pressure

$$\rho_\phi = \frac{\dot{\phi}^2}{2} + V(\phi), \quad p_\phi = \frac{\dot{\phi}^2}{2} - V(\phi).$$

- The equation of state becomes:
$$w = \frac{p_\phi}{\rho_\phi} \approx -1$$
- Hence, at early times the field behaves like dark energy \rightarrow “EDE”

Early Dark Energy

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$



At the critical redshift:


$$\ddot{\phi} + 3H\dot{\phi} + \frac{dV}{d\phi} = 0$$

- As time passes, $H(z)$ decreases, until at some critical redshift z_c :

$$3H\dot{\phi} \approx \frac{dV}{d\phi}$$

Early Dark Energy

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$



At the critical redshift:

$$\ddot{\phi} + 3H\dot{\phi} + \frac{dV}{d\phi} = 0$$

- As time passes, $H(z)$ decreases, until at some critical redshift z_c :

$$3H\dot{\phi} \approx \frac{dV}{d\phi}$$


- Let's approximate:

- $V(\phi) = m^2 f^2 (1 - \cos(\phi/f))^n \approx$

For $n = 1$ and small ϕ : $1 - \cos(\phi/f) \approx \frac{\phi^2}{2f^2}$

Early Dark Energy

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$



At the critical redshift:

$$\ddot{\phi} + 3H\dot{\phi} + \frac{dV}{d\phi} = 0$$

- As time passes, $H(z)$ decreases, until at some critical redshift z_c :

$$3H\phi \approx \frac{dV}{d\phi}$$


- Let's approximate:

$$V(\phi) = m^2 f^2 (1 - \cos(\phi/f))^n \approx \frac{m^2}{2} \phi^2$$

For $n = 1$ and small ϕ : $1 - \cos(\phi/f) \approx \frac{\phi^2}{2f^2}$

Early Dark Energy

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$




At the critical redshift:

$$3H\phi \approx \frac{dV}{d\phi}$$

Early Dark Energy

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$




At the critical redshift:

$$3H\dot{\phi} \approx \frac{dV}{d\phi}$$

- $V(\phi) \approx \frac{m^2}{2}\phi^2 \rightarrow \frac{dV}{d\phi} \approx m^2\phi$:

Early Dark Energy

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$



At the critical redshift:


$$3H\dot{\phi} \approx \frac{dV}{d\phi}$$

- $V(\phi) \approx \frac{m^2}{2}\phi^2 \rightarrow \frac{dV}{d\phi} \approx m^2\phi$:

$$3H \approx m^2$$

Early Dark Energy

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$



At the critical redshift:

$$3H\dot{\phi} \approx \frac{dV}{d\phi}$$

- $V(\phi) \approx \frac{m^2}{2}\phi^2 \rightarrow \frac{dV}{d\phi} \approx m^2\phi$:


$$3H \approx m^2$$

- Using this, one can determine the “mass” of the EDE field:

$$m \sim 10^{-28} \text{ eV}$$

Early Dark Energy

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$



At times after z_c :


$$3H\dot{\phi} \ll \frac{dV}{d\phi}$$

- Hence the Klein-Gordon eq. is:

$$\ddot{\phi} + \frac{dV}{d\phi} \approx 0$$

Early Dark Energy

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$



At times after z_c :

$$3H\dot{\phi} \ll \frac{dV}{d\phi}$$

- Hence the Klein-Gordon eq. is:


$$\ddot{\phi} + \frac{dV}{d\phi} \approx 0$$

- In our approximation:

$$V(\phi) \approx \frac{m^2}{2}\phi^2 \rightarrow \frac{dV}{d\phi} \approx m^2\phi$$

Early Dark Energy

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$



At times after z_c :

$$3H\dot{\phi} \ll \frac{dV}{d\phi}$$

- Hence the Klein-Gordon eq. is:

$$\ddot{\phi} + \frac{dV}{d\phi} \approx 0$$

- In our approximation:

$$V(\phi) \approx \frac{m^2}{2}\phi^2 \rightarrow \frac{dV}{d\phi} \approx m^2\phi$$

- Hence:


$$\ddot{\phi} + m^2\phi \approx 0$$

Early Dark Energy

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$


At times after z_c :

$$\ddot{\phi} + m^2\phi \approx 0$$



Early Dark Energy

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$



At times after z_c :


$$\ddot{\phi} + m^2\phi \approx 0$$

- This is an oscillation!

$$\phi(t) = \phi_0 \cos(mt)$$

Early Dark Energy

$$V(\phi) = V_0 [1 - \cos(\phi/f)]^n$$



At times after z_c :

$$\ddot{\phi} + m^2\phi \approx 0$$

- This is an oscillation!

$$\phi(t) = \phi_0 \cos(mt)$$

- One can now numerically show that the EOS is:

$$\langle w \rangle = \frac{n-1}{n+1}$$

For $n = 3$: $\langle w \rangle = \frac{1}{2}$

Early Dark Energy


Free “particle-physics” parameters:

- m mass ($V_0 = m^2 f^2$)
- f “decay constant”
- $\theta_i = \phi_i/f$ initial value of the field
- $(n = 3)$

Early Dark Energy

Free “particle-physics” parameters:


- m mass ($V_0 = m^2 f^2$)
- f “decay constant”
- $\theta_i = \phi_i/f$ initial value of the field
- $(n = 3)$



Free “phenomenological” parameters:


- f_{EDE} fraction of EDE at z_c
- z_c the critical redshift
- $\theta_i = \phi_i/f$ initial value of the field
- $(n = 3)$

Early Dark Energy



The critical redshift z_c determines when the field starts oscillating, hence when Ω_{EDE} peaks


Early Dark Energy



$f_{\text{EDE}} = \Omega_{\text{EDE}}(z_c)$: determines
the height of the peak

The critical redshift z_c determines
when the field starts oscillating,
hence when Ω_{EDE} peaks


Early Dark Energy



The initial value $\theta_i \in [0, \pi]$ determines how fast the field oscillates when it decays:

The closer to π the faster the oscillations

Early Dark Energy



The index $n = 1, 2, 3, \dots$ determines how fast the field decays:

The higher n , the faster the decay

$n = 3$ was shown to fit the data best

Recap Lecture 1

Introduction

- Friedmann equations describe (background) evolution of the components
- Hubble parameter
- distances
- CMB basics
- BAO
- How the CMB constrains H_0


$$\left(\frac{\dot{a}}{a}\right)^2 + \frac{k}{a^2} = \frac{8\pi G}{3}\rho + \frac{\Lambda}{3} \quad (\text{i})$$

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3}(\rho + 3p) + \frac{\Lambda}{3} \quad (\text{ii})$$

$$\dot{\rho} = -3(\rho + p)\frac{\dot{a}}{a} \quad (\text{iii})$$

Hubble tension and EDE

- Early- and late-universe solutions
$$\theta_s = \frac{r_s}{D_A(z^*)}$$
- EDE: scalar field in an expanding universe with a “ $1 - \cos$ ” potential
- EDE increases $H(z)$ before recombination and decays quickly after
- More about EDE: tomorrow



References

- Books: Dodelson&Schmidt “Modern Cosmology”, Weinberg: “Cosmology”, Huterer: “A course in cosmology”
- Komatsu “Physics of the Cosmic Microwave Background” (recorded lecture)
- Reviews about EDE:
 - Poulin et al. “The Ups and Downs of Early Dark Energy solutions to the Hubble tension: A review of models, hints and constraints circa 2023”
 - Kamionkowski&Riess: “The Hubble Tension and Early Dark Energy”