A decade of Polynomial Affine model of Gravity

José Perdiguero Gárate

March 6, 2024

Contents

Contents		
L	Introduction	1
2	The Polynomial Affine model of Gravity 2.1 Building the model	1 1 2 2
3	Analysis of the solutions	2
1	Final remarks	2
1	Introduction	
2	The Polynomial Affine model of Gravity	
2.	1 Building the model	
	$\begin{split} \hat{\Gamma}_{\alpha}{}^{\beta}{}_{\gamma} &= \hat{\Gamma}_{(\alpha}{}^{\beta}{}_{\gamma)} + \hat{\Gamma}_{[\alpha}{}^{\beta}{}_{\gamma]}, \\ &= \Gamma_{\alpha}{}^{\beta}{}_{\gamma} + \mathcal{B}_{\alpha}{}^{\beta}{}_{\gamma} + \delta^{\beta}_{[\gamma}\mathcal{A}_{\alpha]}, \end{split}$	(1)
S	$I = \int dV^{\alpha\beta\gamma\delta} \left[B_1 \mathcal{R}_{\mu\nu}{}^{\mu}{}_{\rho} \mathcal{B}_{\alpha}{}^{\nu}{}_{\beta} \mathcal{B}_{\gamma}{}^{\rho}{}_{\delta} + B_2 \mathcal{R}_{\alpha\beta}{}^{\mu}{}_{\rho} \mathcal{B}_{\gamma}{}^{\nu}{}_{\delta} \mathcal{B}_{\mu}{}^{\rho}{}_{\nu} + B_3 \mathcal{R}_{\mu\nu}{}^{\mu}{}_{\alpha} \mathcal{B}_{\beta}{}^{\nu}{}_{\gamma} \mathcal{B}_{\beta}{}^{\nu}{}_{\gamma} \mathcal{B}_{\beta}{}^{\nu}{}_{\gamma} \mathcal{B}_{\beta}{}^{\nu}{}_{\beta} \mathcal{B}_{\beta}{}^$	\mathcal{A}_{δ}

$$S = \int dV^{\alpha\beta\gamma\delta} \left[B_1 \mathcal{R}_{\mu\nu}{}^{\mu}{}_{\rho} \mathcal{B}_{\alpha}{}^{\nu}{}_{\beta} \mathcal{B}_{\gamma}{}^{\rho}{}_{\delta} + B_2 \mathcal{R}_{\alpha\beta}{}^{\mu}{}_{\rho} \mathcal{B}_{\gamma}{}^{\nu}{}_{\delta} \mathcal{B}_{\mu}{}^{\rho}{}_{\nu} + B_3 \mathcal{R}_{\mu\nu}{}^{\mu}{}_{\alpha} \mathcal{B}_{\beta}{}^{\nu}{}_{\gamma} \mathcal{A}_{\delta} \right. \\ + B_4 \mathcal{R}_{\alpha\beta}{}^{\sigma}{}_{\rho} \mathcal{B}_{\gamma}{}^{\rho}{}_{\delta} \mathcal{A}_{\sigma} + B_5 \mathcal{R}_{\alpha\beta}{}^{\rho}{}_{\rho} \mathcal{B}_{\gamma}{}^{\sigma}{}_{\delta} \mathcal{A}_{\sigma} + C_1 \mathcal{R}_{\mu\alpha}{}^{\mu}{}_{\nu} \nabla_{\beta} \mathcal{B}_{\gamma}{}^{\nu}{}_{\delta} \\ + C_2 \mathcal{R}_{\alpha\beta}{}^{\rho}{}_{\rho} \nabla_{\sigma} \mathcal{B}_{\gamma}{}^{\sigma}{}_{\delta} + D_1 \mathcal{B}_{\nu}{}^{\mu}{}_{\lambda} \mathcal{B}_{\mu}{}^{\nu}{}_{\alpha} \nabla_{\beta} \mathcal{R}_{\gamma}{}^{\lambda}{}_{\delta} + D_2 \mathcal{B}_{\alpha}{}^{\mu}{}_{\beta} \mathcal{B}_{\mu}{}^{\lambda}{}_{\nu} \nabla_{\lambda} \mathcal{B}_{\gamma}{}^{\nu}{}_{\delta} \\ + D_3 \mathcal{B}_{\alpha}{}^{\mu}{}_{\nu} \mathcal{B}_{\beta}{}^{\lambda}{}_{\gamma} \nabla_{\lambda} \mathcal{B}_{\mu}{}^{\nu}{}_{\delta} + D_4 \mathcal{B}_{\alpha}{}^{\lambda}{}_{\beta} \mathcal{B}_{\gamma}{}^{\sigma}{}_{\delta} \nabla_{\lambda} \mathcal{A}_{\sigma} + D_5 \mathcal{B}_{\alpha}{}^{\lambda}{}_{\beta} \mathcal{A}_{\sigma} \nabla_{\lambda} \mathcal{B}_{\gamma}{}^{\sigma}{}_{\delta} \\ + D_6 \mathcal{B}_{\alpha}{}^{\lambda}{}_{\beta} \mathcal{A}_{\gamma} \nabla_{\lambda} \mathcal{A}_{\delta} + D_7 \mathcal{B}_{\alpha}{}^{\lambda}{}_{\beta} \mathcal{A}_{\lambda} \nabla_{\gamma} \mathcal{A}_{\delta} + E_1 \nabla_{\rho} \mathcal{B}_{\alpha}{}^{\rho}{}_{\beta} \nabla_{\sigma} \mathcal{B}_{\gamma}{}^{\sigma}{}_{\delta} \\ + E_2 \nabla_{\rho} \mathcal{B}_{\alpha}{}^{\rho}{}_{\beta} \nabla_{\gamma} \mathcal{A}_{\delta} + F_1 \mathcal{B}_{\alpha}{}^{\mu}{}_{\beta} \mathcal{B}_{\gamma}{}^{\sigma}{}_{\delta} \mathcal{B}_{\mu}{}^{\lambda}{}_{\rho} \mathcal{B}_{\sigma}{}^{\rho}{}_{\lambda} + F_2 \mathcal{B}_{\alpha}{}^{\mu}{}_{\beta} \mathcal{B}_{\gamma}{}^{\nu}{}_{\lambda} \mathcal{B}_{\delta}{}^{\lambda}{}_{\rho} \mathcal{B}_{\mu}{}^{\rho}{}_{\nu} \\ + F_3 \mathcal{B}_{\nu}{}^{\mu}{}_{\lambda} \mathcal{B}_{\mu}{}^{\nu}{}_{\alpha} \mathcal{B}_{\beta}{}^{\lambda}{}_{\gamma} \mathcal{A}_{\delta} + F_4 \mathcal{B}_{\alpha}{}^{\mu}{}_{\beta} \mathcal{B}_{\gamma}{}^{\nu}{}_{\delta} \mathcal{A}_{\mu} \mathcal{A}_{\nu} \right].$$

$$(2)$$

- 2.2 Coupling a scalar field
- 2.3 Building ansatz
- 3 Analysis of the solutions
- 4 Final remarks

References