Graph the Secant and Cosecant Functions

$y = \csc x$ $y = \sec x$


To graph variations of or , we use the graph of the related reciprocal function for reference.

$y = \csc x = \dfrac{1}{\sin x}$ $\sin x = 0$ $\dfrac{1}{\sin x}$ $y = \csc x$

If , then is undefined, and the graph of has a vertical asymptote.

$x = n\pi$

This occurs when x is a multiple of π: for all integers n.



The relative maxima and minima from the sine function correspond to the relative minima and maxima for the cosecant function. Similarly with the secant function below.


x	0	$\frac{\pi}{2}$	π	$\frac{3\pi}{2}$	2π
sin x	0	1	0	−1	0
csc x	Und.	1	Und.	−1	Und.

$y = \sec x = \dfrac{1}{\cos x}$ $\cos x = 0$ $\dfrac{1}{\cos x}$ $y = \sec x$

If , then is undefined, and the graph of has a vertical asymptote.

$\dfrac{\pi}{2}$ $x = \dfrac{(2n+1)\pi}{2}$

This occurs when x is an odd multiple of : for all integers n.



x	0	$\frac{\pi}{2}$	π	$\frac{3\pi}{2}$	2π
cos x	1	0	−1	0	1
sec x	1	Und	−1	Und.	1

© McGraw Hill LLC. All rights reserved. No reproduction or distribution without the prior written consent of McGraw Hill LLC.

© McGraw Hill LLC. All rights reserved. No reproduction or distribution without the prior written consent of McGraw Hill LLC.

Graphs of the Cosecant and Secant Functions		
	$y = \csc x$	$y = \sec x$
Domain	$\{x \mid x \neq n\pi$ for all integers $n\}$	$\left\{x \mid x \neq \frac{(2n+1)\pi}{2}$ for all integers $n\right\}$
Range	$\{y \mid y \leq -1$ or $y \quad 1\}$	$\{y \mid y \leq -1$ or $y \quad 1\}$
Amplitude	None ($y = \csc x$ increases and decreases without bound)	None ($y = \sec x$ increases and decreases without bound)
Period	2π	2π
Vertical Asymptotes	$x = n\pi$ (multiples of π)	$x = \frac{(2n+1)\pi}{2}$ (odd multiples of $\frac{\pi}{2}$)
Symmetry	Origin (csc is an odd function.)	y-axis (sec is an even function.)

1.

a. Graph $y = -2\csc 3x$

amp $= |-2| = 2$

Period $= \frac{2\pi}{3}$

reflect across x-axis

$y = \sin 3x$ has x-intercepts

at multiples of $\frac{\pi}{3}$


$y = \sin 3x \rightarrow$ period $\frac{2\pi}{3}$

asymptotes of $\csc 3x$

$\frac{1}{4}\left(\frac{2\pi}{3}\right) = \frac{\pi}{6}$ start $(0, 0)$


(1) $0 + \frac{\pi}{6} = \frac{\pi}{6} \rightarrow \left(\frac{\pi}{6}, -2\right)$, etc.

$y = -2\sin 3x$



b. Graph $y = -2\csc 3x + 1$

add 1 $\uparrow +1$




© McGraw Hill LLC. All rights reserved. No reproduction or distribution without the prior written consent of McGraw Hill LLC.

Graph the Tangent and Cotangent Functions

$y = \tan x = \dfrac{\sin x}{\cos x}$

x-intercept where $\sin x = 0$; x is a multiple of π: at $x = n\pi$ for integer n.

vertical asymptote where $\cos x = 0$; x is an odd multiple of $\dfrac{\pi}{2}$: at $x = \dfrac{(2n+1)\pi}{2}$ for integer n.



The tangent function is an odd function and is symmetric with respect to the origin.


Each Quadrant I point has a mirror image in Quadrant III.

The period of the tangent function is π, so one complete period occurs between two consecutive vertical asymptotes.

$y = \cot x = \dfrac{\cos x}{\sin x}$

x-intercept where $\cos x = 0$; x is an odd multiple of $\dfrac{\pi}{2}$: at $x = \dfrac{(2n+1)\pi}{2}$ for integer n.

vertical asymptote where $\sin x = 0$; x is a multiple of π: at $x = n\pi$ for integer n.



The cotangent function is an odd function and is symmetric with respect to the origin.

Each Quadrant I point has a mirror image in Quadrant III.

The period of the cotangent function is π, so one complete period occurs between two consecutive vertical asymptotes.

© McGraw Hill LLC. All rights reserved. No reproduction or distribution without the prior written consent of McGraw Hill LLC.

© McGraw Hill LLC. All rights reserved. No reproduction or distribution without the prior written consent of McGraw Hill LLC.

Graphs of the Tangent and Cotangent Functions		
	$y = \tan x$	$y = \cot x$
Domain	$\left\{x \mid x \neq \frac{(2n+1)\pi}{2} \text{ for all integers } n\right\}$	$\{x \mid x \neq n\pi \text{ for all integers } n\}$
Range	All real numbers	All real numbers
Amplitude	None ($y = \tan x$ is unbounded.)	None ($y = \cot x$ is unbounded.)
Period	π	π
Vertical Asymptotes	$x = \frac{(2n+1)\pi}{2}$ (odd multiples of $\frac{\pi}{2}$)	$x = n\pi$ (multiples of π)
Symmetry	Origin (tan is an odd function.)	Origin (cot is an odd function.)

Graphing $y = A\tan(Bx - C) + D$ and $y = A\cot(Bx - C) + D$ with $B > 0$. (If $B < 0$, rewrite the equations.)

The period is $\frac{\pi}{B}$. The phase shift is $\frac{C}{B}$. The vertical shift is D.

$|A|$ is the vertical scaling factor. If $A < 0$, the graph is reflected across the x-axis.

1. Find an interval between two consecutive vertical asymptotes and graph the asymptotes as dashed lines.

   $y = A\tan(Bx - C)$ and $B > 0$, solve the inequality $-\frac{\pi}{2} < Bx - C < \frac{\pi}{2}$.
   $y = A\cot(Bx - C)$ and $B > 0$, solve the inequality $0 < Bx - C < \pi$.

2. Plot an x-intercept halfway between the asymptotes from Step 1.

3. Sketch the general shape of the "parent" function between the asymptotes.
   If more accuracy is desired, consider evaluating the function for x values midway between the x-intercept and the asymptotes found in step 1.

   For $y = \tan x$, the y-coordinates will be $-A$ and A.
   For $y = \cot x$, the y-coordinates will be A and $-A$.

---

2. Graph $y = \cot 4x$


© McGraw Hill LLC. All rights reserved. No reproduction or distribution without the prior written consent of McGraw Hill LLC.

$0 < 4x < \pi$

$0 < x < \frac{\pi}{4}$ (one cycle)

asymptotes $\Rightarrow x = 0,\ x = \frac{\pi}{4}$

$$\text{midpoint} = \frac{0 + \frac{\pi}{4}}{2} = \frac{\pi}{8}\,(x - \text{int})$$




3. Graph $y = 4\tan\left(2x + \frac{\pi}{3}\right)$

$$-\frac{\pi}{2} < 2x + \frac{\pi}{3} < \frac{\pi}{2}$$

$$-\frac{\pi}{2} - \frac{\pi}{3} < 2x < \frac{\pi}{2} - \frac{\pi}{3}$$

$$-\frac{5\pi}{6} < 2x < \frac{\pi}{6}$$

$$-\frac{5\pi}{12} < x < \frac{\pi}{12}$$



$$\frac{-\frac{5\pi}{12} + \frac{\pi}{12}}{2} = \frac{\frac{-4\pi}{12}}{2} = -\frac{\pi}{6}\,(x - \text{int})$$

midpoint

4. Graph $y = -\tan\frac{3\pi}{4}x + 2$

note


© McGraw Hill LLC. All rights reserved. No reproduction or distribution without the prior written consent of McGraw Hill LLC.


$$-\frac{\pi}{2} < \frac{3\pi}{4}x < \frac{\pi}{2}$$

$$-2\pi < 3\pi x < 2\pi$$

$$-\frac{2\pi}{3\pi} < x < \frac{2\pi}{3\pi}$$

$$-\frac{2}{3} < x < \frac{2}{3}$$





midpoint $\dfrac{-\frac{2}{3}+\frac{2}{3}}{2} = 0$ $(0, 0)$ ↑ move up 2 units from $y = -\tan\dfrac{3\pi}{4}x$

© McGraw Hill LLC. All rights reserved. No reproduction or distribution without the prior written consent of McGraw Hill LLC.