

Evaluate the Inverse Sine Function






Recall that a function must be one-to-one to have an inverse *function*. For $y = \sin x,$ any horizontal line taken between $-1 \le y \le 1$ intersects the graph infinitely many times. Therefore, $y = \sin x$ is not a one-to-one function.

Restrict the domain of $y = \sin x$ to the interval $-\frac{\pi}{2} \le x \le \frac{\pi}{2}.$

The graph of the restricted sine function *is* one-to-one and contains the entire range of y values $-1 \le y \le 1.$




The inverse of this restricted sine function is called the inverse sine function and is denoted by $\sin^{-1}$ or arcsin .

Inverse sine function

$y = \sin^{-1} x \quad \Leftrightarrow \quad \sin y = x$

$y = \arcsin x \quad \Leftrightarrow \quad \sin y = x \qquad$ where $-1 \le x \le 1$ and $-\frac{\pi}{2} \le y \le \frac{\pi}{2}$


To evaluate $y = \sin^{-1} x$ or $y = \arcsin x$ means to find an angle y between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$, inclusive, whose sine value is x.



Evaluate.

1. $\sin^{-1}\left(-\frac{1}{2}\right)$

2. $\arcsin\frac{\sqrt{3}}{2}$




3. $\sin^{-1}\frac{5\pi}{4}$


© McGraw Hill LLC. All rights reserved. No reproduction or distribution without the prior written consent of McGraw Hill LLC.

Evaluate the Inverse Cosine and Tangent Functions


Restricted $y = \cos x$



Restricted $y = \tan x$




$y = \cos^{-1} x \quad -1 \le x \le 1$ and $0 \le y \le \pi$




$y = \tan^{-1} x \quad x \in \sim$ and $-\frac{\pi}{2} < y < \frac{\pi}{2}$

Horizontal asymptotes: $y = -\frac{\pi}{2}, \; y = \frac{\pi}{2}$



Evaluate.

remember the signs



4. $\arctan 1$

5. $\cos^{-1}\left(-\frac{1}{2}\right)$

6. $\arccos \frac{\sqrt{3}}{2}$

7. $\tan^{-1} 0$

© McGraw Hill LLC. All rights reserved. No reproduction or distribution without the prior written consent of McGraw Hill LLC.

Approximate Inverse Trigonometric Functions on a Calculator

Use a calculator to approximate the function values in radians or degrees.

8. $\arccos\frac{3}{5}$

9. $\sin^{-1}(-0.457)$

10. $\arctan(-2.09)$

11. Use a calculator to approximate the degree measure (to 1 decimal place) of the angle θ where $\sin\theta = \frac{3}{8}$, $90° < \theta < 180°$.

Compose Trigonometric Functions and Inverse Trigonometric Functions

$\sin\left(\sin^{-1} x\right) = x$ for $-1 \le x \le 1$ $\qquad$ $\sin^{-1}(\sin x) = x$ for $-\frac{\pi}{2} \le x \le \frac{\pi}{2}$

$\tan\left(\tan^{-1} x\right) = x$ for $x \in \sim$ $\qquad$ $\tan^{-1}(\tan x) = x$ for $-\frac{\pi}{2} < x < \frac{\pi}{2}$

$\cos\left(\cos^{-1} x\right) = x$ for $-1 \le x \le 1$ $\qquad$ $\cos^{-1}(\cos x) = x$ for $0 \le x \le \pi$

Evaluate.

12.

a. $\arcsin\left(\sin\left(-\frac{\pi}{6}\right)\right)$

b. $\arcsin\left(\sin\frac{7\pi}{6}\right)$

13.

a. $\cos^{-1}\left(\cos\frac{7\pi}{6}\right)$

b. $\cos^{-1}\left(\cos\left(-\frac{1}{2}\right)\right)$

© McGraw Hill LLC. All rights reserved. No reproduction or distribution without the prior written consent of McGraw Hill LLC.

14. Find the exact value.

a. $\cos\left(\sin^{-1}\frac{40}{41}\right)$

b. $\cos\left(\sin^{-1}\left(-\frac{40}{41}\right)\right)$

15. Write $\tan\left(\arcsin\frac{2}{x}\right)$ for $x > 2$ as an algebraic expression.

Apply Inverse Trigonometric Functions


16. The Eiffel Tower is 1063 feet tall. A measurement is taken of the length of the shadow of the tower and is found to be 310 feet. Approximate the angle of elevation of the Sun to the nearest tenth of a degree.



© McGraw Hill LLC. All rights reserved. No reproduction or distribution without the prior written consent of McGraw Hill LLC.


Evaluate the Inverse Secant, Cosecant, and Cotangent Functions

Restricted $y = \sec x$




$y = \sec^{-1} x \quad \Leftrightarrow \quad \sec y = x$

where $|x| \geq 1$ and $0 \leq y < \frac{\pi}{2}$ or $\frac{\pi}{2} < y \leq \pi$.




Restricted $y = \csc x$




$y = \csc^{-1} x \quad \Leftrightarrow \quad \csc y = x$

where $|x| \geq 1$ and $-\frac{\pi}{2} \leq y < 0$ or $0 < y \leq \frac{\pi}{2}$.




Restricted $y = \cot x$



$y = \cot^{-1} x \quad \Leftrightarrow \quad \cot y = x$

where x is any real number and $0 < y < \pi$.



Evaluate.

17. $\sec^{-1} 2$

18. $\cot^{-1}\left(-\sqrt{3}\right)$

19. $\csc^{-1}\left(-\frac{2\sqrt{3}}{3}\right)$

20. Use a calculator to approximate $\csc^{-1}\frac{8}{5}$ in radians, rounded to 4 decimal places.

© McGraw Hill LLC. All rights reserved. No reproduction or distribution without the prior written consent of McGraw Hill LLC.