(Suggested maximum time: 5 minutes)

Place the following numbers in order, starting with the smallest:



1.4

 $\sqrt{2}$




Which one of the following is **not** a rational number? Explain your answer. **(b)**



3.142

$$\frac{22}{7}$$

 π



Find the values of $\frac{4n^2+1}{13}$, where $n \in \{17, 19, 21\}$. (c) (i)

n	$\frac{4n^2+1}{13}$
17	
19	
21	

State which **one** of your answers is a natural number, and explain your choice. (ii)

` ′				-						-		-	-				

- (a) John thinks that he has a method for finding **all** prime numbers. He says that if he uses the formulas in the table below, he will generate the prime numbers. He also says that these formulas will generate **only** the prime numbers.
 - (i) Complete the table.

p	6 <i>p</i> + 1	6p + 5
0	1	5
1		
2		
3		
4		
5		

(ii) Give two reasons why his method is not fully correct.

Reason 1:

Reason 2:

(b) The Swiss mathematician and physicist, Euler, first noticed (in 1772) that the expression $n^2 - n + 41$ gives a prime number for all positive integer values of n less than 41.

Explain why it does not give a prime number for n = 41.

(a) The sets A, B, and C are as follows:


$$A = \{2, 3, 4, 5, 6\}, B = \{2, 4, 6, 8, 10\}, \text{ and } C = \{1, 4, 8, 12, 14\}.$$

- (i) Complete the Venn diagram.
- (ii) List the elements of each of the following sets:


$$A \cap B =$$


$$B\setminus (A\cap C)=$$

$$(B \setminus A) \cup (B \setminus C) = \underline{\hspace{1cm}}$$




- (iii) Write down a null set, in terms of A, B, and C.
- **(b)** In a table quiz, 100 questions were asked. Team M answered 72 questions correctly. Team N answered 38 questions correctly.
 - (i) Find, with the aid of the Venn diagram, the minimum number of questions answered correctly by both teams.






(ii) Find, with the aid of the Venn diagram, the maximum number of questions answered correctly by both teams.



Junior Certificate 2014



Page	running
. age	

(Suggested maximum time: 10 minutes)

(a) Factorise fully $9a^2 - 6ab + 12ac - 8bc$.



(b) Factorise $9x^2 - 16y^2$.



(c) Use factors to simplify the following: $\frac{2x^2 + 4x}{2x^2 + x - 6}$.




Question 5

(Suggested maximum time: 5 minutes)

Solve the following inequality and show the solution on the number line.


$$-17 \le 1 - 3x < 13, x \in \mathbb{Z}$$




(Suggested maximum time: 10 minutes)


Below are three containers, labelled 1, 2, and 3.

Water is poured into each container at a constant rate, until it is full.







2




3

The three graphs, A, B, and C, show the height of the water, h, in the containers after time t.









(a) Write A, B, and C in the table below to match each container to its corresponding graph.

Container	1	2	3
Graph			

(b) Another container is shown below. Water is also poured into this container at a constant rate until it is full. Sketch the graph you would expect to get when plotting height (h) against time (t) for this container.





(Suggested maximum time: 10 minutes)


Last year Elena had a gross income of €36 960.

She had to pay Universal Social Charge (USC) and income tax on her gross income.

The rates and bands of USC are as follows.

Income band	Rate of USC
Up to €10 036	2%
Between €10 036 and €16 016	4%
Above €16 016	7%

(i) Find the amount of USC that was deducted from Elena's gross income last year.




(ii) The standard rate of income tax was 20% and the higher rate was 41%.

The standard rate cut-off point was €32 800.

Elena paid a total of €4965·60 income tax last year.

Find Elena's tax credits for the year.



(iii) Find Elena's total deduction (USC and income tax) as a percentage of her gross income. Give your answer correct to the nearest percent.




Question 8

(Suggested maximum time: 10 minutes)

The table shows the height, in metres, of a ball at various times after being kicked into the air.

(i) Is the pattern of heights in the table linear, quadratic, or exponential? Explain your answer.

Time (seconds)	0	0.5	1	1.5	2	2.5	3
Height (metres)	0.3	3.4	5.7	7.2	7.9	7.8	6.9



(ii) Estimate the height of the ball after 3.5 seconds.




(iii) Estimate the total time the ball spends in the air. Justify your answer.

Jack and Sarah are going on a school tour to England. They investigate how much different amounts of sterling (\pounds) will cost them in euro (€). They each go to a different bank.

Their results are shown in the table below.

Amount of sterling (£)	Cost in euro (€) for Jack	Cost in euro (€) for Sarah
20	33	24
40	56	48
60	79	72
80	102	96


(i) On the grid below, draw graphs to show how much the sterling will cost Jack and Sarah, for up to £80.




Sl	lope																			
_ E	xpla	natio	n:																	
XX 7 · 4	1		C	1	_			. 1	4 1	r 1					C					
Write sterli																		n a	moı	int c
	-1- ₀ .							5 01			15 9			J	10					
Write sterli																		en	amo	ount
																		ven	amo	ount
																		ven	amo	ount
																		ven	amo	ount
																		ven	amo	ount
																		ven	amo	ount
sterli	ng.	State	e cle	arly	the	mea	anin	g of	any	lette	ers y	ou us	se in	n you	ur fo	rmul	a.			
Using	ng.	State	rmu	las f	rom	mea	anin	g of	any v), o	or other	ers y	ou us	ind	the a	ur fo	rmul	a.			
sterli	ng.	State	rmu	las f	rom	mea	anin	g of	any v), o	or other	ers y	ou us	ind	the a	ur fo	rmul	a.			
Using	ng.	State	rmu	las f	rom	mea	anin	g of	any v), o	or other	ers y	ou us	ind	the a	ur fo	rmul	a.			
Using	ng.	State	rmu	las f	rom	mea	anin	g of	any v), o	or other	ers y	ou us	ind	the a	ur fo	rmul	a.			
Using	ng.	State	rmu	las f	rom	mea	anin	g of	any v), o	or other	ers y	ou us	ind	the a	ur fo	rmul	a.			
Using	ng.	State	rmu	las f	rom	mea	anin	g of	any v), o	or other	ers y	ou us	ind	the a	ur fo	rmul	a.			
Using	ng.	State	rmu	las f	rom	mea	anin	g of	any v), o	or other	ers y	ou us	ind	the a	ur fo	rmul	a.			
Using	ng.	State	rmu	las f	rom	mea	anin	g of	any v), o	or other	ers y	ou us	ind	the a	ur fo	rmul	a.			
Using	ng.	State	rmu	las f	rom	mea	anin	g of	any v), o	or other	ers y	ou us	ind	the a	ur fo	rmul	a.			


(Suggested maximum time: 10 minutes)


(a) The graphs of the functions $f(x) = x^2 + 2x - 3$ and $g(x) = -x^2 - 2x + 3$ are shown below. Identify each graph by writing f(x) or g(x) in the space provided below the graph.





(b) The graphs of the functions y = h(x) and y = k(x) are shown below.





Write down the roots of each function.

Hence, or otherwise, write down an equation for each function.

Roots of h(x):

Equation: h(x) =

Roots of k(x):

(Suggested maximum time: 10 minutes)

x is a real number.


One new number is formed by increasing x by 1.

A second new number is formed by decreasing x by 2.


(i) Write down each of these new numbers, in terms of x.

Increase <i>x</i> by 1:										
increase x by 1.										
Decrease <i>x</i> by 2:										
Decrease x by 2.										

(ii) The product of these two new numbers is 1. Use this information to write an equation in x.



(iii) Solve this equation to find the two possible values of x. Give each of your answers correct to 3 decimal places.



Question 12

(Suggested maximum time: 15 minutes)

(a) Simplify (6x-3)(2x-1).

(b) Simplify $(3x^3 - 2x^2 - 3x + 2) \div (x - 1)$.




(c) (i) Solve the simultaneous equations:

$$2x - 3y = 18$$

$$5x + 9y = -10.$$





(ii) Verify your answer to (c)(i).




(Suggested maximum time: 5 minutes)


(i) Use the diagram on the right to calculate the value of x. Give your answer in surd form.






(ii) Use the diagram below to calculate the value of y. Give your answer in surd form.





(iii) A rectangle with sides of length x and y is drawn using the values of x and y from parts (i) and (ii), as shown below.


Write the **perimeter** of this rectangle in the form $a\sqrt{2}$, where $a \in \mathbb{N}$.





(Suggested maximum time: 20 minutes)

(i) g is the function $g: x \mapsto x-1$, where $x \in \mathbb{R}$. Find the value of each of the following.



(ii) f is the function $f: x \mapsto 2x^2 - x - 6$, where $x \in \mathbb{R}$.

Using the same axes and scales, draw the graphs of the functions y = f(x) and y = g(x) in the domain $-2 \le x \le 3$.

