C-20

MEMBERSHIP

Stars in catalog	29
with Gaia GSP-Phot	9 (31%)
with radial velocity	8
Stream id (sID)	1

GEOMETRY · great-circle fit

Centroid RA, Dec	358.9, 10.3 deg
Centroid l, b	101.2, -50.2 deg
Pole RA, Dec	95.0, 30.9 deg
Length (φ1, 2–98%)	18.6 deg
Width σ(φ2), robust	0.08 deg

KINEMATICS

pmRA* med ± σ	-0.65 ± 0.23 mas/yr
pmDE med ± σ	1.52 ± 0.27 mas/yr
μφ1* med ± σ	-1.59 ± 0.33 mas/yr
μφ2 med ± σ	0.25 ± 0.14 mas/yr
Parallax (cat.) med	0.045 ± 0.209 mas
Parallax (wtd, ZP-corr.)	0.120 ± 0.027 mas
Distance	d = 8.3 ± 1.9 kpc
Physical length	2.70 kpc
RV median ± σ(MAD)	-115.1 ± 0.9 km/s
RV gradient dV/dφ1	-1.50 km/s/deg


PHOTOMETRY · POPULATIONS

G0 range (median)	13.9–19.9 (19.1)
(BP−RP)0 median	0.71
GSP-Phot [M/H] (n=9)	-1.79 ± 1.28
GSP-Phot Teff median	5390 K
RUWE > 1.4	0.0 %
PM clusters (largest)	2 (15 stars)


IBATA+2024 · TABLE 3

n catalog | paper	29 | 29
n_RV catalog | paper	8 | 8
<[Fe/H]> paper	-2.93
Stellar mass (paper)	1200 Msun
Pole offset vs paper	0.0 deg

Sky map · catalog members (n = 29) · size ∝ brightness · ring = has RV




Stream frame (great-circle fit) · length 18.6° · width σ 0.08°




C-20

Kinematics · proper motion · parallax · radial velocity


Proper motion (n = 29)




μφ1* vs φ1 · σ = 0.33




μφ2 vs φ1 · σ = 0.14




Parallax (catalog, n = 29)



Radial velocity vs φ1 (n = 8)




V_hel distribution




C-20


CMD · dereddened (n = 29)




HR diagram · ϖ>0 (n = 7/9)




G0 & GSP-Phot coverage




Star types from logg (n = 9)



Kiel diagram (n = 9)



Sky · GSP-Phot [M/H] (n = 9)




GSP-Phot (Gaia low-resolution BP/RP) parameters are prior-dominated for faint, distant halo stars (G ≥ 17): logg, [M/H] and ages are indicative only and typically biased toward dwarfs.


C-20

Metallicity · age · astrometric quality

[M/H] GSP-Phot (n = 9)



Teff vs [M/H] (n = 9)




FLAME age (n = 2)

FLAME ages for only 2 star(s)
(FLAME needs good parallaxes)


Age–metallicity (n = 2)

n < 3

RUWE (n = 9)




Parallax quality (n = 9)




C-20

PM clustering · HDBSCAN · 2 cluster(s) · noise 31%



Proper-motion directions (arrows) coloured by cluster



Cluster summary

id	n	pmRA	pmDE	σpmRA	[M/H]	n_RV
noise	9	-0.90	1.66	0.55	-1.38	0
C1	15	-0.56	1.43	0.16	-1.76	5
C2	5	-0.76	1.80	0.01	-3.09	3

C-20

Per-cluster proper-motion vector maps

