

Fjorm

MEMBERSHIP	
Stars in catalog	297
with Gaia GSP-Phot	120 (40%)
with radial velocity	29
Stream id (sID)	47
GEOMETRY · great-circle fit	
Centroid RA, Dec	216.5, 39.0 deg
Centroid l, b	56.0, 64.2 deg
Pole RA, Dec	103.0, 19.5 deg
Length (φ1, 2–98%)	99.5 deg
Width σ(φ2), robust	2.22 deg
KINEMATICS	
pmRA* med ± σ	-0.39 ± 2.51 mas/yr
pmDE med ± σ	4.63 ± 1.20 mas/yr
μφ1* med ± σ	4.49 ± 1.14 mas/yr
μφ2 med ± σ	2.35 ± 1.32 mas/yr
Parallax (cat.) med	0.159 ± 0.208 mas
Parallax (wtd, ZP-corr.)	0.207 ± 0.013 mas
Distance	d = 4.8 ± 0.3 kpc
Physical length	8.40 kpc
RV median ± σ(MAD)	-71.4 ± 29.5 km/s
RV gradient dV/dφ1	0.93 km/s/deg
PHOTOMETRY · POPULATIONS	
G0 range (median)	13.0–20.0 (19.1)
(BP−RP)0 median	0.74
GSP-Phot [M/H] (n=120)	-2.16 ± 0.79
GSP-Phot Teff median	6192 K
RUWE > 1.4	0.0 %
PM clusters (largest)	12 (99 stars)
IBATA+2024 · TABLE 3	
n catalog | paper	297 | 297
n_RV catalog | paper	29 | 29
<[Fe/H]> paper	-2.23
Stellar mass (paper)	1700 Msun
Pole offset vs paper	0.1 deg

Sky map · catalog members (n = 297) · size ∝ brightness · ring = has RV





Stream frame (great-circle fit) · length 99.5° · width σ 2.22°





Fjorm


Kinematics · proper motion · parallax · radial velocity
















Fjorm


CMD · dereddened (n = 297)




HR diagram · ϖ>0 (n = 111/120)




G0 & GSP-Phot coverage




Star types from logg (n = 120)



Kiel diagram (n = 120)



Sky · GSP-Phot [M/H] (n = 120)




GSP-Phot (Gaia low-resolution BP/RP) parameters are prior-dominated for faint, distant halo stars (G ≥ 17): logg, [M/H] and ages are indicative only and typically biased toward dwarfs.


Fjorm

Metallicity · age · astrometric quality


[M/H] GSP-Phot (n = 120)




Teff vs [M/H] (n = 120)




FLAME age (n = 7)




Age–metallicity (n = 7)



RUWE (n = 120)




Parallax quality (n = 120)




Fjorm

PM clustering · HDBSCAN · 12 cluster(s) · noise 22%



Proper-motion directions (arrows) coloured by cluster




Cluster summary

id	n	pmRA	pmDE	σpmRA	[M/H]	n_RV
noise	66	1.88	4.00	2.71	-2.13	6
C1	99	-1.53	4.81	0.63	-2.15	12
C2	18	-2.66	2.77	0.21	-2.08	1
C3	17	-0.67	6.01	0.39	-2.40	4
C4	14	1.20	5.83	0.24	-2.44	1
C5	14	2.30	4.61	0.26	-1.39	0
C6	13	0.21	5.76	0.09	-2.56	1
C7	13	1.65	5.34	0.30	-1.93	2
C8	12	3.97	2.95	0.11	-2.10	2


Fjorm

Per-cluster proper-motion vector maps


Cluster C1 · n = 99
pm = (-1.53, 4.81) mas/yr




Cluster C2 · n = 18
pm = (-2.66, 2.77) mas/yr




Cluster C3 · n = 17
pm = (-0.67, 6.01) mas/yr



Cluster C4 · n = 14
pm = (1.20, 5.83) mas/yr



Cluster C5 · n = 14
pm = (2.30, 4.61) mas/yr



Noise · n = 66

