

Gjoll

MEMBERSHIP	
Stars in catalog	607
with Gaia GSP-Phot	269 (44%)
with radial velocity	40
Stream id (sID)	34
GEOMETRY · great-circle fit	
Centroid RA, Dec	133.7, -46.1 deg
Centroid l, b	259.1, 8.4 deg
Pole RA, Dec	158.4, 43.5 deg
Length (φ1, 2–98%)	103.1 deg
Width σ(φ2), robust	0.71 deg
KINEMATICS	
pmRA* med ± σ	8.61 ± 1.07 mas/yr
pmDE med ± σ	-2.24 ± 0.88 mas/yr
μφ1* med ± σ	8.73 ± 1.14 mas/yr
μφ2 med ± σ	-1.38 ± 0.62 mas/yr
Parallax (cat.) med	0.227 ± 0.188 mas
Parallax (wtd, ZP-corr.)	0.235 ± 0.011 mas
Distance	d = 4.2 ± 0.2 kpc
Physical length	7.64 kpc
RV median ± σ(MAD)	75.9 ± 135.3 km/s
RV gradient dV/dφ1	7.97 km/s/deg
PHOTOMETRY · POPULATIONS	
G0 range (median)	12.4–20.0 (18.7)
(BP−RP)0 median	0.82
GSP-Phot [M/H] (n=269)	-2.20 ± 0.71
GSP-Phot Teff median	5386 K
RUWE > 1.4	0.4 %
PM clusters (largest)	15 (382 stars)
IBATA+2024 · TABLE 3	
n catalog | paper	607 | 607
n_RV catalog | paper	40 | 40
<[Fe/H]> paper	-1.63
Stellar mass (paper)	2100 Msun
Pole offset vs paper	0.1 deg

Sky map · catalog members (n = 607) · size ∝ brightness · ring = has RV





Stream frame (great-circle fit) · length 103.1° · width σ 0.71°





Gjoll


Kinematics · proper motion · parallax · radial velocity
















Gjoll


CMD · dereddened (n = 607)




HR diagram · ϖ>0 (n = 249/269)




G0 & GSP-Phot coverage




Star types from logg (n = 269)



Kiel diagram (n = 269)



Sky · GSP-Phot [M/H] (n = 269)




GSP-Phot (Gaia low-resolution BP/RP) parameters are prior-dominated for faint, distant halo stars (G ≳ 17): logg, [M/H] and ages are indicative only and typically biased toward dwarfs.


Gjoll

Metallicity · age · astrometric quality


[M/H] GSP-Phot (n = 269)




Teff vs [M/H] (n = 269)




FLAME age (n = 14)




Age–metallicity (n = 14)



RUWE (n = 269)




Parallax quality (n = 269)




Gjoll

PM clustering · HDBSCAN · 15 cluster(s) · noise 11%



Proper-motion directions (arrows) coloured by cluster




Cluster summary

id	n	pmRA	pmDE	σpmRA	[M/H]	n_RV
noise	65	20.58	-20.02	5.53	-1.92	7
C1	382	8.31	-1.96	0.50	-2.50	2
C2	29	23.10	-23.67	0.30	-2.11	7
C3	23	21.56	-22.49	0.51	-1.79	5
C4	15	22.54	-18.57	0.29	-1.73	3
C5	12	15.57	-16.12	1.19	-2.04	3
C6	12	19.63	-20.53	0.12	-1.98	0
C7	11	20.30	-21.37	0.32	-2.06	3
C8	10	6.94	-2.45	0.33	-2.17	0


Gjoll

Per-cluster proper-motion vector maps


Cluster C1 · n = 382
pm = (8.31, -1.96) mas/yr




Cluster C2 · n = 29
pm = (23.10, -23.67) mas/yr




Cluster C3 · n = 23
pm = (21.56, -22.49) mas/yr



Cluster C4 · n = 15
pm = (22.54, -18.57) mas/yr



Cluster C5 · n = 12
pm = (15.57, -16.12) mas/yr



Noise · n = 65

