

New-26

MEMBERSHIP	
Stars in catalog	43
with Gaia GSP-Phot	30 (70%)
with radial velocity	4
Stream id (sID)	82
GEOMETRY · great-circle fit	
Centroid RA, Dec	340.6, 29.8 deg
Centroid l, b	92.2, -25.1 deg
Pole RA, Dec	203.9, 51.7 deg
Length (φ1, 2–98%)	9.3 deg
Width σ(φ2), robust	0.42 deg
KINEMATICS	
pmRA* med ± σ	-3.51 ± 0.63 mas/yr
pmDE med ± σ	-5.02 ± 0.81 mas/yr
μφ1* med ± σ	-5.44 ± 0.99 mas/yr
μφ2 med ± σ	-2.97 ± 0.25 mas/yr
Parallax (cat.) med	0.224 ± 0.168 mas
Parallax (wtd, ZP-corr.)	0.224 ± 0.017 mas
Distance	d = 4.5 ± 0.3 kpc
Physical length	0.72 kpc
RV median ± σ(MAD)	-136.5 ± 91.1 km/s
RV gradient dV/dφ1	—
PHOTOMETRY · POPULATIONS	
G0 range (median)	14.9–19.8 (17.7)
(BP−RP)0 median	0.80
GSP-Phot [M/H] (n=30)	-0.99 ± 0.33
GSP-Phot Teff median	5502 K
RUWE > 1.4	0.0 %
PM clusters (largest)	4 (14 stars)
IBATA+2024 · TABLE 3	
n catalog | paper	43 | 43
n_RV catalog | paper	4 | 4
<[Fe/H]> paper	-0.17
Stellar mass (paper)	300 Msun
Pole offset vs paper	0.2 deg
Remark	Velocities unclear

Sky map · catalog members (n = 43) · size ∝ brightness · ring = has RV





Stream frame (great-circle fit) · length 9.3° · width σ 0.42°





New-26


Kinematics · proper motion · parallax · radial velocity
















New-26

Photometry · stellar types


CMD · dereddened (n = 43)




HR diagram · ϖ>0 (n = 30/30)




G0 & GSP-Phot coverage




Star types from logg (n = 30)



Kiel diagram (n = 30)



Sky · GSP-Phot [M/H] (n = 30)




GSP-Phot (Gaia low-resolution BP/RP) parameters are prior-dominated for faint, distant halo stars (G ≥ 17): logg, [M/H] and ages are indicative only and typically biased toward dwarfs.


New-26

Metallicity · age · astrometric quality

[M/H] GSP-Phot (n = 30)



Teff vs [M/H] (n = 30)




FLAME age (n = 2)

FLAME ages for only 2 star(s)
(FLAME needs good parallaxes)


Age–metallicity (n = 2)

n < 3

RUWE (n = 30)




Parallax quality (n = 30)




New-26

PM clustering · HDBSCAN · 4 cluster(s) · noise 12%



Proper-motion directions (arrows) coloured by cluster




Cluster summary

id	n	pmRA	pmDE	σpmRA	[M/H]	n_RV
noise	5	-3.62	-5.67	0.32	-0.91	0
C1	14	-3.51	-5.00	0.29	-0.93	1
C2	12	-4.47	-5.72	0.23	-1.01	1
C3	7	-2.48	-4.27	0.12	-1.02	2
C4	5	-3.28	-4.56	0.05	-0.99	0


New-26

Per-cluster proper-motion vector maps


Cluster C1 · n = 14
pm = (-3.51, -5.00) mas/yr




Cluster C2 · n = 12
pm = (-4.47, -5.72) mas/yr



Cluster C3 · n = 7
pm = (-2.48, -4.27) mas/yr



Cluster C4 · n = 5
pm = (-3.28, -4.56) mas/yr



Noise · n = 5

