

New-9

MEMBERSHIP	
Stars in catalog	10
with Gaia GSP-Phot	6 (60%)
with radial velocity	7
Stream id (sID)	29
GEOMETRY · great-circle fit	
Centroid RA, Dec	145.2, 30.3 deg
Centroid l, b	196.8, 48.4 deg
Pole RA, Dec	309.5, 58.7 deg
Length (φ1, 2–98%)	7.2 deg
Width σ(φ2), robust	0.82 deg
KINEMATICS	
pmRA* med ± σ	10.62 ± 0.70 mas/yr
pmDE med ± σ	-6.81 ± 0.18 mas/yr
μφ1* med ± σ	11.49 ± 0.53 mas/yr
μφ2 med ± σ	-5.19 ± 0.18 mas/yr
Parallax (cat.) med	0.254 ± 0.128 mas
Parallax (wtd, ZP-corr.)	0.228 ± 0.033 mas
Distance	d = 4.4 ± 0.6 kpc
Physical length	0.55 kpc
RV median ± σ(MAD)	120.5 ± 13.5 km/s
RV gradient dV/dφ1	0.20 km/s/deg
PHOTOMETRY · POPULATIONS	
G0 range (median)	16.2–19.7 (18.1)
(BP−RP)0 median	0.76
GSP-Phot [M/H] (n=6)	-1.59 ± 0.19
GSP-Phot Teff median	6218 K
RUWE > 1.4	0.0 %
PM clusters (largest)	none
IBATA+2024 · TABLE 3	
n catalog | paper	10 | 10
n_RV catalog | paper	7 | 7
<[Fe/H]> paper	-1.95
Stellar mass (paper)	38 Msun
Pole offset vs paper	4.4 deg

Sky map · catalog members (n = 10) · size ∝ brightness · ring = has RV





Stream frame (great-circle fit) · length 7.2° · width σ 0.82°





New-9


Kinematics · proper motion · parallax · radial velocity
















New-9


CMD · dereddened (n = 10)




HR diagram · ϖ>0 (n = 6/6)




G0 & GSP-Phot coverage




Star types from logg (n = 6)



Kiel diagram (n = 6)



Sky · GSP-Phot [M/H] (n = 6)




GSP-Phot (Gaia low-resolution BP/RP) parameters are prior-dominated for faint, distant halo stars (G ≥ 17): logg, [M/H] and ages are indicative only and typically biased toward dwarfs.


New-9

Metallicity · age · astrometric quality

[M/H] GSP-Phot (n = 6)



Teff vs [M/H] (n = 6)




FLAME age (n = 0)

FLAME ages for only 0 star(s)
(FLAME needs good parallaxes)


Age–metallicity (n = 0)

n < 3

RUWE (n = 6)




Parallax quality (n = 6)




New-9

PM clustering · HDBSCAN · 0 cluster(s) · noise 100%



Proper-motion directions (arrows) coloured by cluster



Cluster summary

id	n	pmRA	pmDE	σpmRA	[M/H]	n_RV
noise	10	10.62	-6.81	0.70	-1.59	7