

Slidr

MEMBERSHIP	
Stars in catalog	330
with Gaia GSP-Phot	207 (63%)
with radial velocity	34
Stream id (sID)	39
GEOMETRY · great-circle fit	
Centroid RA, Dec	165.7, 7.2 deg
Centroid l, b	245.9, 58.1 deg
Pole RA, Dec	272.1, 63.9 deg
Length (φ1, 2–98%)	30.8 deg
Width σ(φ2), robust	2.32 deg
KINEMATICS	
pmRA* med ± σ	-22.03 ± 2.85 mas/yr
pmDE med ± σ	-5.86 ± 1.06 mas/yr
μφ1* med ± σ	-17.46 ± 2.50 mas/yr
μφ2 med ± σ	-14.59 ± 1.71 mas/yr
Parallax (cat.) med	0.302 ± 0.205 mas
Parallax (wtd, ZP-corr.)	0.333 ± 0.012 mas
Distance	d = 3.0 ± 0.1 kpc
Physical length	1.61 kpc
RV median ± σ(MAD)	-92.0 ± 44.6 km/s
RV gradient dV/dφ1	-3.84 km/s/deg
PHOTOMETRY · POPULATIONS	
G0 range (median)	12.3–20.0 (18.5)
(BP−RP)0 median	0.92
GSP-Phot [M/H] (n=207)	-1.77 ± 0.39
GSP-Phot Teff median	5557 K
RUWE > 1.4	0.0 %
PM clusters (largest)	19 (27 stars)
IBATA+2024 · TABLE 3	
n catalog | paper	330 | 330
n_RV catalog | paper	34 | 34
<[Fe/H]> paper	-1.70
Stellar mass (paper)	840 Msun
Pole offset vs paper	0.2 deg

Sky map · catalog members (n = 330) · size ∝ brightness · ring = has RV





Stream frame (great-circle fit) · length 30.8° · width σ 2.32°





Slidr


Kinematics · proper motion · parallax · radial velocity
















Slidr


CMD · dereddened (n = 330)




HR diagram · ϖ>0 (n = 199/207)




G0 & GSP-Phot coverage




Star types from logg (n = 207)



Kiel diagram (n = 207)



Sky · GSP-Phot [M/H] (n = 207)




GSP-Phot (Gaia low-resolution BP/RP) parameters are prior-dominated for faint, distant halo stars (G ≳ 17): logg, [M/H] and ages are indicative only and typically biased toward dwarfs.


Slidr

Metallicity · age · astrometric quality


[M/H] GSP-Phot (n = 207)




Teff vs [M/H] (n = 207)




FLAME age (n = 9)




Age–metallicity (n = 9)



RUWE (n = 207)




Parallax quality (n = 207)




Slidr

PM clustering · HDBSCAN · 19 cluster(s) · noise 39%



Proper-motion directions (arrows) coloured by cluster




Cluster summary

id	n	pmRA	pmDE	σpmRA	[M/H]	n_RV
noise	129	-21.11	-5.69	3.53	-1.72	9
C1	27	-27.15	-8.15	0.65	-1.79	7
C2	22	-18.14	-4.63	0.37	-1.67	1
C3	15	-25.47	-6.68	0.49	-1.94	3
C4	13	-23.27	-6.25	0.23	-1.94	2
C5	11	-24.21	-6.78	0.15	-1.64	3
C6	11	-22.02	-6.90	0.36	-1.97	1
C7	11	-22.70	-6.11	0.15	-1.63	1
C8	11	-19.17	-4.58	0.20	-2.00	1


Slidr

Per-cluster proper-motion vector maps


Cluster C1 · n = 27
pm = (-27.15, -8.15) mas/yr




Cluster C2 · n = 22
pm = (-18.14, -4.63) mas/yr




Cluster C3 · n = 15
pm = (-25.47, -6.68) mas/yr



Cluster C4 · n = 13
pm = (-23.27, -6.25) mas/yr



Cluster C5 · n = 11
pm = (-24.21, -6.78) mas/yr



Noise · n = 129

