

Sylgr

MEMBERSHIP	
Stars in catalog	256
with Gaia GSP-Phot	146 (57%)
with radial velocity	33
Stream id (sID)	42
GEOMETRY · great-circle fit	
Centroid RA, Dec	176.9, -3.1 deg
Centroid l, b	274.6, 56.6 deg
Pole RA, Dec	93.1, 59.8 deg
Length (φ1, 2–98%)	24.7 deg
Width σ(φ2), robust	1.38 deg
KINEMATICS	
pmRA* med ± σ	-15.25 ± 4.13 mas/yr
pmDE med ± σ	-14.14 ± 4.25 mas/yr
μφ1* med ± σ	-20.22 ± 5.66 mas/yr
μφ2 med ± σ	-4.74 ± 1.67 mas/yr
Parallax (cat.) med	0.296 ± 0.241 mas
Parallax (wtd, ZP-corr.)	0.314 ± 0.014 mas
Distance	d = 3.2 ± 0.1 kpc
Physical length	1.37 kpc
RV median ± σ(MAD)	-204.3 ± 12.2 km/s
RV gradient dV/dφ1	-4.48 km/s/deg
PHOTOMETRY · POPULATIONS	
G0 range (median)	13.7–20.0 (18.7)
(BP−RP)0 median	0.89
GSP-Phot [M/H] (n=146)	-2.29 ± 0.78
GSP-Phot Teff median	5722 K
RUWE > 1.4	0.0 %
PM clusters (largest)	9 (105 stars)
IBATA+2024 · TABLE 3	
n catalog | paper	256 | 256
n_RV catalog | paper	33 | 33
<[Fe/H]> paper	-2.92
Stellar mass (paper)	700 Msun
Pole offset vs paper	0.6 deg
Remark	High-velocity scatter

Sky map · catalog members (n = 256) · size ∝ brightness · ring = has RV





Stream frame (great-circle fit) · length 24.7° · width σ 1.38°





Sylgr


Kinematics · proper motion · parallax · radial velocity
















Sylgr


CMD · dereddened (n = 256)




HR diagram · ϖ>0 (n = 135/146)




G0 & GSP-Phot coverage




Star types from logg (n = 146)



Kiel diagram (n = 146)



Sky · GSP-Phot [M/H] (n = 146)




GSP-Phot (Gaia low-resolution BP/RP) parameters are prior-dominated for faint, distant halo stars (G ≳ 17): logg, [M/H] and ages are indicative only and typically biased toward dwarfs.


Sylgr

Metallicity · age · astrometric quality


[M/H] GSP-Phot (n = 146)




Teff vs [M/H] (n = 146)




FLAME age (n = 4)




Age–metallicity (n = 4)



RUWE (n = 146)




Parallax quality (n = 146)




Sylgr

PM clustering · HDBSCAN · 9 cluster(s) · noise 18%



Proper-motion directions (arrows) coloured by cluster



Cluster summary

id	n	pmRA	pmDE	σpmRA	[M/H]	n_RV
noise	46	-17.62	-16.64	7.07	-2.00	0
C1	105	-13.31	-12.21	1.52	-2.60	25
C2	22	-16.05	-15.16	0.41	-2.15	3
C3	21	-18.08	-17.18	0.40	-2.34	0
C4	13	-21.42	-20.54	0.70	-2.60	0
C5	13	-20.07	-19.04	0.30	-2.43	1
C6	12	-16.92	-16.41	0.41	-2.62	0
C7	10	-19.19	-18.41	0.63	-2.12	0
C8	7	-19.73	-17.61	0.34	-2.29	0

Sylgr

Per-cluster proper-motion vector maps

