

Reider's Theorem, More and Less

January 2026

Let S be a smooth projective surface over $\mathbb{C}$.

1 Strategy

Going back to the ideas of [1], we will prove variants of Reider's theorem by finding suitable Bridgeland stability conditions.

Specifically, to show a line bundle $\omega_S(C)$ is d -very ample, we will get the vanishing

$$H^1(S, \mathcal{I}_Z \otimes \omega_S(C)) = 0,$$

as Z ranges over all length d subschemes. This cohomology group can be rewritten (with the help of Serre duality) as

$$\begin{aligned} \operatorname{Ext}_{\mathcal{O}_S}^1(\mathcal{O}_S(-C), \mathcal{I}_Z \otimes \omega_S) &= \operatorname{Ext}_{\mathcal{O}_S}^1(\mathcal{I}_Z, \mathcal{O}_S(-C))^\vee \\ &= \operatorname{Hom}_{D^b(S)}(\mathcal{I}_Z, \mathcal{O}_S(-C)[1])^\vee. \end{aligned}$$

To preclude any nonzero morphism $\mathcal{I}_Z \rightarrow \mathcal{O}_S(-C)[1]$ in the derived category, we will then find a Bridgeland stability condition with $\mathcal{I}_Z, \mathcal{O}_S(-C)[1]$ both (semi)stable but such that $\mathcal{I}_Z$ has greater slope.

2 Stability

2.1 Review

We use notation from [2], which includes a choice of smooth coordinates for a “slice” (of a chart) of the stability manifold. Specifically, we get a stability condition for each (x, y) inside the parabola $x > \frac{1}{2}y^2$, which is the data of a central charge $Z_{(x,y)}$ and a heart of a bounded t-structure $\mathcal{A}_{-y}$.

$$Z_{(x,y)}(E) = (-\operatorname{ch}_2(E), c_1(E) \cdot H) + (\operatorname{rk} E)(x, y)$$

$$\mathcal{A}_{-y} = \{E \in D^b(S) \mid -\mu_{\min}(\mathcal{H}^0(E)) < y \leq -\mu_{\max}(\mathcal{H}^{-1}(E)), \mathcal{H}^i(E) = 0 \text{ for } i \neq -1, 0\}$$

We recall here that as we increase y , we make the following trade:

- We *gain* H -semistable coherent sheaves with low H -slope, like $\mathcal{O}_S(-nC)$
- We *lose* shifts of H -semistable coherent sheaves with low H -slope, like $\mathcal{O}_S(-nC)[1]$.

In particular, an Mumford H -stable coherent sheaf $\mathcal{F}$ appears in the category when $y > -\mu(\mathcal{F})$.

Note that when $\mathcal{F}$ is *not* H -semistable, we have to wait a while after we lose $\mathcal{F}[1]$ before we gain $\mathcal{F}$. For instance, on $\mathbb{P}^2$ for $\mathcal{F} = \mathcal{O}_{\mathbb{P}^2}(-1) \oplus \mathcal{O}_{\mathbb{P}^2}(1)$, we have Harder-Narasimhan filtration

$$0 \leq \mathcal{O}_{\mathbb{P}^2}(1) \leq \mathcal{O}_{\mathbb{P}^2}(-1) \oplus \mathcal{O}_{\mathbb{P}^2}(1),$$

so has maximal H -slope $\mu_{\max}(\mathcal{F}) = 1$ and minimal slope $\mu_{\min}(\mathcal{F}) = -1$. Therefore, $\mathcal{F}[1]$ leaves the category once $y > -(1)$, but $\mathcal{F}$ only joins the category once $y > -(-1)$.

Also important is a description of (numerical) walls, which are lines through *points of indeterminacy* $I(E)$ of objects $E \in D^b(S)$, which are points $I(E) \in \mathbb{R}^2$ such that $Z_{I(E)}(E) = 0$. These have coordinates

$$I(E) = \left(\frac{\text{ch}_2 E}{\text{rk} E}, -\mu E \right).$$

Any H -semistable coherent sheaf $\mathcal{F}$ (or a shift, which has identical point of indeterminacy) has $I(\mathcal{F})$ outside or on the parabola—this is a version of the Bogomolov inequality. Alternatively, we can rephrase this argument as follows: if $\mathcal{F}$ is H -semistable, then $\mathcal{F} \in \mathcal{A}_{-\mu\mathcal{F}}$ so it is in the category at $I(\mathcal{F})$, so (x, y) cannot give a stability condition because the central charge sends an object to 0. Thus, (x, y) is outside the parabola.

When $\mathcal{F}$ fails to be semistable, like our example $\mathcal{F} = \mathcal{O}_{\mathbb{P}^2}(-1) \oplus \mathcal{O}_{\mathbb{P}^2}(1)$, we might have $I(\mathcal{F})$ inside the parabola, but it will occur at a y such that $\mathcal{F} \notin \mathcal{A}_{-y}$. For example, for this $\mathcal{F}$,

$$I(\mathcal{F}) = \left(\frac{\text{ch}_2(\mathcal{O}_{\mathbb{P}^2}(-1)) + \text{ch}_2(\mathcal{O}_{\mathbb{P}^2}(1))}{2}, \frac{-(-1) - (1)}{2} \right) = \left(\frac{1}{2}, 0 \right),$$

and indeed, $\mathcal{O}_{\mathbb{P}^2}(-1) \oplus \mathcal{O}_{\mathbb{P}^2}(1) \notin \mathcal{A}_0$ as discussed earlier.

2.2 $\mathcal{O}_S(-C)$ and Bogomolov-sharp Sheaves

Now, we analyze when $\mathcal{O}_S(-C)[1]$ is (semi)stable. First, we describe how, for an H -semistable torsion free coherent sheaf F , might F or $F[1]$ be destabilized. In particular, we need to know what a sub/quotient looks like in one of our hearts $\mathcal{A}_{-y}$.

Lemma 2.1. Let F be as above. Then,

- (1) If $K \leq F$ in $\mathcal{A}_{-y}$ with cokernel Q , then, by the long exact sequence in $\text{Coh}(S)$ cohomology,

$$0 \rightarrow \mathcal{H}^{-1}(K) \rightarrow 0 \rightarrow \mathcal{H}^{-1}(Q) \rightarrow \mathcal{H}^0(K) \rightarrow F \rightarrow \mathcal{H}^0(Q) \rightarrow 0$$

K is itself a sheaf.

- (2) If $F[1] \twoheadrightarrow Q$ is a surjection in $\mathcal{A}_{-y}$ with kernel K , then, by the long exact sequence in $\text{Coh}(S)$ cohomology,

$$0 \rightarrow \mathcal{H}^{-1}(K) \rightarrow F \rightarrow \mathcal{H}^{-1}(Q) \rightarrow \mathcal{H}^0(K) \rightarrow 0 \rightarrow \mathcal{H}^0(Q) \rightarrow 0,$$

$Q[-1]$ is itself a sheaf.

Lemma 2.2. Let F as before. Suppose there exists (x, y) such that F is semistabilized by some proper subobject $E \leq F$ in $\mathcal{A}_{-y}$. Then, $I(E)$ lies above $I(F)$, i.e. $-\mu(E) > -\mu(F)$, and $I(E)$ is on the same side of the parabola as $I(F)$.

Proof. Since $E, F \in \mathcal{A}_{-y}$, we know

$$y > -\mu(E), -\mu(F).$$

CONFUSED. By virtue of being in the category $E \in \mathcal{A}_{-y}$, we get $I(E)$ outside the parabola. $\square$

Now, we apply this lemma to get the stability of $\mathcal{O}_S(-C)[1]$ and friends, using the observation that

$$I(\mathcal{O}_S(-C)[1]) = \left(\frac{C^2}{2}, C^2 \right)$$

is exactly on the parabola. In this sense, $\mathcal{O}_S(-C)[1]$ is Bogomolov-sharp (the inequality became an equality), and we will show such Bogomolov-sharp object are stable in the following sense.

Proposition 2.3. Let $F \in \text{Coh}(S)$ be H -stable and torsion free such that $I(F)$ lands on the parabola $x = \frac{1}{2}y^2$. Then,

- (1) F is (x, y) semistable for all (x, y) in the parabola with $y > -\mu(F)$.
- (2) $F[1]$ is (x, y) semistable for all (x, y) in the parabola with $y \leq -\mu(F)$.

Proof. We show (1). Let (x, y) be inside the parabola with $y > -\mu(F)$, and suppose there exists a short exact sequence

$$0 \rightarrow E \rightarrow F \rightarrow F/E \rightarrow 0$$

in $\mathcal{A}_{-y}$ with E destabilizing F .

To get a contradiction, our strategy is as follows.

- Bound the positions of $I(E), I(F) \in \mathbb{R}^2$, forming a (numerical) wall for F .
- Argue that (x, y) is above this wall (i.e., on the “compact” side), since E destabilizes F .
- Contradict that (x, y) is inside the parabola.

By lemma 2.1, E is itself a coherent sheaf. Since $F \in \mathcal{A}_{-y}$, we need $y > -\mu(F)$. Since $E \in \mathcal{A}_{-y}$, we also need $y > -\mu_{\min}(E) > -\mu(E)$.

We want to argue that $-\mu(E) > -\mu(F)$ is actually

We show that E cannot destabilize. Our strategy is to show that any wall through $F[1]$ is disjoint from these (x, y) .

Suppose $F[1]$ is destabilized at some (x, y) in the parabola with $y \leq -\mu(F)$, witnessed by a short exact sequence

$$0 \rightarrow E \rightarrow F[1] \rightarrow F[1]/E \rightarrow 0$$

in $\mathcal{A}_{-y}$, such that $\nu_{(x,y)}(E) > \nu_{(x,y)}(F)$. In particular, for E to even be in the category, we need

$$-\mu(E) < y,$$

so $I(E)$ is strictly below both (x, y) and $I(F)$.

□

References

- [1] Daniele Arcara and Aaron Bertram. *{Reider}'s Theorem and {Thaddeus} Pairs Revisited*. Apr. 22, 2009. DOI: 10.48550/arXiv.0904.3500. arXiv: 0904.3500[math]. URL: <http://arxiv.org/abs/0904.3500> (visited on 02/07/2026).
- [2] Aaron Bertram et al. *Two New Extensions of Reider's Theorem on Algebraic Surfaces*. Apr. 23, 2026. DOI: 10.48550/arXiv.2604.21242. arXiv: 2604.21242[math]. URL: <http://arxiv.org/abs/2604.21242> (visited on 04/24/2026).