

Julia Sets

Theory and Algorithms

Josh Lipschultz & Ricky LeVan
MATH/CAAM 435



April 15, 2014

A quick recap of last week:

A quick recap of last week:

Definition

The stable set of a complex polynomial $P : \mathbb{C} \rightarrow \mathbb{C}$, denoted $S(P)$, is the complement of $J(C)$.

A quick recap of last week:

Definition

The stable set of a complex polynomial $P : \mathbb{C} \rightarrow \mathbb{C}$, denoted $S(P)$, is the complement of $J(C)$.

Definition

An orbit is bounded if there exists a K such that $|Q_C^{on}(z)| < K$ for all n . Otherwise the orbit is unbounded.

A quick recap of last week:

Definition

The stable set of a complex polynomial $P : \mathbb{C} \rightarrow \mathbb{C}$, denoted $S(P)$, is the complement of $J(C)$.

Definition

An orbit is bounded if there exists a K such that $|Q_c^n(z)| < K$ for all n . Otherwise the orbit is unbounded.

Remark

The points of S^1 are supersensitive under Q_0 . That is, any open ball around $z \in S^1$ has the property that $\bigcup_{n=0}^{\infty} Q_0^n(z) = \mathbb{C} \setminus \{p\}$ for at most one point p .

Definition

The Julia Set J_c is the boundary of the filled Julia set K_c . (The filled Julia set is the set of bounded points of Q_c .) We could alternatively define J_c as the closure of the set of repelling points of Q_c .

Definition

The Julia Set J_c is the boundary of the filled Julia set K_c . (The filled Julia set is the set of bounded points of Q_c .) We could alternatively define J_c as the closure of the set of repelling points of Q_c .

Remark

For the quadratic map $Q_0(z) = z^2$ we saw chaotic behavior only on S^1 by angle doubling ($\theta \rightarrow 2\theta$). We also saw that $|Q_0(z)| \rightarrow \infty$ for all $|z| > 1$ and $|Q_0(z)| \rightarrow 0$ for all $|z| < 1$.

Definition

The Julia Set J_c is the boundary of the filled Julia set K_c . (The filled Julia set is the set of bounded points of Q_c .) We could alternatively define J_c as the closure of the set of repelling points of Q_c .

Remark

For the quadratic map $Q_0(z) = z^2$ we saw chaotic behavior only on S^1 by angle doubling ($\theta \rightarrow 2\theta$). We also saw that $|Q_0(z)| \rightarrow \infty$ for all $|z| > 1$ and $|Q_0(z)| \rightarrow 0$ for all $|z| < 1$.

Remark

Finally, we learned that for $Q_{-2}(z) = z^2 - 2$, we have $J_2 = K_2 = [-2, 2]$, so any $z \in \mathbb{C} \setminus [-2, 2] \rightarrow \infty$ as we compose $Q_{-2}(z)$ infinitely many times.

- Goal: discuss and prove parts of:

Theorem

If $|c|$ is sufficiently large, Λ , the set of points whose entire forward orbits lie within the circle $|z| = |c|$, is a Cantor set on which Q_c is topologically conjugate to the shift map on two symbols. All points in $\mathbb{C} - \Lambda$ tend to ∞ under iteration of Q_c . Hence, $J_c = K_c$.

