

# Julia Sets

## Theory and Algorithms

Josh Lipschultz & Ricky LeVan  
*MATH/CAAM 435*



April 15, 2014

A quick recap of last week:

A quick recap of last week:

## Definition

*The stable set of a complex polynomial  $P : \mathbb{C} \rightarrow \mathbb{C}$ , denoted  $S(P)$ , is the complement of  $J(C)$ .*

A quick recap of last week:

## Definition

*The stable set of a complex polynomial  $P : \mathbb{C} \rightarrow \mathbb{C}$ , denoted  $S(P)$ , is the complement of  $J(C)$ .*

## Definition

*An orbit is bounded if there exists a  $K$  such that  $|Q_C^{on}(z)| < K$  for all  $n$ . Otherwise the orbit is unbounded.*

A quick recap of last week:

## Definition

*The stable set of a complex polynomial  $P : \mathbb{C} \rightarrow \mathbb{C}$ , denoted  $S(P)$ , is the complement of  $J(C)$ .*

## Definition

*An orbit is bounded if there exists a  $K$  such that  $|Q_c^n(z)| < K$  for all  $n$ . Otherwise the orbit is unbounded.*

## Remark

*The points of  $S^1$  are supersensitive under  $Q_0$ . That is, any open ball around  $z \in S^1$  has the property that  $\bigcup_{n=0}^{\infty} Q_0^n(z) = \mathbb{C} \setminus \{p\}$  for at most one point  $p$ .*

## Definition

*The Julia Set  $J_c$  is the boundary of the filled Julia set  $K_c$ . (The filled Julia set is the set of bounded points of  $Q_c$ .) We could alternatively define  $J_c$  as the closure of the set of repelling points of  $Q_c$ .*

## Definition

*The Julia Set  $J_c$  is the boundary of the filled Julia set  $K_c$ . (The filled Julia set is the set of bounded points of  $Q_c$ .) We could alternatively define  $J_c$  as the closure of the set of repelling points of  $Q_c$ .*

## Remark

*For the quadratic map  $Q_0(z) = z^2$  we saw chaotic behavior only on  $S^1$  by angle doubling ( $\theta \rightarrow 2\theta$ ). We also saw that  $|Q_0(z)| \rightarrow \infty$  for all  $|z| > 1$  and  $|Q_0(z)| \rightarrow 0$  for all  $|z| < 1$ .*

## Definition

*The Julia Set  $J_c$  is the boundary of the filled Julia set  $K_c$ . (The filled Julia set is the set of bounded points of  $Q_c$ .) We could alternatively define  $J_c$  as the closure of the set of repelling points of  $Q_c$ .*

## Remark

*For the quadratic map  $Q_0(z) = z^2$  we saw chaotic behavior only on  $S^1$  by angle doubling ( $\theta \rightarrow 2\theta$ ). We also saw that  $|Q_0(z)| \rightarrow \infty$  for all  $|z| > 1$  and  $|Q_0(z)| \rightarrow 0$  for all  $|z| < 1$ .*

## Remark

*Finally, we learned that for  $Q_{-2}(z) = z^2 - 2$ , we have  $J_2 = K_2 = [-2, 2]$ , so any  $z \in \mathbb{C} [-2, 2] \rightarrow \infty$  as we compose  $Q_{-2}(z)$  infinitely many times.*



- Goal: discuss and prove parts of:

## Theorem

*If  $|c|$  is sufficiently large,  $\Lambda$ , the set of points whose entire forward orbits lie within the circle  $|z| = |c|$ , is a Cantor set on which  $Q_c$  is topologically conjugate to the shift map on two symbols. All points in  $\mathbb{C} - \Lambda$  tend to  $\infty$  under iteration of  $Q_c$ . Hence,  $J_c = K_c$ .*







