

Homework Due 10/5/17 (7 problems):

Section 4.1 pages 169 - 170; 1, 6(b), 7(f), 9(a), 11, 12, 15

#1

Mark each statement True or False. Justify each answer.

- a. If (s_n) is a sequence and $s_i = s_j$ then $i = j$.

False.

Let: $(s_n) = \{1^n\}$

- b. If $s_n \rightarrow s$, then, for every $\epsilon > 0$, $\exists N \in \mathbb{N}$ st $n \geq N$ implies $|s_n - s| < \epsilon$.

True.

A sequence $\{s_n\}$ is said to **converge** to $s \in \mathbb{R}$ provided that $\forall \epsilon > 0$

$\exists N \in \mathbb{N} \leq n$ st

$$|s_n - s| < \epsilon$$

This is the definition of convergence, so this implies that $s_n \rightarrow s$

- c. If $s_n \rightarrow k$ and $t_n \rightarrow k$, then $s_n = t_n \forall n \in \mathbb{N}$.

False.

Let: $s_n = \sum_{i=0}^{\infty} \frac{1}{2^i}$, $t_n = 2 - \sum_{i=0}^{\infty} \frac{1}{2^i}$

- d. Every convergent sequence is bounded.

By Theorem 4.1.13, this is true.

6(b)

Definition 4.1.2

A sequence $\{s_n\}$ is said to **converge** to $s \in \mathbb{R}$ provided that $\forall \epsilon > 0$

$\exists N \in \mathbb{N}$ st

$n \geq N \longrightarrow |s_n - s| < \epsilon$

Using only definition 4.1.2, prove the following:

For $k > 0, k \in \mathbb{R}$, $\lim_{n \rightarrow \infty} \left(\frac{1}{n^k}\right) = 0$

Proof.

Let: $\{s_n\} = \frac{1}{n^k}, s = 0$

$$|s_n - s| = \left|\frac{1}{n^k} - 0\right| = \left|\frac{1}{n^k}\right|$$

Want to show: $\forall \epsilon > 0, \exists N \in \mathbb{N}$ st $n \geq N \longrightarrow \left|\frac{1}{n^k}\right| < \epsilon$

Let: $\epsilon > 0, N \in \mathbb{N}, k \in \mathbb{R} > 0$

Want to show: $\exists N \in \mathbb{N}$ st $\left|\frac{1}{N^k}\right| < \epsilon$

Let: $\left|\frac{1}{N^k}\right| < \epsilon$

$$\frac{1}{|N^k|} < \epsilon$$

$$\frac{1}{\epsilon} < |N^k|$$

$$|N^k| = N^k \text{ since } N \in \mathbb{N} \text{ and } k > 0 \text{ (1)}$$

$$\frac{1}{\epsilon} < N^k$$

$$\left(\frac{1}{\epsilon}\right)^{\frac{1}{k}} < N$$

If N is the ceiling of $\left(\frac{1}{\epsilon}\right)^{\frac{1}{k}} + 1$, then N exists.

Want to show: $\left|\frac{1}{(N+1)^k}\right| < \epsilon$

If we know that $\left|\frac{1}{N^k}\right| < \epsilon$,

then showing

$$\left|\frac{1}{(N+1)^k}\right| < \left|\frac{1}{N^k}\right|$$

shows

$$\left|\frac{1}{(N+1)^k}\right| < \epsilon$$

$$\left|\frac{1}{(N+1)^k}\right| < \left|\frac{1}{N^k}\right|$$

$$\frac{1}{|(N+1)^k|} < \frac{1}{|N^k|}$$

$$|N^k| < |(N+1)^k|$$

From (1),

$$|N^k| = N^k < |(N+1)^k| = (N+1)^k$$

$$N^k < (N+1)^k$$

This is true since $N \in \mathbb{N}$ and $k > 0$

So, $\left|\frac{1}{N^k}\right|$ decreases as N grows.

Since $\forall \epsilon > 0, \exists N \in \mathbb{N}$ st $n \geq N$ implies $\left|\frac{1}{n^k}\right| < \epsilon$,

$$\lim_{n \rightarrow \infty} \frac{1}{n^k} = 0$$

□

7(f)

Using any of the results in this section (4.1), prove the following:

If $|x| < 1$, then $\lim_{n \rightarrow \infty} x^n = 0$

Proof.

$|x| < 1$ implies $0 \leq |x| < 1$ **(1)**

Let: $s_n = x^n$, $s = 0$

Want to show: $\forall \epsilon > 0, \exists N \in \mathbb{N}$ st $n \geq N$ implies $|s_n - s| < \epsilon$

Let: $\epsilon > 0$

$|s_n - s| < \epsilon = |x^n| < \epsilon$

Want to show: $\exists N \in \mathbb{N}$ st $|x^N| < \epsilon$

$|x^N| < \epsilon$

$||x^N|| < |\epsilon|$

We know that because of **(1)** and because $N \in \mathbb{N}$,

$|x^{N+1}| < |x^N|$

We also know that $\epsilon > 0$

So, $0 < |x^{N+k}| < \dots < |x^{N+1}| < |x^N|$ where $k \in \mathbb{N}$

□

9(a)

For each of the following, prove or give a counter example:

If (s_n) converges to s , then $(|s_n|)$ converges to $|s|$.

Proof.

If s_n converges to s , then by definition,

$\forall \epsilon > 0, \exists N \in \mathbb{N}$ st $N \leq n$ implies $|s_n - s| < \epsilon$

Want to show: $||s_n| - |s|| < \epsilon$

Case 1: s_n and s are the same sign.

$||s_n| - |s|| = |s_n - s|$

Therefore, $N \leq n$ implies $||s_n| - |s|| < \epsilon$

If we let $s_n = |s_n|$ and $|s| = s$, then $|s_n|$ converges to $|s|$.

Case 2: s_n and s are different signs.

$||s_n| - |s|| \leq |s_n - s| < \epsilon$

$||s_n| - |s|| < \epsilon$

Therefore, $N \leq n$ implies $||s_n| - |s|| < \epsilon$

If we let $s_n = |s_n|$ and $|s| = s$, then $|s_n|$ converges to $|s|$.

Hence, result.

□

11

Given the sequence (s_n) , $k \in \mathbb{N}$, let (t_n) be the sequence defined by $t_n = s_{n+k}$. That is, the terms in (t_n) are the same as that of the terms in (s_n) after the first k terms have been skipped. Prove that (t_n) converges iff (s_n) converges, and if they converge, show that $\lim t_n = \lim s_n$. Thus, the convergence of a sequence is not affected by omitting (or changing) a finite number of terms.

Proof.

\longrightarrow

(t_n) converges $\longrightarrow (s_n)$ converges

If t_n converges, then by definition,

$\forall \epsilon > 0, \exists N \in \mathbb{N}$ st $n \geq N$ implies $|t_n - L| < \epsilon$

Since $t_n = s_{n+k}$,

we know that s_{n+k} converges.

Let: $n_1 = n + k$

So,

$\forall \epsilon > 0, \exists N \in \mathbb{N}$ st $n_1 \geq N$ implies $|s_{n_1} - L| < \epsilon$

$\forall \epsilon > 0, \exists N \in \mathbb{N}$ st $n + k \geq N$ implies $|s_{n+k} - L| < \epsilon$

$\forall \epsilon > 0, \exists N \in \mathbb{N}$ st $n \geq N - k$ implies $|s_{n+k} - L| < \epsilon$

Notice that $N - k \in \mathbb{N}$. Let's call it N_1

$\forall \epsilon > 0, \exists N_1 \in \mathbb{N}$ st $n \geq N_1$ implies $|s_{n+k} - L| < \epsilon$

Since there is still a natural number N_1 st $n \geq N_1$ implies $|s_{n+k} - L| < \epsilon$,

If t_n converges, then s_n converges.

\longleftarrow

(s_n) converges $\longrightarrow (t_n)$ converges

If s_n converges, then by definition,

$\forall \epsilon > 0, \exists N \in \mathbb{N}$ st $n \geq N$ implies $|s_n - L| < \epsilon$

Since $t_n = s_{n+k}$, $t_{n-k} = s_n$

So since s_n converges, t_{n-k} converges.

If we let $n_1 = n - k$,

$\forall \epsilon > 0, \exists N \in \mathbb{N}$ st $n_1 \geq N$ implies $|t_{n_1} - L| < \epsilon$

$\forall \epsilon > 0, \exists N \in \mathbb{N}$ st $n - k \geq N$ implies $|t_{n-k} - L| < \epsilon$

$\forall \epsilon > 0, \exists N \in \mathbb{N}$ st $n \geq (N + k)$ implies $|t_{n-k} - L| < \epsilon$

Notice that $N + k \in \mathbb{N}$. Let's call it N_1

$\forall \epsilon > 0, \exists N_1 \in \mathbb{N}$ st $n \geq N_1$ implies $|t_{n-k} - L| < \epsilon$

Since there is still a natural number N_1 st $n \geq N_1$ implies $|t_{n-k} - L| < \epsilon$,

If s_n converges, then t_n converges.

□

12

a. Assume that $\lim s_n = 0$. If (t_n) is a bounded sequence, prove that $\lim(s_n t_n) = 0$.

If $\lim s_n = 0$, then by definition,

$\forall \epsilon > 0, \exists N \in \mathbb{N}$ st if $n \geq N$, then $|s_n - 0| < \epsilon$

If t_n is a bounded sequence, then $\forall n \in \mathbb{N}$, $a \leq t_n \leq b$, where $a, b \in \mathbb{R}$

We know that t_n will always be between two constants a and b , so let's let $t_n = c$, where $a \leq c \leq b$.

Since s_n converges,

$\forall \epsilon > 0, \exists N \in \mathbb{N}$ st if $n \geq N$, then $|s_n - 0| < \epsilon$

can be simplified to

$\forall \epsilon > 0, \exists N \in \mathbb{N}$ st if $n \geq N$, then $|s_n| < \epsilon$

Want to show: $\lim(s_n t_n) = 0$

$\lim(s_n t_n) = 0$ if

$\forall \epsilon > 0, \exists N \in \mathbb{N}$ st if $n \geq N$, then $|s_n t_n| < \epsilon$

Since we let $t_n = c$, some bounded real number, this is equivalent to

$\forall \epsilon > 0, \exists N \in \mathbb{N}$ st if $n \geq N$, then $|cs_n| < \epsilon$

$\forall \epsilon > 0, \exists N \in \mathbb{N}$ st if $n \geq N$, then $|s_n| < |c|\epsilon$

$\forall \epsilon > 0, \exists N \in \mathbb{N}$ st if $n \geq N$, then $|s_n| < \epsilon_1$

which is equivalent to

$\forall \epsilon > 0, \exists N \in \mathbb{N}$ st if $n \geq N$, then $|s_n| < \epsilon$

Hence, result.

- b. Show by example that the boundedness of (t_n) is a necessary condition in part (a).

If $\lim s_n = 0$, then by definition,

$\forall \epsilon > 0, \exists N \in \mathbb{N}$ st if $n \geq N$, then $|s_n - 0| < \epsilon$

$\forall \epsilon > 0, \exists N \in \mathbb{N}$ st if $n \geq N$, then $|s_n| < \epsilon$

However, if we let t_n be unbounded (i.e. let $t_n = e^n$), this doesn't work. See below:

$s_n t_n$ is bounded if

$\forall \epsilon > 0, \exists N \in \mathbb{N}$ st if $n \geq N$, then $|s_n t_n| < \epsilon$

Suppose: $s_n = \frac{1}{n}$

Then $s_n t_n = \frac{e^n}{n}$

Since e^n grows faster than $\frac{1}{n}$, $s_n t_n$ grows overall as n approaches infinity.

Hence, the boundedness of t_n is necessary.

15

- Prove that x is an accumulation point of a set S iff \exists a sequence (s_n) of points in $S \setminus \{x\}$ st (s_n) converges to x .
- Prove that a set S is closed iff, whenever (s_n) is a convergent sequence of points in S , it follows that $\lim s_n$ is in S .