

Homework Due 10/12/17: (13 problems) Section 4.2 pages 177 - 178; 1, 2, 4, 5(a)(c)(e)(g)(i)(k), 9, 10, 17, 18 (for 5(i) define t_n to be $1/\sqrt{n}$, and then show that $1/\sqrt{n} \rightarrow 0$)

Corollary 4.2.5

If $\{t_n\}$ converges to t and $t_n \geq 0 \forall n \in \mathbb{N}$, then $t \geq 0$

Example 4.2.6

If $\{t_n\}$ converges to t and $t_n \geq 0 \forall n \in \mathbb{N}$, then

$$\lim_{n \rightarrow \infty} \sqrt{t_n} = \sqrt{t}$$

Proof.

Side Note

For $\epsilon > 0$, $\exists N \in \mathbb{N}$ st
 $|\sqrt{t_n} - \sqrt{t}| < \epsilon \forall n \geq N$

Notice that $\lim_{n \rightarrow \infty} t_n = t$, $t \geq 0$

Case **(i)**:

$$\begin{aligned} |\sqrt{t_n} - \sqrt{t}| &= \frac{|(\sqrt{t_n} - \sqrt{t})(\sqrt{t_n} + \sqrt{t})|}{|\sqrt{t_n} + \sqrt{t}|} \\ &= \frac{|t_n - t|}{\sqrt{t_n} + \sqrt{t}} \\ &\leq \frac{|t_n - t|}{\sqrt{t}} \\ &= \left(\frac{1}{\sqrt{t}}\right)|t_n - t| \end{aligned}$$

For $\epsilon > 0$, $\exists N \in \mathbb{N}$ st
 $|t_n - t| < \sqrt{t} \times \epsilon$, $\forall n \geq N$ **(2)**

Side Note

$$\begin{aligned} \sqrt{t} + \sqrt{t_n} &\geq \sqrt{t} \\ \frac{1}{\sqrt{t} + \sqrt{t_n}} &\leq \frac{1}{\sqrt{t}} \\ \sqrt{t_n} &\geq 0 \\ \sqrt{t} &> 0 \end{aligned}$$

So, $\sqrt{t_n} + \sqrt{t} > 0 \forall n \in \mathbb{N}$

From **(1)** and **(2)**,

$$|\sqrt{t_n} - \sqrt{t}| \leq \frac{|t_n - t|}{\sqrt{t}} < \frac{\sqrt{t} \times \epsilon}{\sqrt{t}} = \epsilon, \forall n \geq N$$

Hence, result in this case.

Side Note

If

$$|s_n - s| \leq k|a_n| \quad \forall n \geq N$$

and if

$$\lim_{n \rightarrow \infty} a_n = 0,$$

$$\text{then } \lim_{n \rightarrow \infty} s_n = s$$

Case (ii): $t = 0$

Then, for $\epsilon > 0$, $\exists N \in \mathbb{N}$ st

$$t_n = |t_n - 0| < \epsilon^2, \quad \forall n \geq N$$

Thus, $\sqrt{t_n} < \epsilon, \quad \forall n \geq N$

In other words,

$$|\sqrt{t_n} - 0| < \epsilon, \quad \forall n \geq N$$

$$\text{So, } \lim_{n \rightarrow \infty} \sqrt{t_n} = 0 = \sqrt{t}$$

Hence, result.

□

Theorem 4.2.7 - "The Ratio Test"

Suppose that $\{s_n\}$ is a sequence of **positive** terms (i.e. $s_n > 0, \forall n \in \mathbb{N}$) and $\lim_{n \rightarrow \infty} \frac{s_{n+1}}{s_n} = L$.

If $L < 1$, then $\lim_{n \rightarrow \infty} s_n = 0$

Proof.

$$\text{For } \epsilon = \frac{1-L}{2} > 0,$$

$$\exists N \in \mathbb{N} \text{ st}$$

$$\left| \frac{s_{n+1}}{s_n} - L \right| < \frac{1-L}{2}, \quad \forall n \geq N$$

So,

$$\frac{s_{n+1}}{s_n} = \left| \frac{s_{n+1}}{s_n} \right| = \left| \left(\frac{s_{n+1}}{s_n} - L \right) + L \right| \leq \left| \frac{s_{n+1}}{s_n} - L \right| + |L| < \left(\frac{1-L}{2} \right) + L = \frac{1+L}{2} = \frac{1}{2} + \frac{L}{2} < \frac{1}{2} + \frac{1}{2} \text{ (which is 1)}$$

$$\text{Define } c = \frac{1+L}{2}$$

Then,

$$s_n \times \frac{s_{n+1}}{s_n} < c s_n, \quad \forall n \geq N \text{ where } c < 1$$

$$\text{So, } s_{n+1} < c s_n, \quad \forall n \geq N$$

Now

$$s_{N+1} < c^1 s_N$$

$$s_{N+2} < c s_{N+1} < c^2 s_N$$

$$s_{N+3} < c s_{N+2} < c^3 s_N, \text{ etc.}$$

So,

$$s_{N+k} \leq c^k s_N, \quad \forall k \in \mathbb{N} \cup \{0\}$$

Thus,

$$s_m \leq c^{m-N} s_N \quad \forall m \geq N$$

$$s_m \leq c^m \frac{s_N}{c^N} \quad \forall m \geq N$$

$$N + k = m$$

$$k = m - N$$

$$|s_m - 0| = \left(\frac{s_N}{c^N}\right) \quad (1)$$

Side Note

Theorem 4.1.8

If $|s_m - s| \leq k|a_m|$ and

$$\lim_{m \rightarrow \infty} a_m = 0$$

then $\lim s_m = s$ Also, recall HW 5 7(f): If $|x| < 1$, then $\lim_{n \rightarrow \infty} (x^n) = 0$ From (1), it follows by Example 7(f) pg 170 and Theorem 4.1.8, that $\lim_{n \rightarrow \infty} s_n = 0$ **Example 5(g):** $s_n = \frac{1-n}{2^n} = \frac{1}{2^n} - \frac{n}{2^n}$ (or, $v_n - u_n$)Suppose $u_n = \frac{n}{2^n} > 0 \forall n \in \mathbb{N}$,

$$\frac{t_{n+1}}{t_n} = \frac{n+1}{2^{n+1}} \times \frac{2^n}{n} = \frac{1}{2} \frac{n(1+\frac{1}{n})}{n} = \frac{1}{2} \frac{1+\frac{1}{n}}{1} = \frac{1+\frac{1}{n}}{2} = \frac{1}{2} + \frac{1}{2n}$$

Which approaches $\frac{1}{2}$ as $n \rightarrow \infty$ **Definition 4.2.9****Infinite Limits:**A sequence $\{s_n\}$ is said to **diverge** to ∞ , written as $\lim_{n \rightarrow \infty} s_n = \infty$, provided that $\forall M \in \mathbb{R}, \exists N \in \mathbb{N}$ st $s_n > M, \forall n \geq N$ (i.e. $s_n = (-1)^n$ Similarly, $\{s_n\}$ diverges to $-\infty$, written as $\lim_{n \rightarrow \infty} s_n = -\infty$, if, provided thatfor every $M \in \mathbb{R}, \exists N(M) \in \mathbb{N}$ st $s_n < M, \forall n \geq N$ **Theorem 4.2.12**Suppose that $\{s_n\}, \{t_n\}$ are sequences st $s_n \leq t_n \forall n \in \mathbb{N}$ a. If $\lim_{n \rightarrow \infty} s_n = \infty$, then $\lim_{n \rightarrow \infty} t_n = \infty$ b. If $\lim_{n \rightarrow \infty} s_n = -\infty$, then $\lim_{n \rightarrow \infty} t_n = -\infty$

On the homework, the proof, using the definition, about "one comment away" from being done.

Theorem 4.2.13**Let:** $\{s_n\}$ be a sequence of **positive** numbersThen $\lim_{n \rightarrow \infty} s_n = \infty$ iff $\lim_{n \rightarrow \infty} \frac{1}{s_n} = 0$

Proof:

 \longrightarrow Suppose that $\lim_{n \rightarrow \infty} s_n = \infty$ **Want to show:** $\lim_{n \rightarrow \infty} \frac{1}{s_n} = 0$

Side Note

 $\forall \epsilon > 0, \exists N \in \mathbb{N}$ st

$$\left| \frac{1}{s_n} - 0 \right| = \frac{1}{s_n} < \epsilon$$

(which implies that $s_n > \frac{1}{\epsilon}$)

$$\forall n \geq N$$

$$\forall \epsilon > 0, \exists N \in \mathbb{N} \text{ st}$$

$$s_n > \frac{1}{\epsilon}, \forall n \geq N$$

Hence,

$$|\frac{1}{s_n} - 0| = \frac{1}{s_n} < \epsilon, \forall n \geq N$$

$$\text{Which shows that } \lim_{n \rightarrow \infty} \frac{1}{s_n} = 0$$

←

$$\text{Conversely, assume that } \lim_{n \rightarrow \infty} \frac{1}{s_n} = 0$$

$$\textbf{Want to show: } \lim_{n \rightarrow \infty} s_n = \infty$$

Side Note

$$\text{For } M \in \mathbb{R}, \exists N \in \mathbb{N} \text{ st}$$

$$\frac{1}{s_n} < \frac{1}{M}$$

$$s_n > M$$

$$\forall n \geq N$$

$$\textbf{Let: } M \in \mathbb{R}, M > 0$$

$$\text{Then } \exists N(M) \in \mathbb{N} \text{ st}$$

$$\frac{1}{s_n} = |\frac{1}{s_n} - 0| < \frac{1}{M} \forall n \geq N$$

$$\text{Hence, } s_n > M, \forall n \geq N.$$

Hence, result.

□