

Pinned group summary: SO-n6-q2

Pinning-Sympy automated output

July 14, 2026

Contents

1	Basic group attributes	3
1.1	Symmetric bilinear form	3
1.2	Lie algebra	3
1.3	Torus	3
2	Root system	4
2.1	Dynkin diagram	4
2.2	Cartan matrix	4
2.3	Roots	4
2.4	Coroots	4
2.5	Linear combinations of roots	4
3	Root spaces	6
3.1	Dimensions	6
3.2	Root space table	6
4	Root subgroups	8
4.1	Root subgroup table	8
4.2	Homomorphism property	9
5	Commutators	10
5.1	Commutator identities	11
6	Weyl group	12
6.1	Weyl group elements	12
6.2	Weyl group conjugation coefficients	13
6.3	Weyl group conjugation identities	14

1 Basic group attributes

Group	SO-n6-q2
Matrix size	6×6
Rank	2
Defining equations	$X^T B X = B$ and $\det(X) = 1$

1.1 Symmetric bilinear form

Dimension	6
Witt index	2
Primitive element	N/A
Epsilon	N/A

Matrix:
$$\begin{bmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & c_0 & 0 \\ 0 & 0 & 0 & 0 & 0 & c_1 \end{bmatrix}$$

1.2 Lie algebra

The defining equations for the Lie algebra are: $X^T B + B X = 0$ and $\text{tr}(X) = 0$
A generic element of the Lie algebra is:

$$\begin{bmatrix} x_{00} & x_{01} & 0 & -x_{12} & -c_0 x_{42} & -c_1 x_{52} \\ x_{10} & x_{11} & x_{12} & 0 & -c_0 x_{43} & -c_1 x_{53} \\ 0 & -x_{30} & -x_{00} & -x_{10} & -c_0 x_{40} & -c_1 x_{50} \\ x_{30} & 0 & -x_{01} & -x_{11} & -c_0 x_{41} & -c_1 x_{51} \\ x_{40} & x_{41} & x_{42} & x_{43} & 0 & x_{45} \\ x_{50} & x_{51} & x_{52} & x_{53} & -\frac{c_0 x_{45}}{c_1} & 0 \end{bmatrix}$$

1.3 Torus

The torus has rank 2. A generic element of the torus is

$$\begin{bmatrix} t_0 & 0 & 0 & 0 & 0 & 0 \\ 0 & t_1 & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{t_0} & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{t_1} & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

The rows of the matrix below are trivial characters:

$$\begin{bmatrix} 1 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 1 & 1 & 1 & 1 & 1 & 1 \end{bmatrix}$$

2 Root system

Dynkin type	B_2
Irreducible	True
Reduced	True
Simply laced	False
Number of roots	8

2.1 Dynkin diagram

The root system is irreducible and has Dynkin diagram:



2.2 Cartan matrix

The root system is irreducible and has Cartan matrix:

$$\begin{bmatrix} 2 & -1 \\ -2 & 2 \end{bmatrix}$$

2.3 Roots

Definition 2.1. A root $\alpha \in \Phi$ is **multipliable** if $2\alpha \in \Phi$.

Root	Squared norm	Simple	Sign	Multipliable
(0, 1, 0, 0, 0, 0)	1	Yes	+	No
(1, 0, 0, 0, 0, 0)	1	No	+	No
(-1, 0, 0, 0, 0, 0)	1	No	-	No
(0, -1, 0, 0, 0, 0)	1	No	-	No
(1, -1, 0, 0, 0, 0)	2	Yes	+	No
(1, 1, 0, 0, 0, 0)	2	No	+	No
(-1, -1, 0, 0, 0, 0)	2	No	-	No
(-1, 1, 0, 0, 0, 0)	2	No	-	No

2.4 Coroots

Root	Coroot
(-1, -1, 0, 0, 0, 0)	(-1, -1, 1, 1, 0, 0)
(-1, 0, 0, 0, 0, 0)	(-2, 0, 2, 0, 0, 0)
(-1, 1, 0, 0, 0, 0)	(-1, 1, 1, -1, 0, 0)
(0, -1, 0, 0, 0, 0)	(0, -2, 0, 2, 0, 0)
(0, 1, 0, 0, 0, 0)	(0, 2, 0, -2, 0, 0)
(1, -1, 0, 0, 0, 0)	(1, -1, -1, 1, 0, 0)
(1, 0, 0, 0, 0, 0)	(2, 0, -2, 0, 0, 0)
(1, 1, 0, 0, 0, 0)	(1, 1, -1, -1, 0, 0)

2.5 Linear combinations of roots

Definition 2.2. For two roots α, β in a root system Φ , the set of positive integral linear combinations of α, β that are also roots is denoted $(\alpha, \beta) = \{i\alpha + j\beta \in \Phi : i, j \in \mathbb{Z}_{\geq 1}\}$.

α	β	Pairs (i, j)
(-1, 0, 0, 0, 0, 0)	(0, -1, 0, 0, 0, 0)	(1, 1)
(-1, 0, 0, 0, 0, 0)	(0, 1, 0, 0, 0, 0)	(1, 1)

α	β	Pairs (i, j)
$(0, -1, 0, 0, 0, 0)$	$(1, 0, 0, 0, 0, 0)$	$(1, 1)$
$(0, 1, 0, 0, 0, 0)$	$(1, 0, 0, 0, 0, 0)$	$(1, 1)$
$(-1, -1, 0, 0, 0, 0)$	$(0, 1, 0, 0, 0, 0)$	$(1, 1), (1, 2)$
$(-1, -1, 0, 0, 0, 0)$	$(1, 0, 0, 0, 0, 0)$	$(1, 1), (1, 2)$
$(-1, 0, 0, 0, 0, 0)$	$(1, -1, 0, 0, 0, 0)$	$(1, 1), (2, 1)$
$(-1, 0, 0, 0, 0, 0)$	$(1, 1, 0, 0, 0, 0)$	$(1, 1), (2, 1)$
$(-1, 1, 0, 0, 0, 0)$	$(0, -1, 0, 0, 0, 0)$	$(1, 1), (1, 2)$
$(-1, 1, 0, 0, 0, 0)$	$(1, 0, 0, 0, 0, 0)$	$(1, 1), (1, 2)$
$(0, -1, 0, 0, 0, 0)$	$(1, 1, 0, 0, 0, 0)$	$(1, 1), (2, 1)$
$(0, 1, 0, 0, 0, 0)$	$(1, -1, 0, 0, 0, 0)$	$(1, 1), (2, 1)$

3 Root spaces

3.1 Dimensions

The table below lists each root along with the dimension of the associated root space (and root subgroup).

Root (α)	Dimension of root space (d_α)
$(-1, -1, 0, 0, 0, 0)$	1
$(-1, 1, 0, 0, 0, 0)$	1
$(1, -1, 0, 0, 0, 0)$	1
$(1, 1, 0, 0, 0, 0)$	1
$(-1, 0, 0, 0, 0, 0)$	2
$(0, -1, 0, 0, 0, 0)$	2
$(0, 1, 0, 0, 0, 0)$	2
$(1, 0, 0, 0, 0, 0)$	2

3.2 Root space table

The table below lists each root along with a generic element of the associated root space (inside the Lie algebra).
Include a definition here? INCOMPLETE

Root	Generic element of root space
$(-1, -1, 0, 0, 0, 0)$	$\begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -u_0 & 0 & 0 & 0 & 0 \\ u_0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$
$(-1, 0, 0, 0, 0, 0)$	$\begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -c_0 u_1 & -c_1 u_0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ u_1 & 0 & 0 & 0 & 0 & 0 \\ u_0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$
$(-1, 1, 0, 0, 0, 0)$	$\begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ u_0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -u_0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$
$(0, -1, 0, 0, 0, 0)$	$\begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -c_0 u_0 & -c_1 u_1 \\ 0 & u_0 & 0 & 0 & 0 & 0 \\ 0 & u_1 & 0 & 0 & 0 & 0 \end{bmatrix}$
$(0, 1, 0, 0, 0, 0)$	$\begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -c_0 u_1 & -c_1 u_0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & u_1 & 0 & 0 \\ 0 & 0 & 0 & u_0 & 0 & 0 \end{bmatrix}$

Root	Generic element of root space
$(1, -1, 0, 0, 0, 0)$	$\begin{bmatrix} 0 & u_0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -u_0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$
$(1, 0, 0, 0, 0, 0)$	$\begin{bmatrix} 0 & 0 & 0 & 0 & -c_0 u_0 & -c_1 u_1 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & u_0 & 0 & 0 & 0 \\ 0 & 0 & u_1 & 0 & 0 & 0 \end{bmatrix}$
$(1, 1, 0, 0, 0, 0)$	$\begin{bmatrix} 0 & 0 & 0 & -u_0 & 0 & 0 \\ 0 & 0 & u_0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$

4 Root subgroups

4.1 Root subgroup table

The table below lists each root along with a generic element of the associated root subgroup.

Root	Generic element of root subgroup
$(-1, -1, 0, 0, 0, 0)$	$\begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & -u_0 & 1 & 0 & 0 & 0 \\ u_0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$
$(-1, 0, 0, 0, 0, 0)$	$\begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ -\frac{c_0 u_1^2}{2} - \frac{c_1 u_0^2}{2} & 0 & 1 & 0 & -c_0 u_1 & -c_1 u_0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ u_1 & 0 & 0 & 0 & 1 & 0 \\ u_0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$
$(-1, 1, 0, 0, 0, 0)$	$\begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ u_0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & -u_0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$
$(0, -1, 0, 0, 0, 0)$	$\begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & -\frac{c_0 u_0^2}{2} - \frac{c_1 u_1^2}{2} & 0 & 1 & -c_0 u_0 & -c_1 u_1 \\ 0 & u_0 & 0 & 0 & 1 & 0 \\ 0 & u_1 & 0 & 0 & 0 & 1 \end{bmatrix}$
$(0, 1, 0, 0, 0, 0)$	$\begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & -\frac{c_0 u_1^2}{2} - \frac{c_1 u_0^2}{2} & -c_0 u_1 & -c_1 u_0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & u_1 & 1 & 0 \\ 0 & 0 & 0 & u_0 & 0 & 1 \end{bmatrix}$
$(1, -1, 0, 0, 0, 0)$	$\begin{bmatrix} 1 & u_0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & -u_0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$
$(1, 0, 0, 0, 0, 0)$	$\begin{bmatrix} 1 & 0 & -\frac{c_0 u_0^2}{2} - \frac{c_1 u_1^2}{2} & 0 & -c_0 u_0 & -c_1 u_1 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & u_0 & 0 & 1 & 0 \\ 0 & 0 & u_1 & 0 & 0 & 1 \end{bmatrix}$

Root	Generic element of root subgroup
(1, 1, 0, 0, 0, 0)	$\begin{bmatrix} 1 & 0 & 0 & -u_0 & 0 & 0 \\ 0 & 1 & u_0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$

4.2 Homomorphism property

For each root $\alpha \in \Phi$, the associated root subgroup map $X_\alpha : V_\alpha \rightarrow G$ is a pseudo-homomorphism with respect to the additive structure on V_α and the group operation (i.e. matrix multiplication) in G . That is, for any $u, v \in V_\alpha$,

$$X_\alpha(u) \cdot X_\alpha(v) = X_\alpha(u + v) \cdot \prod_{\substack{i > 1 \\ i\alpha \in \Phi}} X_{i\alpha}(q_\alpha^i(u, v)) \quad (1)$$

for some homogeneous polynomial maps $q_\alpha^i : V_\alpha \times V_\alpha \rightarrow V_{i\alpha}$. Thus X_α is a group homomorphism if and only the product on the right hand side reduces to 1. The quantities $q_\alpha^i(u, v)$ are called **homomorphism defect coefficients**. For any root α of a root system Φ of any group considered in the Pinning-Sympy package, the set

$$\{i\alpha \in \Phi : i \in \mathbb{Z}_{\geq 2}\}$$

is either empty or a singleton set consisting of $\{2\alpha\}$. The non-empty case only arises when Φ is a non-reduced root system (of Dynkin type BC), and when α is a multipliable root.

Since the root system for this group is reduced, there are no multipliable roots and the product is always empty, hence each X_α is a group homomorphism there are no homomorphism defect coefficients.

5 Commutators

The Steinberg/Chevalley commutator formula says that for two roots $\alpha, \beta \in \Phi$, the commutator of two root group elements $X_\alpha(u)$ and $X_\beta(v)$ is controlled by the structure of Φ , in particular by which positive integer linear combinations $i\alpha + j\beta$ are elements of Φ . Recall that

$$(\alpha, \beta) = \{i\alpha + j\beta \in \Phi : i, j \in \mathbb{Z}_{\geq 1}\}$$

Specifically, we have the commutator formula

$$[X_\alpha(u), X_\beta(v)] = \prod_{(\alpha, \beta)} X_{i\alpha + j\beta} \left(N_{ij}^{\alpha\beta}(u, v) \right)$$

for some homogeneous polynomial functions $N_{ij}^{\alpha\beta} : V_\alpha \times V_\beta \rightarrow V_{i\alpha + j\beta}$. We repeat here the table of linear combinations of roots. The quantity $N_{ij}^{\alpha\beta}(u, v)$ is called a **commutator coefficient**. The table below gives these coefficients.

α	β	i	j	$i\alpha + j\beta$	$N_{ij}^{\alpha\beta}(u, v)$
(-1, 0, 0, 0, 0, 0)	(0, -1, 0, 0, 0, 0)	1	1	(-1, -1, 0, 0, 0, 0)	$c_0 u_0 v_1 + c_1 u_1 v_0$
(-1, 0, 0, 0, 0, 0)	(0, 1, 0, 0, 0, 0)	1	1	(-1, 1, 0, 0, 0, 0)	$c_0 u_0 v_0 + c_1 u_1 v_1$
(0, -1, 0, 0, 0, 0)	(-1, 0, 0, 0, 0, 0)	1	1	(-1, -1, 0, 0, 0, 0)	$-c_0 u_1 v_0 - c_1 u_0 v_1$
(0, -1, 0, 0, 0, 0)	(1, 0, 0, 0, 0, 0)	1	1	(1, -1, 0, 0, 0, 0)	$c_0 u_1 v_0 + c_1 u_0 v_1$
(0, 1, 0, 0, 0, 0)	(-1, 0, 0, 0, 0, 0)	1	1	(-1, 1, 0, 0, 0, 0)	$-c_0 u_0 v_0 - c_1 u_1 v_1$
(0, 1, 0, 0, 0, 0)	(1, 0, 0, 0, 0, 0)	1	1	(1, 1, 0, 0, 0, 0)	$-c_0 u_0 v_0 - c_1 u_1 v_1$
(1, 0, 0, 0, 0, 0)	(0, -1, 0, 0, 0, 0)	1	1	(1, -1, 0, 0, 0, 0)	$-c_0 u_0 v_1 - c_1 u_1 v_0$
(1, 0, 0, 0, 0, 0)	(0, 1, 0, 0, 0, 0)	1	1	(1, 1, 0, 0, 0, 0)	$c_0 u_0 v_0 + c_1 u_1 v_1$
(-1, -1, 0, 0, 0, 0)	(0, 1, 0, 0, 0, 0)	1	1	(-1, 0, 0, 0, 0, 0)	$[-u_0 v_0 \quad -u_0 v_1]$
(-1, -1, 0, 0, 0, 0)	(0, 1, 0, 0, 0, 0)	1	2	(-1, 1, 0, 0, 0, 0)	$\frac{u_0(c_0 v_0^2 + c_1 v_1^2)}{2}$
(-1, -1, 0, 0, 0, 0)	(1, 0, 0, 0, 0, 0)	1	1	(0, -1, 0, 0, 0, 0)	$[u_0 v_1 \quad u_0 v_0]$
(-1, -1, 0, 0, 0, 0)	(1, 0, 0, 0, 0, 0)	1	2	(1, -1, 0, 0, 0, 0)	$\frac{u_0(-c_0 v_0^2 - c_1 v_1^2)}{2}$
(-1, 0, 0, 0, 0, 0)	(1, -1, 0, 0, 0, 0)	1	1	(0, -1, 0, 0, 0, 0)	$[u_1 v_0 \quad u_0 v_0]$
(-1, 0, 0, 0, 0, 0)	(1, -1, 0, 0, 0, 0)	2	1	(-1, -1, 0, 0, 0, 0)	$\frac{v_0(c_0 u_0^2 + c_1 u_1^2)}{2}$
(-1, 0, 0, 0, 0, 0)	(1, 1, 0, 0, 0, 0)	1	1	(0, 1, 0, 0, 0, 0)	$[-u_0 v_0 \quad -u_1 v_0]$
(-1, 0, 0, 0, 0, 0)	(1, 1, 0, 0, 0, 0)	2	1	(-1, 1, 0, 0, 0, 0)	$\frac{v_0(-c_0 u_0^2 - c_1 u_1^2)}{2}$
(-1, 1, 0, 0, 0, 0)	(0, -1, 0, 0, 0, 0)	1	1	(-1, 0, 0, 0, 0, 0)	$[-u_0 v_1 \quad -u_0 v_0]$
(-1, 1, 0, 0, 0, 0)	(0, -1, 0, 0, 0, 0)	1	2	(-1, -1, 0, 0, 0, 0)	$\frac{u_0(c_0 v_1^2 + c_1 v_0^2)}{2}$
(-1, 1, 0, 0, 0, 0)	(1, 0, 0, 0, 0, 0)	1	1	(0, 1, 0, 0, 0, 0)	$[u_0 v_0 \quad u_0 v_1]$
(-1, 1, 0, 0, 0, 0)	(1, 0, 0, 0, 0, 0)	1	2	(1, 1, 0, 0, 0, 0)	$\frac{u_0(c_0 v_0^2 + c_1 v_1^2)}{2}$
(0, -1, 0, 0, 0, 0)	(-1, 1, 0, 0, 0, 0)	1	1	(-1, 0, 0, 0, 0, 0)	$[u_1 v_0 \quad u_0 v_0]$
(0, -1, 0, 0, 0, 0)	(-1, 1, 0, 0, 0, 0)	2	1	(-1, -1, 0, 0, 0, 0)	$\frac{v_0(-c_0 u_1^2 - c_1 u_0^2)}{2}$
(0, -1, 0, 0, 0, 0)	(1, 1, 0, 0, 0, 0)	1	1	(1, 0, 0, 0, 0, 0)	$[u_1 v_0 \quad u_0 v_0]$
(0, -1, 0, 0, 0, 0)	(1, 1, 0, 0, 0, 0)	2	1	(1, -1, 0, 0, 0, 0)	$\frac{v_0(c_0 u_1^2 + c_1 u_0^2)}{2}$
(0, 1, 0, 0, 0, 0)	(-1, -1, 0, 0, 0, 0)	1	1	(-1, 0, 0, 0, 0, 0)	$[u_0 v_0 \quad u_1 v_0]$
(0, 1, 0, 0, 0, 0)	(-1, -1, 0, 0, 0, 0)	2	1	(-1, 1, 0, 0, 0, 0)	$\frac{v_0(-c_0 u_0^2 - c_1 u_1^2)}{2}$
(0, 1, 0, 0, 0, 0)	(1, -1, 0, 0, 0, 0)	1	1	(1, 0, 0, 0, 0, 0)	$[-u_0 v_0 \quad -u_1 v_0]$
(0, 1, 0, 0, 0, 0)	(1, -1, 0, 0, 0, 0)	2	1	(1, 1, 0, 0, 0, 0)	$\frac{v_0(c_0 u_0^2 + c_1 u_1^2)}{2}$
(1, -1, 0, 0, 0, 0)	(-1, 0, 0, 0, 0, 0)	1	1	(0, -1, 0, 0, 0, 0)	$[-u_0 v_1 \quad -u_0 v_0]$
(1, -1, 0, 0, 0, 0)	(-1, 0, 0, 0, 0, 0)	1	2	(-1, -1, 0, 0, 0, 0)	$\frac{u_0(-c_0 v_0^2 - c_1 v_1^2)}{2}$

α	β	i	j	$i\alpha + j\beta$	$N_{ij}^{\alpha\beta}(u, v)$
(1, -1, 0, 0, 0, 0)	(0, 1, 0, 0, 0, 0)	1	1	(1, 0, 0, 0, 0, 0)	$\begin{bmatrix} u_0 v_0 & u_0 v_1 \end{bmatrix}$
(1, -1, 0, 0, 0, 0)	(0, 1, 0, 0, 0, 0)	1	2	(1, 1, 0, 0, 0, 0)	$\frac{u_0(-c_0 v_0^2 - c_1 v_1^2)}{2}$
(1, 0, 0, 0, 0, 0)	(-1, -1, 0, 0, 0, 0)	1	1	(0, -1, 0, 0, 0, 0)	$\begin{bmatrix} -u_1 v_0 & -u_0 v_0 \end{bmatrix}$
(1, 0, 0, 0, 0, 0)	(-1, -1, 0, 0, 0, 0)	2	1	(1, -1, 0, 0, 0, 0)	$\frac{v_0(c_0 u_0^2 + c_1 u_1^2)}{2}$
(1, 0, 0, 0, 0, 0)	(-1, 1, 0, 0, 0, 0)	1	1	(0, 1, 0, 0, 0, 0)	$\begin{bmatrix} -u_0 v_0 & -u_1 v_0 \end{bmatrix}$
(1, 0, 0, 0, 0, 0)	(-1, 1, 0, 0, 0, 0)	2	1	(1, 1, 0, 0, 0, 0)	$\frac{v_0(-c_0 u_0^2 - c_1 u_1^2)}{2}$
(1, 1, 0, 0, 0, 0)	(-1, 0, 0, 0, 0, 0)	1	1	(0, 1, 0, 0, 0, 0)	$\begin{bmatrix} u_0 v_0 & u_0 v_1 \end{bmatrix}$
(1, 1, 0, 0, 0, 0)	(-1, 0, 0, 0, 0, 0)	1	2	(-1, 1, 0, 0, 0, 0)	$\frac{u_0(c_0 v_0^2 + c_1 v_1^2)}{2}$
(1, 1, 0, 0, 0, 0)	(0, -1, 0, 0, 0, 0)	1	1	(1, 0, 0, 0, 0, 0)	$\begin{bmatrix} -u_0 v_1 & -u_0 v_0 \end{bmatrix}$
(1, 1, 0, 0, 0, 0)	(0, -1, 0, 0, 0, 0)	1	2	(1, -1, 0, 0, 0, 0)	$\frac{u_0(-c_0 v_1^2 - c_1 v_0^2)}{2}$

5.1 Commutator identities

Below is an explicit formulation of the commutator formula identities.

CommutatorIdentitiesPlaceholder

6 Weyl group

Background theoretical info on Weyl group, especially as realized inside the group, including info on conjugation between root subgroups INCOMPLETE

6.1 Weyl group elements

The table below lists roots along with an associated Weyl group element.

Root	Weyl group element
$(-1, -1, 0, 0, 0, 0)$	$\begin{bmatrix} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$
$(-1, 0, 0, 0, 0, 0)$	$\begin{bmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 \end{bmatrix}$
$(-1, 1, 0, 0, 0, 0)$	$\begin{bmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$
$(0, -1, 0, 0, 0, 0)$	$\begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 \end{bmatrix}$
$(0, 1, 0, 0, 0, 0)$	$\begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 \end{bmatrix}$
$(1, -1, 0, 0, 0, 0)$	$\begin{bmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$
$(1, 0, 0, 0, 0, 0)$	$\begin{bmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 \end{bmatrix}$

Root	Weyl group element
(1, 1, 0, 0, 0, 0)	$\begin{bmatrix} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$

6.2 Weyl group conjugation coefficients

The table below lists roots along with the coefficients obtain in performing conjugation by an associated Weyl element.

Root (α)	Root (β)	$\gamma = \sigma_\alpha(\beta)$	Weyl conjugation coefficient
(-1, -1, 0, 0, 0, 0)	(-1, -1, 0, 0, 0, 0)	(1, 1, 0, 0, 0, 0)	$-u_0$
(-1, -1, 0, 0, 0, 0)	(-1, 0, 0, 0, 0, 0)	(0, 1, 0, 0, 0, 0)	$\begin{bmatrix} u_0 & u_1 \end{bmatrix}$
(-1, -1, 0, 0, 0, 0)	(-1, 1, 0, 0, 0, 0)	(-1, 1, 0, 0, 0, 0)	$-u_0$
(-1, -1, 0, 0, 0, 0)	(0, -1, 0, 0, 0, 0)	(1, 0, 0, 0, 0, 0)	$\begin{bmatrix} u_1 & u_0 \end{bmatrix}$
(-1, -1, 0, 0, 0, 0)	(0, 1, 0, 0, 0, 0)	(-1, 0, 0, 0, 0, 0)	$\begin{bmatrix} u_0 & u_1 \end{bmatrix}$
(-1, -1, 0, 0, 0, 0)	(1, -1, 0, 0, 0, 0)	(1, -1, 0, 0, 0, 0)	$-u_0$
(-1, -1, 0, 0, 0, 0)	(1, 0, 0, 0, 0, 0)	(0, -1, 0, 0, 0, 0)	$\begin{bmatrix} u_1 & u_0 \end{bmatrix}$
(-1, -1, 0, 0, 0, 0)	(1, 1, 0, 0, 0, 0)	(-1, -1, 0, 0, 0, 0)	$-u_0$
(-1, 0, 0, 0, 0, 0)	(-1, -1, 0, 0, 0, 0)	(1, -1, 0, 0, 0, 0)	$-u_0$
(-1, 0, 0, 0, 0, 0)	(-1, 0, 0, 0, 0, 0)	(1, 0, 0, 0, 0, 0)	$\begin{bmatrix} u_0 & -u_1 \end{bmatrix}$
(-1, 0, 0, 0, 0, 0)	(-1, 1, 0, 0, 0, 0)	(1, 1, 0, 0, 0, 0)	u_0
(-1, 0, 0, 0, 0, 0)	(0, -1, 0, 0, 0, 0)	(0, -1, 0, 0, 0, 0)	$\begin{bmatrix} -u_0 & u_1 \end{bmatrix}$
(-1, 0, 0, 0, 0, 0)	(0, 1, 0, 0, 0, 0)	(0, 1, 0, 0, 0, 0)	$\begin{bmatrix} u_0 & -u_1 \end{bmatrix}$
(-1, 0, 0, 0, 0, 0)	(1, -1, 0, 0, 0, 0)	(-1, -1, 0, 0, 0, 0)	$-u_0$
(-1, 0, 0, 0, 0, 0)	(1, 0, 0, 0, 0, 0)	(-1, 0, 0, 0, 0, 0)	$\begin{bmatrix} u_0 & -u_1 \end{bmatrix}$
(-1, 0, 0, 0, 0, 0)	(1, 1, 0, 0, 0, 0)	(-1, 1, 0, 0, 0, 0)	u_0
(-1, 1, 0, 0, 0, 0)	(-1, -1, 0, 0, 0, 0)	(-1, -1, 0, 0, 0, 0)	$-u_0$
(-1, 1, 0, 0, 0, 0)	(-1, 0, 0, 0, 0, 0)	(0, -1, 0, 0, 0, 0)	$\begin{bmatrix} u_1 & u_0 \end{bmatrix}$
(-1, 1, 0, 0, 0, 0)	(-1, 1, 0, 0, 0, 0)	(1, -1, 0, 0, 0, 0)	u_0
(-1, 1, 0, 0, 0, 0)	(0, -1, 0, 0, 0, 0)	(-1, 0, 0, 0, 0, 0)	$\begin{bmatrix} u_1 & u_0 \end{bmatrix}$
(-1, 1, 0, 0, 0, 0)	(0, 1, 0, 0, 0, 0)	(1, 0, 0, 0, 0, 0)	$\begin{bmatrix} u_0 & u_1 \end{bmatrix}$
(-1, 1, 0, 0, 0, 0)	(1, -1, 0, 0, 0, 0)	(-1, 1, 0, 0, 0, 0)	u_0
(-1, 1, 0, 0, 0, 0)	(1, 0, 0, 0, 0, 0)	(0, 1, 0, 0, 0, 0)	$\begin{bmatrix} u_0 & u_1 \end{bmatrix}$
(-1, 1, 0, 0, 0, 0)	(1, 1, 0, 0, 0, 0)	(1, 1, 0, 0, 0, 0)	$-u_0$
(0, -1, 0, 0, 0, 0)	(-1, -1, 0, 0, 0, 0)	(-1, 1, 0, 0, 0, 0)	u_0
(0, -1, 0, 0, 0, 0)	(-1, 0, 0, 0, 0, 0)	(-1, 0, 0, 0, 0, 0)	$\begin{bmatrix} u_0 & -u_1 \end{bmatrix}$
(0, -1, 0, 0, 0, 0)	(-1, 1, 0, 0, 0, 0)	(-1, -1, 0, 0, 0, 0)	u_0
(0, -1, 0, 0, 0, 0)	(0, -1, 0, 0, 0, 0)	(0, 1, 0, 0, 0, 0)	$\begin{bmatrix} u_1 & -u_0 \end{bmatrix}$
(0, -1, 0, 0, 0, 0)	(0, 1, 0, 0, 0, 0)	(0, -1, 0, 0, 0, 0)	$\begin{bmatrix} -u_1 & u_0 \end{bmatrix}$
(0, -1, 0, 0, 0, 0)	(1, -1, 0, 0, 0, 0)	(1, 1, 0, 0, 0, 0)	$-u_0$
(0, -1, 0, 0, 0, 0)	(1, 0, 0, 0, 0, 0)	(1, 0, 0, 0, 0, 0)	$\begin{bmatrix} u_0 & -u_1 \end{bmatrix}$
(0, -1, 0, 0, 0, 0)	(1, 1, 0, 0, 0, 0)	(1, -1, 0, 0, 0, 0)	$-u_0$
(0, 1, 0, 0, 0, 0)	(-1, -1, 0, 0, 0, 0)	(-1, 1, 0, 0, 0, 0)	u_0
(0, 1, 0, 0, 0, 0)	(-1, 0, 0, 0, 0, 0)	(-1, 0, 0, 0, 0, 0)	$\begin{bmatrix} u_0 & -u_1 \end{bmatrix}$
(0, 1, 0, 0, 0, 0)	(-1, 1, 0, 0, 0, 0)	(-1, -1, 0, 0, 0, 0)	u_0
(0, 1, 0, 0, 0, 0)	(0, -1, 0, 0, 0, 0)	(0, 1, 0, 0, 0, 0)	$\begin{bmatrix} u_1 & -u_0 \end{bmatrix}$
(0, 1, 0, 0, 0, 0)	(0, 1, 0, 0, 0, 0)	(0, -1, 0, 0, 0, 0)	$\begin{bmatrix} -u_1 & u_0 \end{bmatrix}$

Root (α)	Root (β)	$\gamma = \sigma_\alpha(\beta)$	Weyl conjugation coefficient
(0, 1, 0, 0, 0, 0)	(1, -1, 0, 0, 0, 0)	(1, 1, 0, 0, 0, 0)	$-u_0$
(0, 1, 0, 0, 0, 0)	(1, 0, 0, 0, 0, 0)	(1, 0, 0, 0, 0, 0)	$\begin{bmatrix} u_0 & -u_1 \end{bmatrix}$
(0, 1, 0, 0, 0, 0)	(1, 1, 0, 0, 0, 0)	(1, -1, 0, 0, 0, 0)	$-u_0$
(1, -1, 0, 0, 0, 0)	(-1, -1, 0, 0, 0, 0)	(-1, -1, 0, 0, 0, 0)	$-u_0$
(1, -1, 0, 0, 0, 0)	(-1, 0, 0, 0, 0, 0)	(0, -1, 0, 0, 0, 0)	$\begin{bmatrix} u_1 & u_0 \end{bmatrix}$
(1, -1, 0, 0, 0, 0)	(-1, 1, 0, 0, 0, 0)	(1, -1, 0, 0, 0, 0)	u_0
(1, -1, 0, 0, 0, 0)	(0, -1, 0, 0, 0, 0)	(-1, 0, 0, 0, 0, 0)	$\begin{bmatrix} u_1 & u_0 \end{bmatrix}$
(1, -1, 0, 0, 0, 0)	(0, 1, 0, 0, 0, 0)	(1, 0, 0, 0, 0, 0)	$\begin{bmatrix} u_0 & u_1 \end{bmatrix}$
(1, -1, 0, 0, 0, 0)	(1, -1, 0, 0, 0, 0)	(-1, 1, 0, 0, 0, 0)	u_0
(1, -1, 0, 0, 0, 0)	(1, 0, 0, 0, 0, 0)	(0, 1, 0, 0, 0, 0)	$\begin{bmatrix} u_0 & u_1 \end{bmatrix}$
(1, -1, 0, 0, 0, 0)	(1, 1, 0, 0, 0, 0)	(1, 1, 0, 0, 0, 0)	$-u_0$
(1, 0, 0, 0, 0, 0)	(-1, -1, 0, 0, 0, 0)	(1, -1, 0, 0, 0, 0)	$-u_0$
(1, 0, 0, 0, 0, 0)	(-1, 0, 0, 0, 0, 0)	(1, 0, 0, 0, 0, 0)	$\begin{bmatrix} u_0 & -u_1 \end{bmatrix}$
(1, 0, 0, 0, 0, 0)	(-1, 1, 0, 0, 0, 0)	(1, 1, 0, 0, 0, 0)	u_0
(1, 0, 0, 0, 0, 0)	(0, -1, 0, 0, 0, 0)	(0, -1, 0, 0, 0, 0)	$\begin{bmatrix} -u_0 & u_1 \end{bmatrix}$
(1, 0, 0, 0, 0, 0)	(0, 1, 0, 0, 0, 0)	(0, 1, 0, 0, 0, 0)	$\begin{bmatrix} u_0 & -u_1 \end{bmatrix}$
(1, 0, 0, 0, 0, 0)	(1, -1, 0, 0, 0, 0)	(-1, -1, 0, 0, 0, 0)	$-u_0$
(1, 0, 0, 0, 0, 0)	(1, 0, 0, 0, 0, 0)	(-1, 0, 0, 0, 0, 0)	$\begin{bmatrix} u_0 & -u_1 \end{bmatrix}$
(1, 0, 0, 0, 0, 0)	(1, 1, 0, 0, 0, 0)	(-1, 1, 0, 0, 0, 0)	u_0
(1, 1, 0, 0, 0, 0)	(-1, -1, 0, 0, 0, 0)	(1, 1, 0, 0, 0, 0)	$-u_0$
(1, 1, 0, 0, 0, 0)	(-1, 0, 0, 0, 0, 0)	(0, 1, 0, 0, 0, 0)	$\begin{bmatrix} u_0 & u_1 \end{bmatrix}$
(1, 1, 0, 0, 0, 0)	(-1, 1, 0, 0, 0, 0)	(-1, 1, 0, 0, 0, 0)	$-u_0$
(1, 1, 0, 0, 0, 0)	(0, -1, 0, 0, 0, 0)	(1, 0, 0, 0, 0, 0)	$\begin{bmatrix} u_1 & u_0 \end{bmatrix}$
(1, 1, 0, 0, 0, 0)	(0, 1, 0, 0, 0, 0)	(-1, 0, 0, 0, 0, 0)	$\begin{bmatrix} u_0 & u_1 \end{bmatrix}$
(1, 1, 0, 0, 0, 0)	(1, -1, 0, 0, 0, 0)	(1, -1, 0, 0, 0, 0)	$-u_0$
(1, 1, 0, 0, 0, 0)	(1, 0, 0, 0, 0, 0)	(0, -1, 0, 0, 0, 0)	$\begin{bmatrix} u_1 & u_0 \end{bmatrix}$
(1, 1, 0, 0, 0, 0)	(1, 1, 0, 0, 0, 0)	(-1, -1, 0, 0, 0, 0)	$-u_0$

6.3 Weyl group conjugation identities

Below is an explicit formulation of the Weyl group conjugation identities.

WeylConjugationIdentitiesPlaceholder