

DATA 605 - Assignment 2

Joshua Sturm

02/05/2018

Problem Set 1

1. Show that $A^T A \neq A A^T$ in general. (Proof and demonstration.)

Solution

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then $A^T = \begin{bmatrix} a & c \\ b & d \end{bmatrix}$.

The product of $A A^T$ will be: $\begin{bmatrix} a \times a + b \times b & a \times c + b \times d \\ c \times a + d \times b & c \times c + d \times d \end{bmatrix}$.

The product of $A^T A$ will be: $\begin{bmatrix} a \times a + c \times c & a \times b + c \times d \\ a \times b + c \times d & b \times b + d \times d \end{bmatrix}$.

Thus, we can see that matrix multiplication does not have the commutative property.

In the event that A is not a square matrix, then the proof is trivial. Say B is an $m \times n$ matrix, then B^T will be an $n \times m$ matrix. $B B^T$ will thus be an $m \times m$ matrix, which is a square matrix.

2. For a special type of square matrix A , we get $A^T A = A A^T$. Under what conditions could this be true? (Hint: The identity matrix I is an example of such a matrix.)

Solution

Theorem: For any matrix A , $A A^T$ and $A^T A$ are symmetric matrices.

The product of two symmetric matrices is symmetric only if both matrices commute, i.e. $A A^T = A^T A$.

(https://en.wikipedia.org/wiki/Symmetric_matrix#Properties) / This is only possible when the matrix is invertible.

Problem Set 3

Write an R function to factorize a square matrix A into LU or LDU, whichever you prefer. You don't have to worry about permuting rows of A and you can assume that A is less than 5x5, if you need to hard-code any variables in your code

```
# Assume the user inputs a square matrix.
```

```
factorize <- function(mat){  
  L <- diag(nrow(mat)) # Create lower diagonal matrix of the same size  
  U <- mat  
  
  # If the leading entry [1,1] is not > 1, swap it with a row that has a pivot  
  if (U[1,1] == 0){  
    for (i in U[2,1]:U[nrow,1]){
```

```

    if (U[i,1] != 0){
      U[1,] <- U[i,]
    }
  }
}
# Loop through n-1 columns, n rows, get the multiplier,
# Multiply to get the Upper Triangular matrix, and plug
# the multiplier into the Lower Triangular Matrix
for (k in 1:(nrow(U)-1)){
  for (j in (k+1):(nrow(U))){
    if (U[j,k] != 0){
      mult <- U[j,k] / U[k,k]
      U[j,] <- U[j,] - (mult * U[k,])
      L[j,k] <- mult
    }
  }
}

factored.matrix <- list(L, U)

return(factored.matrix)
}

```