

Learning Lean 4

Joshua W. Kelly

2026-7-17

Abstract

Working notes and exercises from learning the Lean 4 theorem prover, following *An Introduction to Lean 4* by Cosme Llópez and Gong. Covers basic syntax and natural-deduction proofs, paired with a runnable Lean library.

Contents

1	Basic Syntax	1
1.1	Basic syntax	1
1.1.1	Comments	1
1.1.2	Inspecting terms and types with <code>#check</code>	1
1.1.3	Printing definitions with <code>#print</code>	1
1.1.4	Definitions and functions	1
2	Natural Deduction	2
2.1	Natural deduction	2
2.1.1	5.5 Reduction to the absurd. $P \Rightarrow Q, \neg Q \vdash \neg P$	2
2.1.2	Conjunction introduction. $P, P \Rightarrow Q \vdash P \wedge Q$	2
3	Linear Algebra	3
3.1	Linear algebra	3
3.1.1	Vectors	3
3.1.2	Matrices	3
3.1.3	Linear maps	4
3.1.4	A worked computation	4
3.1.5	Exercises	4

1 Basic Syntax

1.1 Basic syntax

The runnable version of this chapter lives in `LearningLean/BasicSyntax.lean`.

1.1.1 Comments

```
-- This is a single-line comment.
```

```
/-  
This is a multi-line comment.  
Here is another line.  
-/
```

1.1.2 Inspecting terms and types with `#check`

`#check` asks Lean for the type of an expression without evaluating it.

```
#check true           -- Bool  
#check 42             -- Nat  
#check "Hello, World!" -- String  
#check Type           -- Type 1  
#check Nat            -- Type  
#check ['h', 'e', 'l', 'l', 'o'] -- List Char
```

1.1.3 Printing definitions with `#print`

`#print` shows how something is defined.

```
#print Bool
```

Type itself cannot be printed with `#print`, because it is a type rather than a definition.

1.1.4 Definitions and functions

```
def pi : Float := 3.1415926
```

```
#check pi
```

```
-- An anonymous function (lambda) taking two natural numbers.
```

```
#check λ (a b : Nat) => a + b
```

2 Natural Deduction

2.1 Natural deduction

The Lean proofs for this chapter live in `LearningLean/NaturalDeduction.lean`, inside the `Exercises` namespace.

2.1.1 5.5 Reduction to the absurd. $P \Rightarrow Q, \neg Q \vdash \neg P$

This is a very useful technique. Validity of $P \Rightarrow Q, \neg Q \vdash \neg P$ is proved with:

1	$P \Rightarrow Q$	
2	$\neg Q$	
3	$\begin{array}{ l} P \end{array}$	H
4	$\begin{array}{ l} Q \end{array}$	$E \Rightarrow 1,3$
5	$\begin{array}{ l} \neg Q \end{array}$	IT 2
6	$\neg P$	$I \neg 3,4,5$

The sub-derivation (lines 3–5) discharges the hypothesis P : assuming P , [$\Rightarrow$ -elimination] on lines 1 and 3 gives Q , while line 2 reiterates $\neg Q$. The contradiction lets us introduce $\neg P$ ($\neg$ -introduction over lines 3, 4, 5).

In Lean this is theorem `T55`, where $\neg P$ is definitionally $P \rightarrow \text{False}$, so the whole proof is just function composition:

```
theorem T55 (h1 : P → Q) (h2 : ¬Q) : ¬P :=  
  (h2 ∘ h1)
```

2.1.2 Conjunction introduction. $P, P \Rightarrow Q \vdash P \wedge Q$

Theorem `T51` builds the pair (proof of P , proof of Q) directly, using the anonymous-constructor notation `{_, _}`:

```
theorem T51 (h1 : P) (h2 : P → Q) : P ∧ Q :=  
  {h1, h2 h1}
```

3 Linear Algebra

3.1 Linear algebra

A scratch chapter for linear-algebra notes — enough structure to flesh out later, and enough variety (numbered equations, a table, admonitions, and interleaved Lean) to show the book template in action. The runnable code lives in `LearningLean/LinearAlgebra.lean`.

Note

This chapter is a work in progress. Sections marked **To flesh out** are placeholders.

3.1.1 Vectors

A vector in R^n is an ordered list of n real numbers. We write column vectors as $\vec{u} = (u_1, \dots, u_n)$ with each $u_i \in R$.

The **dot product** of $\vec{u}, \vec{v} \in R^n$ is the scalar

$$\vec{u} \cdot \vec{v} = \sum_{i=1}^n u_i v_i.$$

For a concrete computation, take $\vec{u} = (1, 2, 3)$ and $\vec{v} = (4, 5, 6)$. Then by equation (1),

$$\vec{u} \cdot \vec{v} = 1 \cdot 4 + 2 \cdot 5 + 3 \cdot 6 = 32.$$

In Lean, a vector is a function `Fin n → ℝ`, and `![...]` is notation for one:

```
def u : Fin 3 → ℝ := ![1, 2, 3]
def v : Fin 3 → ℝ := ![4, 5, 6]

#check u [?] v      -- `[?]' is Mathlib's dot product, of type ℝ
```

3.1.2 Matrices

An $m \times n$ matrix $A \in R^{m \times n}$ is a rectangular array of scalars. For example,

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \in R^{2 \times 2}.$$

The **matrix–vector product** $A\vec{x}$ takes a vector $\vec{x} \in R^n$ to a vector in R^m . With $\vec{x} = (5, 6)$:

$$A\vec{x} = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \begin{bmatrix} 5 \\ 6 \end{bmatrix} = \begin{bmatrix} 1 \cdot 5 + 2 \cdot 6 \\ 3 \cdot 5 + 4 \cdot 6 \end{bmatrix} = \begin{bmatrix} 17 \\ 39 \end{bmatrix}.$$

A few operations and their Mathlib spellings:

Operation	Math	Lean / Mathlib
Dot product	$\vec{u} \cdot \vec{v}$	<code>u · v</code>
Matrix–vector product	$A\vec{x}$	<code>A *v x</code>
Matrix product	AB	<code>A * B</code>
Transpose	$A^\top$	<code>Aᵀ</code>

```
def A : Matrix (Fin 2) (Fin 2) ℝ := !![1, 2; 3, 4]

#check A *v ![5, 6] -- matrix-vector product, of type `Fin 2 → ℝ`
```

3.1.3 Linear maps

A **linear map** $T : R^n \rightarrow R^m$ is a function satisfying $T(\alpha\vec{u} + \vec{v}) = \alpha T(\vec{u}) + T(\vec{v})$ for all scalars α and vectors $\vec{u}, \vec{v}$. Every such map is represented by a matrix, and matrix multiplication corresponds to composition of the maps.

Note

To flesh out: state and prove that `Matrix.mulVec A` is a `LinearMap`, and connect it to `Matrix.toLin`.

3.1.4 A worked computation

To flesh out. A step-by-step example — e.g. solving $A\vec{x} = \vec{b}$ by row reduction — narrated alongside the corresponding Lean definitions, so the computation and its formalization sit side by side.

3.1.5 Exercises

Note

To flesh out. Add exercises here, e.g.:

1. Show the dot product is symmetric: $\vec{u} \cdot \vec{v} = \vec{v} \cdot \vec{u}$.
2. Prove $(A^\top)^\top = A$ in Lean using `Matrix.transpose_transpose`.