

OPTIMIZERS OF THREE-POINT ENERGIES AND NEARLY ORTHOGONAL SETS

DMITRIY BILYK, DAMIR FERIZOVIĆ, ALEXEY GLAZYRIN, RYAN W. MATZKE,
 JOSIAH PARK, AND OLEKSANDR VLASIUK

(Communicated by Yuan Xu)

ABSTRACT. This paper is devoted to spherical measures and point configurations optimizing three-point energies. Our main goal is to extend the classic optimization problems based on pairs of distances between points to the context of three-point potentials. In particular, we study three-point analogues of the sphere packing problem and the optimization problem for p -frame energies based on three points. It turns out that both problems are inherently connected to the problem of nearly orthogonal sets by Erdős. As the outcome, we provide a new solution of the Erdős problem from the three-point packing perspective. We also show that the orthogonal basis uniquely minimizes the p -frame three-point energy when $0 < p < 1$ in all dimensions. The arguments make use of multivariate polynomials employed in semidefinite programming and based on the classical Gegenbauer polynomials. For $p = 1$, we completely solve the analogous problem on the circle. As for higher dimensions, we show that the Hausdorff dimension of minimizers is not greater than $d - 2$ for measures on \mathbb{S}^{d-1} .

CONTENTS

1. Introduction	4016
2. Notation and preliminaries	4017
3. Three-point packing problem and nearly orthogonal sets	4019
4. Minimal energy for multivariate p -frame potentials	4022
5. Isotropic measures without obtuse angles	4026
6. Support of a minimizer for 1-frame energy has dimension not greater than $d - 2$	4029
7. Appendix	4030
Acknowledgments	4031
References	4031

Received by the editors March 21, 2023, and, in revised form, January 23, 2024.

2020 Mathematics Subject Classification. Primary 52C17, 90C22; Secondary 52A40, 05D05.

Key words and phrases. Potential energy minimization, optimal measures, positive definite kernels, spherical codes, tight frames, isotropic measures, nearly orthogonal sets.

The first author was supported by the NSF grant DMS-2054606 and Simons Collaboration Grant 712810. The second author was supported by the Methusalem grant METH/21/03—long term structural funding of the Flemish Government. The third author was supported by the NSF grant DMS-2054536. The fourth author was supported by the Doctoral Dissertation Fellowship of the University of Minnesota, the Austrian Science Fund FWF project F5503 part of the Special Research Program (SFB) “Quasi-Monte Carlo Methods: Theory and Applications”, and NSF Postdoctoral Fellowship Grant 2202877. The sixth author was supported by an AMS-Simons Travel Grant.

1. INTRODUCTION

Various interesting point configurations in metric spaces are realized as optimizers of energies defined by two-point potentials. For an N -point configuration ω_N on the unit sphere, \mathbb{S}^{d-1} , the energy is defined as

$$(1.1) \quad E_K(\omega_N) = \frac{1}{N^2} \sum_{x,y \in \omega_N} K(x,y),$$

where K is the function defining a two-point potential. Typically, K depends on the Euclidean distance between points x and y . A very interesting example, closely connected to unit norm tight frames and spherical designs, is the 2-frame potential $K(x,y) = |\langle x,y \rangle|^2$, or simply *frame potential*, introduced by Benedetto and Fickus [BF], and later generalized, by Ehler and Okoudjou [EO], to the p -frame potential $|\langle x,y \rangle|^p$, for $p \in (0,\infty)$, which has been studied further in, e.g., [BE, BCGKO, BGMPV1, BGMPV2, CGGKO, Gl, GP, H1, H2, WO, XX]. See [BHS] for an extensive introduction on energy optimization for two-point potentials.

In this paper we study three-point potentials, i.e. analogous energies depending on interactions of triples of points rather than pairwise interactions. That is, we shall be concerned with the minimization of three-point discrete energies and energy integrals:

$$E_K(\omega_N) = \frac{1}{N^3} \sum_{x,y,z \in \omega_N} K(x,y,z) \quad \text{and} \\ I_K(\mu) = \int_{\mathbb{S}^{d-1}} \int_{\mathbb{S}^{d-1}} \int_{\mathbb{S}^{d-1}} K(x,y,z) d\mu(x) d\mu(y) d\mu(z),$$

see Section 2 for precise definitions. Such energies arise in various applications and have been previously studied in [BFGMPV, BFGMPV1].

The notion of the p -frame potential is naturally extended to the three-point case by defining the three-point p -frame potential as

$$K(x,y,z) = |uvz|^p = |\langle y,z \rangle \langle x,z \rangle \langle x,y \rangle|^p.$$

In this paper, we are mostly interested in the case $p = 1$, as well as $0 < p < 1$. As we shall see, even in this case the problem of describing minimizers appears to be highly non-trivial and inherently connected to the problem of Erdős about nearly orthogonal sets and packing problems which we describe below.

1.1. Nearly orthogonal sets. A (multi-)set of non-zero vectors in \mathbb{R}^d is called *nearly orthogonal* if, for any three distinct vectors from the set, at least two of them are orthogonal. Erdős asked about the maximum size of nearly orthogonal sets. In 1991, Rosenfeld showed that the maximum size of such a set is $2d$ [Ros]. Other proofs of this result are given in [Pud] and [Dea], and we provide an alternative proof in Theorem 3.1. Nearly orthogonal sets of size $2d$ in \mathbb{S}^{d-1} are called *Rosenfeld sets*. One might be tempted to believe that Rosenfeld sets in \mathbb{R}^d are necessarily unions of two orthogonal bases but, in fact, for $d \geq 5$ this is not true (see [Dea] for examples) and the full characterization of Rosenfeld sets is an open problem.

We would also like to remark that Rosenfeld [Ros] used the term “almost orthogonal sets”, but “nearly orthogonal” seems to be more common in this context in later literature, although both are still used (see, e.g., [AlSz, Pol]).

1.2. Packing problems on the sphere. The classic packing problem on the sphere, or the problem of determining an optimal spherical code, consists in finding the maximal number of points in the configuration $\omega_N \in \mathbb{S}^{d-1}$ such that for any distinct points $x, y \in \omega_N$, $\langle x, y \rangle \leq \alpha$. This condition ensures that distinct points of ω_N are separated by a spherical distance of at least $\arccos \alpha$. In the case $\alpha = 1/2$, the problem is known as a *kissing number* problem that, despite its rich history, is solved only in a handful of small dimensions [SvdW, Lev, OS, Mus].

We study the three-point analogue of the packing problem. Particularly, we are interested in finding the maximal number of points in the configuration $\omega_N \in \mathbb{S}^{d-1}$ such that for any distinct points $x, y, z \in \omega_N$, $\langle x, y \rangle \langle x, z \rangle \langle y, z \rangle \leq \alpha$. In Section 3, we will show that the three-point packing problem for $\alpha = 0$ is surprisingly similar to the classic two-point packing problem for $\alpha = 0$ and the optimal configurations are precisely Rosenfeld sets.

1.3. Outline of the paper. In Section 2 we present some relevant background information, in particular, about isotropic measures, frames, frame energy, and semidefinite programming bounds. In Section 3 we explore the connections between nearly orthogonal sets and three-point packing bounds, giving an alternative proof and generalizing Rosenfeld's result (Theorem 3.1). Section 4 is devoted to the three-point p -frame energy: in particular, we show that in all dimensions for $0 < p < 1$ any minimizer of the p -frame energy is a uniform distribution over an orthonormal basis, up to symmetries (Theorem 4.4) and on \mathbb{S}^1 minimizers of the 1-frame energy are convex combinations of uniform distributions over two orthonormal bases (Theorem 4.6). In Section 5 we use a spherical version of Jung's inequality [D] to show that any isotropic measure without obtuse angles in its support is a uniform distribution over an orthonormal basis. Finally, in Section 6 we show that the support of a measure minimizing the three-point 1-frame energy must have codimension at least one (Theorem 6.2).

2. NOTATION AND PRELIMINARIES

The notation in the paper follows [BFGMPV, BFGMPV1]. All potentials are defined for triples of points (x, y, z) on the unit sphere \mathbb{S}^{d-1} . For brevity, throughout the paper we use the notation $u = \langle y, z \rangle$, $v = \langle x, z \rangle$, $t = \langle x, y \rangle$, where $\langle \cdot, \cdot \rangle$ denotes the standard Euclidean inner product. We denote by $\mathcal{P}(\mathbb{S}^{d-1})$ the set of Borel probability measures on \mathbb{S}^{d-1} . The normalized uniform distribution over \mathbb{S}^{d-1} is denoted by σ . A measure μ is called *balanced* if its center of mass is at the origin, that is, $\int_{\mathbb{S}^{d-1}} x d\mu(x) = 0$.

2.1. Three-point energies. Let $\omega_N = \{z_1, z_2, \dots, z_N\}$ be an N -point configuration (multiset) in \mathbb{S}^{d-1} , for $N \geq 3$. Given a continuous three-point kernel $K : (\mathbb{S}^{d-1})^3 \rightarrow \mathbb{R}$, the discrete K -energy of ω_N is defined to be

$$(2.1) \quad E_K(\omega_N) := \frac{1}{N^3} \sum_{i=1}^N \sum_{j=1}^N \sum_{k=1}^N K(z_i, z_j, z_k).$$

Similarly, we define the energy for a measure $\mu \in \mathcal{P}(\mathbb{S}^{d-1})$:

$$(2.2) \quad I_K(\mu) = \int_{\mathbb{S}^{d-1}} \int_{\mathbb{S}^{d-1}} \int_{\mathbb{S}^{d-1}} K(x, y, z) d\mu(x) d\mu(y) d\mu(z).$$

Under the normalization in (2.1), $E_K(\omega_N) = I_K(\mu)$ for the discrete measure $\mu = \frac{1}{N} \sum_{x \in \omega_N} \delta_x$. Generally, we are interested in finding point configurations and measures optimizing the energy for a given potential K .

2.2. Isotropic energies and frame potentials. A measure $\mu \in \mathcal{P}(\mathbb{S}^{d-1})$ is called *isotropic* if

$$(2.3) \quad \int_{\mathbb{S}^{d-1}} xx^T d\mu(x) = \frac{1}{d} I_d,$$

where I_d is the unit $d \times d$ matrix. This is equivalent to the statement that for any $y \in \mathbb{S}^{d-1}$,

$$(2.4) \quad \int_{\mathbb{S}^{d-1}} \langle x, y \rangle^2 d\mu(x) = \frac{1}{d}.$$

One can easily show that isotropic measures are precisely the minimizers of the 2-frame energy, see e.g. [BGMPV1].

Lemma 2.1. *For any $\mu \in \mathcal{P}(\mathbb{S}^{d-1})$,*

$$(2.5) \quad \int_{\mathbb{S}^{d-1}} \int_{\mathbb{S}^{d-1}} \langle x, y \rangle^2 d\mu(x) d\mu(y) \geq \frac{1}{d}.$$

The equality holds if and only if μ is isotropic.

Applying Lemma 2.1 to the uniform distribution over a discrete configuration $\omega_N = \{x_1, \dots, x_N\}$, $N \geq d$, we get the result of Benedetto and Fickus in the discrete setting [BF, Theorem 7.1]:

$$(2.6) \quad \sum_{i=1}^N \sum_{j=1}^N \langle x_i, x_j \rangle^2 \geq \frac{N^2}{d},$$

with equality if and only if for each $y \in \mathbb{R}^d$

$$\sum_{i=1}^N \langle x_i, y \rangle^2 = \frac{N}{d} \|y\|^2,$$

i.e. ω_N is a *unit norm tight frame*. The bound (2.6) is essentially a special case of the results of Welch [W] and Sidel'nikov [Sid, Corollary 1].

2.3. Semidefinite programming and three-point bounds. For a variety of optimization problems for two-point energies, the linear programming method serves as the main machinery (see, e.g., [Del, KL, Y, CK, BGMPV1]). Naturally, optimization problems for three-point energies should employ the three-point generalization of this method. This generalization was developed for the spherical case by Bachoc and Vallentin [BV] who used it to obtain new bounds for the kissing number problem. They produced a class of infinite matrices and associated polynomials of the form

$$(2.7) \quad (Y_m^d)_{i+1, j+1}(x, y, z) := Y_{m, i, j}^d(x, y, z) := P_i^{d+2m}(u) P_j^{d+2m}(v) Q_m^{d-1}(u, v, t),$$

where $m, i, j \in \mathbb{N}_0$, P_m^h is the Gegenbauer polynomial of degree m on \mathbb{S}^{h-1} , normalized by $P_m^h(1) = 1$, so that, in particular, $P_0^h(t) = 1$, $P_1^h(t) = t$, $P_2^h(t) = \frac{ht^2-1}{h-1}$, and

$$(2.8) \quad Q_m^{d-1}(u, v, t) = ((1-u^2)(1-v^2))^{\frac{m}{2}} P_m^{d-1} \left(\frac{t-uv}{\sqrt{(1-u^2)(1-v^2)}} \right).$$

For convenience, we include here the upper left 3×3 , 2×2 , and 1×1 submatrices of infinite matrices Y_0^d , Y_1^d , and Y_2^d respectively:

$$\begin{pmatrix} 1 & v & \frac{dv^2-1}{d-1} \\ u & uv & u\frac{dv^2-1}{d-1} \\ \frac{du^2-1}{d-1} & \frac{du^2-1}{d-1}v & \frac{du^2-1}{d-1}\frac{dv^2-1}{d-1} \end{pmatrix},$$

$$\begin{pmatrix} t-uv & v(t-uv) \\ u(t-uv) & uv(t-uv) \end{pmatrix}, \left(\frac{(d-1)(t-uv)^2 - (1-u^2)(1-v^2)}{d-2} \right).$$

Symmetrizing by all permutations π over the variables x , y , and z , Bachoc and Vallentin defined the following symmetric matrices and associated polynomials

$$(S_m^d)_{i+1,j+1}(x,y,z) := S_{m,i,j}^d(x,y,z) := \frac{1}{6} \sum_{\pi} Y_{m,i,j}^d(\pi(x), \pi(y), \pi(z)).$$

The following property of these matrices allows one to use them for optimization purposes. For any $\mu \in \mathcal{P}(\mathbb{S}^{d-1})$ and $e \in \mathbb{S}^{d-1}$, the infinite matrices

$$\int_{\mathbb{S}^{d-1}} \int_{\mathbb{S}^{d-1}} Y_m^d(x, y, e) d\mu(x) d\mu(y)$$

and

$$S_m^d(\mu) := \int_{\mathbb{S}^{d-1}} \int_{\mathbb{S}^{d-1}} \int_{\mathbb{S}^{d-1}} S_m^d(x, y, z) d\mu(x) d\mu(y) d\mu(z)$$

are positive semidefinite, that is, all principal minors (formed by finite submatrices) are non-negative. This property leads to the following energy minimization theorem (more details and justification can be found in Sections 3 and 4 of [BFGMPV1]).

Theorem 2.2. *Let $n \in \mathbb{N}_0$. For each $m \leq n$, let A_m be an infinite, symmetric, positive semidefinite matrix with finitely many non-zero entries, with the additional requirement that A_0 has only zeros in its first row and first column. Let*

$$K(x, y, z) = \sum_{m=0}^n \text{Tr}(S_m^d(x, y, z) A_m).$$

Then σ is a minimizer of I_K over probability measures on the sphere \mathbb{S}^{d-1} and $I_K(\sigma) = 0$.

Observe that when the matrices A_m are diagonal, the kernel K is simply a non-negative linear combination of the diagonal entries of the matrices S_m^d . Due to the nature of the bounds from Theorem 2.2 and their most common use, we will refer to them as semidefinite programming bounds throughout the paper.

3. THREE-POINT PACKING PROBLEM AND NEARLY ORTHOGONAL SETS

We use semidefinite programming bounds to prove a three-point packing bound generalizing the aforementioned result of Rosenfeld [Ros]. Observe that this also gives an alternative proof of Rosenfeld's result.

Theorem 3.1. *Assume for any three distinct points x, y, z of the set $\omega_N \subset \mathbb{S}^{d-1}$,*

$$\langle x, y \rangle \langle x, z \rangle \langle y, z \rangle \leq 0.$$

Then $N \leq 2d$. The equality $N = 2d$ is achieved only if ω_N is nearly orthogonal, i.e. a Rosenfeld set.

Proof. Following the notation from Section 2.3, we use

$$6 \frac{(d-1)^2}{d^2} S_{0,2,2}^d = 2(u^2 v^2 + u^2 t^2 + v^2 t^2) - \frac{4}{d}(u^2 + v^2 + t^2) + \frac{6}{d^2}$$

and

$$6 S_{1,1,1}^d = 6uv t - 2(u^2 v^2 + u^2 t^2 + v^2 t^2).$$

Considering the kernel $K = 6 \frac{(d-1)^2}{d^2} S_{0,2,2}^d + 6 S_{1,1,1}^d$, we notice that it can be written in the form $K(x, y, z) = \text{Tr}(S_0^d(x, y, z) A_0) + \text{Tr}(S_1^d(x, y, z) A_1)$, where A_0 and A_1 are infinite positive semidefinite rank-one matrices of the form $A_0 = 6 \frac{(d-1)^2}{d^2} a_0 a_0^T$ with $a_0 = (0, 0, 1, 0, 0, \dots)^T$ and $A_1 = 6 a_1 a_1^T$ with $a_1 = (0, 1, 0, 0, \dots)$, respectively. Summing up the values of this kernel over all triples of points in $\omega_N = \{x_1, \dots, x_N\}$ amounts to computing the expression $N^3 I_K(\mu_N)$, where μ_N is the probability measure assigning equal weights to points x_1, \dots, x_N . Due to Theorem 2.2, such summation must produce a non-negative number, whence

$$\begin{aligned} 0 &\leq \sum_{i,j,k=1}^N 6 \left(\frac{(d-1)^2}{d^2} S_{0,2,2}^d + S_{1,1,1}^d \right) (x_i, x_j, x_k) \\ &= \sum_{i,j,k=1}^N \left(6 \langle x_i, x_j \rangle \langle x_i, x_k \rangle \langle x_j, x_k \rangle - \frac{4}{d} (\langle x_i, x_j \rangle^2 + \langle x_i, x_k \rangle^2 + \langle x_j, x_k \rangle^2) + \frac{6}{d^2} \right). \end{aligned}$$

All triple product terms for distinct i, j, k are non-positive so we can eliminate them without changing the validity of the inequality (equality in this step is achieved if and only if ω_N is nearly orthogonal). The remaining terms are split into four cases: $i = j = k$, $i = j \neq k$, $i = k \neq j$, and $j = k \neq i$. The sum of triple product terms is N for the first group and $\sum_{i,j=1}^N \langle x_i, x_j \rangle^2 - N$ for the other three groups. The same sum shows up for double product terms. Overall, we have

$$\begin{aligned} 0 &\leq 6 \left(N + 3 \left(\sum_{i,j=1}^N \langle x_i, x_j \rangle^2 - N \right) \right) - \frac{4}{d} 3N \sum_{i,j=1}^N \langle x_i, x_j \rangle^2 + \frac{6}{d^2} N^3 \\ &= \left(18 - \frac{12N}{d} \right) \sum_{i,j=1}^N \langle x_i, x_j \rangle^2 - 12N + \frac{6N^3}{d^2}. \end{aligned}$$

If $18 - \frac{12N}{d}$ is non-negative, then $N \leq \frac{3}{2}d$ so the statement of the theorem is true. If it is negative, we note that $\sum_{i,j=1}^N \langle x_i, x_j \rangle^2$ is the frame energy of the set and must be at least $\frac{N^2}{d}$ by the discrete version of Lemma 2.1, i.e. inequality (2.6). Substituting this value in the inequality, we obtain

$$\left(18 - \frac{12N}{d} \right) \frac{N^2}{d} - 12N + \frac{6N^3}{d^2} \geq 0.$$

Dividing by $6N$ and factoring, we reach the inequality

$$(2 - N/d)(N/d - 1) \geq 0$$

that immediately implies the statement of the theorem.

Finally, we note that the size of ω_N is exactly $2d$ only when it is nearly orthogonal, i.e. a Rosenfeld set (and the converse is obvious). \square

Further analyzing the case of equality in the proof of Theorem 3.1, we observe that the frame energy of the set ω_N is necessarily N^2/d , which according to (2.6) means that ω_N has to be a tight frame.

Corollary 3.2. *Each nearly orthogonal set on \mathbb{S}^{d-1} of size $2d$ (Rosenfeld set) is a tight frame.*

Theorem 3.1 mirrors the classic packing bound: the size of a configuration in \mathbb{S}^{d-1} with $\langle x, y \rangle \leq 0$ for any two distinct points is no greater than $2d$ and the set of vertices of the d -dimensional crosspolytope is the unique (up to orthogonal transformations) set satisfying this bound [A, Sz, Ran]. This packing bound complements the initial result of Davenport and Hajós who showed that the size of a configuration in \mathbb{S}^{d-1} with $\langle x, y \rangle < 0$ for any two distinct points is no greater than $d + 1$ [DH]. The result of Davenport and Hajós can be easily extended to the three-point case.

Theorem 3.3. *Assume for any three distinct points x, y, z of the set $\omega_N \subset \mathbb{S}^{d-1}$,*

$$\langle x, y \rangle \langle x, z \rangle \langle y, z \rangle < 0.$$

Then $N \leq d + 1$.

Proof. Fix a point $x \in \omega_N$. Since changing any point $y \in \omega_N$ to its opposite $-y$ does not affect the condition of the theorem, we may, without loss of generality, assume that $\langle x, y \rangle < 0$ for each $y \in \omega_N \setminus \{x\}$. The condition $\langle x, y \rangle \langle x, z \rangle \langle y, z \rangle < 0$ then implies that $\langle y, z \rangle < 0$ for any distinct $y, z \in \omega_N \setminus \{x\}$. Thus, all the inner products in ω_N are negative. By the result of Davenport and Hajós [DH] mentioned above, the size of the set is no greater than $d + 1$. \square

Finally, it is a simple fact that if $\langle x, y \rangle < -\varepsilon$ for each distinct $x, y \in \omega_N$, then $N \leq 1 + \frac{1}{\varepsilon}$, which is independent of the dimension d . This can be quickly derived from the inequality $\|\sum x_i\|^2 \geq 0$. This fact also easily generalizes to the multivariate setting.

Lemma 3.4. *Let $\varepsilon > 0$. Assume for any three distinct points x, y, z of the set $\omega_N \subset \mathbb{S}^{d-1}$,*

$$\langle x, y \rangle \langle x, z \rangle \langle y, z \rangle \leq -\varepsilon.$$

Then $N \leq 1 + \frac{1}{\varepsilon}$.

Proof. As the value of $\langle x, y \rangle \langle x, z \rangle \langle y, z \rangle$ doesn't change when any of the vectors is changed to its opposite, arguing as in the proof of Theorem 3.3, we may assume that all inner products between distinct elements of ω_N are negative. But $|\langle x, y \rangle \langle x, z \rangle \langle y, z \rangle| \leq |\langle x, y \rangle|$, therefore $\langle x, y \rangle < -\varepsilon$ for each distinct $x, y \in \omega_N$. Thus, the aforementioned fact implies the statement of the lemma. \square

The Erdős problem on nearly orthogonal sets gave rise to active investigations of orthogonal representations of graphs, see e.g. [Dea]. Since the proof of Theorem 3.1 demonstrates that the machinery of semidefinite bounds is effective for the Erdős problem, it would be interesting to find out whether any other problems regarding orthogonal representations of graphs can be solved in a similar manner. A clear obstacle to this approach is the complexity of functions (2.8) involved in k -point semidefinite bounds.

4. MINIMAL ENERGY FOR MULTIVARIATE p -FRAME POTENTIALS

We now turn our attention to the multivariate p -frame energy, i.e. the energy with the potential $K(x, y, z) = |\langle x, y \rangle \langle x, z \rangle \langle y, z \rangle|^p = |uvt|^p$ for $p > 0$. While this section primarily focuses on the case $p \leq 1$, we first quickly address the case when the p -frame potential is a polynomial. Corollary 5.2 in [BFGMPV] implies

Proposition 4.1. *If $p \in 2\mathbb{N}$, then σ minimizes the three-point p -frame energy.*

As the first step in understanding the set of minimizers for $p = 1$, we give a description of minimizing measures for the potential uvt .

Lemma 4.2. *For any $\mu \in \mathcal{P}(\mathbb{S}^{d-1})$,*

$$I_{uvt}(\mu) = \int_{\mathbb{S}^{d-1}} \int_{\mathbb{S}^{d-1}} \int_{\mathbb{S}^{d-1}} \langle x, y \rangle \langle y, z \rangle \langle z, x \rangle d\mu(x) d\mu(y) d\mu(z) \geq \frac{1}{d^2}$$

and the equality holds if and only if μ is isotropic.

We shall give two different proofs of this fact, as both are quite instructive: one based on linear algebra and another one based on the semidefinite programming bounds of Theorem 2.2.

Proof 1 (Linear algebra). If we denote $x = (x^{(1)}, \dots, x^{(d)})$, then

$$uvt = \sum_{k,l,m=1}^d x^{(k)} x^{(l)} y^{(l)} y^{(m)} z^{(m)} z^{(k)}.$$

Therefore, setting $a_{k,l} = \int_{\mathbb{S}^{d-1}} x^{(k)} x^{(l)} d\mu(x)$, we see that

$$I_{uvt}(\mu) = \sum_{k,l,m=1}^d a_{k,l} a_{l,m} a_{m,k} = \text{Tr}(A^3),$$

where $A = (a_{k,l}) = \int_{\mathbb{S}^{d-1}} xx^T d\mu(x)$ is obviously positive semidefinite.

For any positive semidefinite $d \times d$ matrix A with eigenvalues $\lambda_i \geq 0$, using Hölder's inequality, one obtains

$$\text{Tr}(A) = \sum_{i=1}^d \lambda_i \leq \left(\sum_{i=1}^d \lambda_i^3 \right)^{\frac{1}{3}} \cdot d^{\frac{2}{3}} = \left(\text{Tr}(A^3) \right)^{\frac{1}{3}} \cdot d^{\frac{2}{3}}, \quad \text{i.e.} \quad \text{Tr}(A^3) \geq \frac{(\text{Tr}(A))^3}{d^2},$$

with equality if and only if A is a multiple of the identity.

In our case, $\text{Tr}(A) = \sum_{i=1}^d \int_{\mathbb{S}^{d-1}} (x^{(i)})^2 d\mu(x) = 1$, therefore $I_{uvt}(\mu) = \text{Tr}(A^3) \geq \frac{1}{d^2}$, and the equality is achieved if and only if $A = \frac{1}{d}I_d$, i.e. μ is isotropic. \square

Proof 2 (Semidefinite programming). From the proof of Theorem 3.1, we see that

$$uvt = \frac{(d-1)^2}{d^2} S_{0,2,2}^d + S_{1,1,1}^d + \frac{2}{3d}(u^2 + v^2 + t^2) - \frac{1}{d^2}.$$

When integrating the right hand side, the values for $S_{0,2,2}^d$ and $S_{1,1,1}^d$ are non-negative due to Theorem 2.2. The integral of $u^2 + v^2 + t^2$ is at least $\frac{3}{d}$ by Lemma 2.1. Overall, we get

$$\int_{\mathbb{S}^{d-1}} \int_{\mathbb{S}^{d-1}} \int_{\mathbb{S}^{d-1}} \langle x, y \rangle \langle x, z \rangle \langle y, z \rangle d\mu(x) d\mu(y) d\mu(z) \geq \frac{2}{3d} \cdot \frac{3}{d} - \frac{1}{d^2} = \frac{1}{d^2}.$$

The equality can hold only when μ is isotropic due to Lemma 2.1.

On the other hand, using equations (2.3)–(2.4), it is easy to check that the energy is precisely $\frac{1}{d^2}$ for all isotropic measures (or that the integrals of both $S_{0,2,2}^d$ and $S_{1,1,1}^d$ vanish on isotropic measures). Indeed,

$$\begin{aligned} I_{uv\ell}(\mu) &= \int_{\mathbb{S}^{d-1}} \int_{\mathbb{S}^{d-1}} \int_{\mathbb{S}^{d-1}} x^T y y^T z z^T x \, d\mu(x) d\mu(y) d\mu(z) \\ &= \frac{1}{d} \int_{\mathbb{S}^{d-1}} \int_{\mathbb{S}^{d-1}} x^T y y^T x \, d\mu(x) d\mu(y) \\ &= \frac{1}{d} \int_{\mathbb{S}^{d-1}} \int_{\mathbb{S}^{d-1}} |\langle x, y \rangle|^2 \, d\mu(x) d\mu(y) = \frac{1}{d^2}. \end{aligned}$$

□

We remark that this result (although without the full characterization of minimizers) has also been proved in Corollary 5.2 of [BFGMPV] by a different method.

As a direct consequence of Lemma 4.2, we can describe minimizers of the 1-frame potential as follows.

Corollary 4.3. *For any $\mu \in \mathcal{P}(\mathbb{S}^{d-1})$,*

$$\int_{\mathbb{S}^{d-1}} \int_{\mathbb{S}^{d-1}} \int_{\mathbb{S}^{d-1}} |\langle x, y \rangle \langle x, z \rangle \langle y, z \rangle| \, d\mu(x) d\mu(y) d\mu(z) \geq \frac{1}{d^2}$$

and the equality holds if and only if μ is isotropic and $\langle x, y \rangle \langle x, z \rangle \langle y, z \rangle \geq 0$ for any points x, y, z in the support of μ .

Using Corollary 4.3, we can solve the problem for $0 < p < 1$ and show that the orthonormal basis is a unique (up to central symmetry and rotations) minimizer.

Theorem 4.4. *For $0 < p < 1$,*

$$\int_{\mathbb{S}^{d-1}} \int_{\mathbb{S}^{d-1}} \int_{\mathbb{S}^{d-1}} |\langle x, y \rangle \langle x, z \rangle \langle y, z \rangle|^p \, d\mu(x) d\mu(y) d\mu(z) \geq \frac{1}{d^2}$$

and the equality holds if and only if μ is a uniform distribution over an orthonormal basis (up to central symmetry).

Proof. For $0 < p < 1$ and any $\mu \in \mathcal{P}(\mathbb{S}^{d-1})$,

$$|\langle x, y \rangle \langle x, z \rangle \langle y, z \rangle|^p \geq |\langle x, y \rangle \langle x, z \rangle \langle y, z \rangle|,$$

so the lower bound follows from Corollary 4.3. The bound is sharp if and only if μ minimizes the three-point 1-frame energy and for all x, y, z from $\text{supp}(\mu)$, $\langle x, y \rangle \langle x, z \rangle \langle y, z \rangle = 0$ or 1. This condition is not satisfied for the triple (x, x, y) if there are two points x, y such that $\langle x, y \rangle \notin \{0, -1, 1\}$. Therefore, any distinct points in $\text{supp}(\mu)$ are orthogonal or antipodal. The support of an isotropic μ must span \mathbb{R}^d so, due to (2.4), $\mu(\{e_j, -e_j\}) = \frac{1}{d}$ for $j = 1, \dots, d$, for some orthonormal basis e_1, \dots, e_d . □

Remark 4.5. We observe that Theorem 4.4 actually holds for more general kernels than just $K(x, y, z) = |\langle x, y \rangle \langle x, z \rangle \langle y, z \rangle|^p$. The proof above shows that it holds for any kernel K such that $K(x, y, z) \geq |\langle x, y \rangle \langle x, z \rangle \langle y, z \rangle|$ for all $x, y, z \in \mathbb{S}^{d-1}$ with equality only for those triples where $\langle x, y \rangle \langle x, z \rangle \langle y, z \rangle = 0$ or 1.

In particular, it holds for all kernels of the form

$$K(x, y, z) = f(|\langle x, y \rangle|)f(|\langle x, z \rangle|)f(|\langle y, z \rangle|),$$

where a (Borel measurable) function $f : [0, 1] \rightarrow \mathbb{R}$ satisfies $f(0) = 0$, $f(1) = 1$, and f is concave (or, more generally $f(t) > t$ for $0 < t < 1$).

The main difficulty in solving the problem for $p = 1$ is in characterizing all isotropic measures such that $\langle x, y \rangle \langle x, z \rangle \langle y, z \rangle \geq 0$ for any points x, y, z in their support. Note that Rosenfeld sets are discrete minimizers of the 1-frame energy as they are tight frames (Corollary 3.2) and satisfy this condition. The general problem of describing all minimizers is highly non-trivial since, as mentioned above, even the problem of characterizing all Rosenfeld sets is wide-open for $d \geq 5$. However, we can provide a complete description of minimizers for $d = 2$.

Theorem 4.6. *For any $\mu \in \mathcal{P}(\mathbb{S}^1)$,*

$$\int_{\mathbb{S}^1} \int_{\mathbb{S}^1} \int_{\mathbb{S}^1} |\langle x, y \rangle \langle x, z \rangle \langle y, z \rangle| d\mu(x) d\mu(y) d\mu(z) \geq \frac{1}{4}$$

and the equality holds if and only if μ is a convex combination of uniform distributions over two (not necessarily distinct) orthonormal bases (up to central symmetry).

Proof. Let μ be a minimizer and assume there is $x \in \text{supp}(\mu)$ such that $x^\perp \cap \text{supp}(\mu) = \emptyset$. Note that the potential is invariant under central symmetry for any of its arguments. This means that we can rearrange the measure by switching points to their opposites so that $\text{supp}(\mu)$ is entirely contained in an open half-circle centered at x . Assume the left endpoint of the support now is x_1 and the right endpoint is x_2 . Since both $\langle x, x_1 \rangle$ and $\langle x, x_2 \rangle$ are positive, $\langle x_1, x_2 \rangle \geq 0$, due to Corollary 4.3. Take the midpoint y of the circular arc between x_1 and x_2 , contained in the same half-circle around x . The angle between x_1 and x_2 is no greater than $\frac{\pi}{2}$ so for any $z \in \text{supp}(\mu)$, $\langle z, y \rangle \geq \frac{1}{\sqrt{2}}$. By (2.4), since μ is isotropic

$$\int_{\mathbb{S}^1} |\langle z, y \rangle|^2 d\mu(z) = \frac{1}{2}.$$

This may happen only if $\langle z, y \rangle$ is precisely $\frac{1}{\sqrt{2}}$ for any $z \in \text{supp}(\mu)$ and $\langle x_1, x_2 \rangle = 0$. However, x cannot coincide with x_1 or x_2 because it does not have an orthogonal counterpart in the support and, therefore, $\langle x, y \rangle$ is definitely not $\frac{1}{\sqrt{2}}$ which contradicts our assumption.

We conclude that any $x \in \text{supp}(\mu)$ has an orthogonal counterpart. Now we assume that there are at least three pairwise non-orthogonal and pairwise non-opposite points x, y, z in the support of μ . Each of them has an orthogonal counterpart $x^\perp, y^\perp, z^\perp \in \text{supp}(\mu)$. We claim that among these six points, there are three with the negative product of pairwise scalar products. Without loss of generality, we can assume, by switching to opposites if needed, that pairwise angles between three of the points, say, x, y, z are all acute and y is between x and z on the circle. Then $\langle x, y^\perp \rangle \langle x, z \rangle \langle y^\perp, z \rangle < 0$, a contradiction, so we may have no more than two pairwise non-orthogonal and pairwise non-opposite points.

Again switching to opposite points if necessary, we assume that the support of μ consists of only four points (the case of two points is handled later): x, y , and their orthogonal counterparts x^\perp and y^\perp . Assume $\mu(x) = \alpha_1$, $\mu(x^\perp) = \alpha_2$, $\mu(y) = \beta_1$, $\mu(y^\perp) = \beta_2$. Using (2.4) for an arbitrary $w \in \mathbb{S}^{d-1}$, we get

$$\frac{1}{2} = \int_{\mathbb{S}^1} |\langle z, w \rangle|^2 d\mu(z) = \alpha_1 \langle w, x \rangle^2 + \alpha_2 (1 - \langle w, x \rangle^2) + \beta_1 \langle w, y \rangle^2 + \beta_2 (1 - \langle w, y \rangle^2).$$

Hence there exists a linear dependence between 1 , $\langle w, x \rangle^2$, and $\langle w, y \rangle^2$. This may happen only when y is orthogonal or opposite to x or the dependence is trivial. In the latter case, $\alpha_1 = \alpha_2$ and $\beta_1 = \beta_2$ so μ is precisely a convex combination of uniform distributions over two orthonormal bases.

The remaining case is when the support of μ consists of only x and x^\perp and, by the same condition, their weights must be equal. \square

There are certain distinctions between the behavior of the p -frame potentials in the two- and three-input cases: in particular, while in the two-input case the uniform distribution over an orthonormal basis is the unique (up to symmetries) minimizer of the p -frame energy in the range $0 < p < 2$, the discussion in this section suggests that for the three-input energy this is only true for $0 < p < 1$, as Theorem 4.6 indicates that one does not have uniqueness for $p = 1$, and an argument similar to Theorem 4.4 would then show that the orthonormal basis is not a minimizer for $p > 1$. This difference might be partially explained by the fact that the degree of each of the variables in the p -frame potential in the three-input case is twice as large.

In the two-input case, the general conjecture in [BGMPV2] claims that all minimizers of the p -frame energy are discrete when $p \notin 2\mathbb{N}$. It is therefore natural to conjecture that all minimizers of the three-point p -frame energy are also discrete, at least when $p \notin \mathbb{N}$.

4.1. Discrete three-input p -frame energy for $d < N < 2d$ and $N = kd$. We finish this section with a brief discussion of the discrete version of the problem. In the classical two-input case, the problem of finding optimal configurations for the discrete p -frame energy with fixed N has attracted considerable attention recently, see e.g. [CGGKO, GP, XX]. The case $N = d + 1$ is particularly interesting, since it presents natural candidates for minimizers (regular simplex, orthonormal basis with one repeated vector, products of lower-dimensional simplices and bases, etc), as well as, more generally, the case of $d < N < 2d$.

While this is not the main focus of our paper, below we present some results on the minimization of the discrete three-input p -frame energy with fixed N , which draw on connections of the problem to the two-input case and nearly orthogonal sets. To this end, we start with Theorem 4.7:

Theorem 4.7. *If a nearly orthogonal set is a (unique) minimizer of the $2p$ -frame energy among finite sets of a fixed size in \mathbb{S}^{d-1} , then it is a (unique) minimizer of the three-point p -frame energy among finite sets of a fixed size in \mathbb{S}^{d-1} .*

Proof. Let $\omega_N = \{x_1, \dots, x_N\} \subset \mathbb{S}^{d-1}$. Similarly to the proof of Theorem 3.1, we have

$$\begin{aligned} \sum_{i,j,k=1}^N |\langle x_i, x_j \rangle \langle x_j, x_k \rangle \langle x_k, x_i \rangle|^p &= \sum_{i \neq j, j \neq k, k \neq i} |\langle x_i, x_j \rangle \langle x_j, x_k \rangle \langle x_k, x_i \rangle|^p \\ &\quad + 3 \sum_{i,j=1}^N |\langle x_i, x_j \rangle|^{2p} - 2N \\ &\geq 3 \sum_{i,j=1}^N |\langle x_i, x_j \rangle|^{2p} - 2N, \end{aligned}$$

where the exact equality holds precisely for nearly orthogonal sets. The statement of the theorem immediately follows. \square

Theorem 4.7 allows us to find minimizers of three-point p -frame energies in several concrete cases.

Corollary 4.8. *For $p \in (0, \frac{\ln 3}{2 \ln 2})$, the unique minimizer of the three-point p -frame energy among sets in \mathbb{S}^{d-1} of size $d+1$ is the set of points from the same orthonormal basis with exactly one point repeated twice.*

Proof. Due to Theorem 4.7, the corollary follows from a partial case of [XX, Theorem 1.2], which states that the two-input p -frame energy with $N = d + 1$ points is uniquely minimized by an orthonormal basis with one repeated point when $p \in (0, \frac{\ln 3}{\ln 2})$, as well as the fact that this minimizing set is nearly orthogonal. \square

Remark 4.9. The paper [XX] completely classifies the minimizers of the p -frame energy among sets of size $d + 1$ for all values of $p \in (0, 2)$. Unfortunately, Theorem 4.7 is not applicable to other minimizers from this classification as they are not nearly orthogonal sets.

A similar result can be obtained for $d + 1 < N < 2d$:

Corollary 4.10. *For $1 \leq m < d$ and $p \in (0, \frac{\ln(2m+1) - \ln(2m)}{\ln(m+1) - \ln m})$, the unique (up to central symmetry) minimizer of the three-point p -frame energy among sets of size $d + m$ in \mathbb{S}^{d-1} is the set of points from an orthonormal basis with exactly m points repeated twice.*

Proof. Similarly to Corollary 4.8, the statement follows from [GP, Theorem 3.2] and the fact that the minimizing set is nearly orthogonal. \square

Note also that Corollary 4.3 and Theorem 4.4 give the following observation about minimizers of the discrete energy for $N = kd$, $k \geq 1$, and $0 < p \leq 1$:

Corollary 4.11. *For $0 < p < 1$, the unique (up to central symmetry) minimizer of the three-point p -frame energy among sets of size kd in \mathbb{S}^{d-1} is the set of points from an orthonormal basis with every point repeated k times. When $p = 1$ such sets are also minimizers (although not unique).*

Proof. Values of the discrete p -frame energy on sets of size kd coincide with the values of integral p -frame energy on the normalized counting measures of such sets. Hence the first statement immediately follows from Theorem 4.4. For $p = 1$, minimization follows from Corollary 4.3. One can further see from Corollary 4.3 that taking repeated copies of two different orthonormal bases yields another minimizer. \square

5. ISOTROPIC MEASURES WITHOUT OBTUSE ANGLES

Isotropic measures with no obtuse angles between any two points in their support clearly minimize the three-point 1-frame energy (due to Corollary 4.3). In this section we show that only orthonormal bases satisfy this condition.

Theorem 5.1. *If $\langle x, y \rangle \geq 0$ for any x, y in the support of an isotropic measure $\mu \in \mathcal{P}(\mathbb{S}^{d-1})$, then μ is a uniform distribution over an orthonormal basis.*

For the proof of Theorem 5.1, we need three ingredients: the spherical Jung inequality, the procedure of lifting of balanced isotropic measures to a higher dimension, and the linear programming bound on the diameter of a balanced isotropic measure.

Jung's inequality [Jung] shows that the simplex has the maximal circumradius for a Euclidean set of a given diameter. For the first ingredient of the proof, we use the spherical version of this inequality proved by Dekster [D].

Theorem 5.2 (Dekster). *If the spherical distance, i.e. $\arccos(\langle x, y \rangle)$, between any two points of a compact set $C \subset \mathbb{S}^{d-1}$ is not greater than D , $0 \leq D \leq 2 \sin^{-1} \sqrt{\frac{d}{2d-2}}$, then C can be covered by a spherical cap whose radius is a circumradius of a regular simplex in \mathbb{S}^{d-1} with spherical edge length D .*

Note that a regular simplex with edge length $\frac{\pi}{2}$ is formed by the endpoints of an orthonormal basis e_1, \dots, e_d in \mathbb{S}^{d-1} . Denote $z = \frac{1}{\sqrt{d}}(e_1 + \dots + e_d)$. The points on the circumsphere of the simplex are defined by $\langle x, z \rangle = \frac{1}{\sqrt{d}}$. Theorem 5.2 then implies that for any spherical set, where $\langle x, y \rangle \geq 0$ for any pair of points x and y , there exists $z \in \mathbb{S}^{d-1}$ such that $\langle x, z \rangle \geq \frac{1}{\sqrt{d}}$ for any x in the set.

For the next ingredient of the proof of Theorem 5.1, we describe the procedure of lifting of an isotropic measure to a higher dimension. Assume $z \in \mathbb{S}^d$ and take for \mathbb{S}^{d-1} the intersection of \mathbb{S}^d and the hyperplane through the origin orthogonal to z . Define $f : \mathbb{S}^{d-1} \rightarrow \mathbb{S}^d$ as follows:

$$f(x) = \sqrt{\frac{d}{d+1}}x + \frac{1}{\sqrt{d+1}}z.$$

For $\mu \in \mathcal{P}(\mathbb{S}^{d-1})$ define the lifted measure $\mu_l \in \mathcal{P}(\mathbb{S}^d)$ as the pushforward of μ under f .

Lemma 5.3. *A probability measure μ is a balanced isotropic measure in \mathbb{S}^{d-1} if and only if μ_l is an isotropic measure in \mathbb{S}^d .*

Proof. Any $y \in \mathbb{S}^d$ can be represented as $\alpha_y y' + \beta_y z$, where $y' \in \mathbb{S}^{d-1}$ and $\alpha_y^2 + \beta_y^2 =$

1. For any $x' \in \text{supp}(\mu)$, letting $x = f(x')$, $\alpha_x = \sqrt{\frac{d}{d+1}}$ and $\beta_x = \frac{1}{\sqrt{d+1}}$. Then

$$\begin{aligned} \int_{\mathbb{S}^d} \langle x, y \rangle^2 d\mu_l(x) &= \int_{\mathbb{S}^{d-1}} \left(\sqrt{\frac{d}{d+1}} \alpha_y \langle x', y' \rangle + \frac{1}{\sqrt{d+1}} \beta_y \right)^2 d\mu(x') \\ &= \frac{1}{d+1} \int_{\mathbb{S}^{d-1}} \left(d\alpha_y^2 \langle x', y' \rangle^2 + 2\sqrt{d}\alpha_y\beta_y \langle x', y' \rangle + \beta_y^2 \right) d\mu(x') \\ &= \frac{d\alpha_y^2}{d+1} \int_{\mathbb{S}^{d-1}} \langle x', y' \rangle^2 d\mu(x') \\ &\quad + \frac{2\sqrt{d}\alpha_y\beta_y}{d+1} \int_{\mathbb{S}^{d-1}} \langle x', y' \rangle d\mu(x') + \frac{\beta_y^2}{d+1}. \end{aligned}$$

If μ is balanced and isotropic, then, due to (2.4), for any $y' \in \mathbb{S}^{d-1}$,

$$\begin{aligned} & \frac{d\alpha_y^2}{d+1} \int_{\mathbb{S}^{d-1}} \langle x', y' \rangle^2 d\mu(x') + \frac{2\sqrt{d}\alpha_y\beta_y}{d+1} \int_{\mathbb{S}^{d-1}} \langle x', y' \rangle d\mu(x') + \frac{\beta_y^2}{d+1} \\ &= \frac{d\alpha_y^2}{d+1} \cdot \frac{1}{d} + 0 + \frac{\beta_y^2}{d+1} \\ &= \frac{\alpha_y^2 + \beta_y^2}{d+1} = \frac{1}{d+1}, \end{aligned}$$

so (2.4) is satisfied and μ_l is isotropic in \mathbb{S}^d .

For the other direction, assume μ_l is isotropic. Taking $y \in \mathbb{S}^{d-1}$, i.e. $\alpha_y = 1$ and $\beta_y = 0$, we get

$$\frac{1}{d+1} = \frac{d}{d+1} \int_{\mathbb{S}^{d-1}} \langle x', y' \rangle^2 d\mu(x'),$$

so, again by (2.4), μ is isotropic in \mathbb{S}^{d-1} . Now we use that both μ_l and μ are isotropic and take an arbitrary $\alpha_y \neq 0, 1$ to get

$$\frac{1}{d+1} = \frac{d\alpha_y^2}{d+1} \cdot \frac{1}{d} + \frac{2\sqrt{d}\alpha_y\beta_y}{d+1} \int_{\mathbb{S}^{d-1}} \langle x', y' \rangle d\mu(x') + \frac{\beta_y^2}{d+1},$$

so

$$\frac{2\sqrt{d}\alpha_y\beta_y}{d+1} \int_{\mathbb{S}^{d-1}} \langle x', y' \rangle d\mu(x') = 0,$$

meaning that μ is balanced. \square

Remark 5.4. Essentially the same lifting construction for discrete sets was used, for example, in [BGOY] to construct two-distance tight frames and in [Ball1] to make use of the geometric Brascamp-Lieb inequality.

The next result provides the bound on the diameter of the support of a balanced isotropic measure confined to a sphere. The spirit of the proof resembles the linear programming approach, where a carefully constructed polynomial typically leads to a required bound. Similar optimization results for isotropic measures were also obtained in [GL].

Theorem 5.5. *Let $\mu \in \mathcal{P}(\mathbb{S}^{d-1})$ be a balanced isotropic measure. If $\langle x, y \rangle \geq -\frac{1}{d}$ for any $x, y \in \text{supp}(\mu)$, then μ is a uniform distribution over a regular simplex.*

Proof. Define $P(t) = (t-1)(t+\frac{1}{d})$ and note that $P(\langle x, y \rangle) \leq 0$ for any $x, y \in \text{supp}(\mu)$. Then

$$(5.1) \quad \int_{\mathbb{S}^{d-1}} \int_{\mathbb{S}^{d-1}} P(\langle x, y \rangle) d\mu(x) d\mu(y) \leq 0$$

and it is strictly less than 0 if $P(\langle x, y \rangle) < 0$ for some $x, y \in \text{supp}(\mu)$.

On the other hand, $P(\langle x, y \rangle) = \langle x, y \rangle^2 - \frac{d-1}{d} \langle x, y \rangle - \frac{1}{d}$. Since μ is balanced and isotropic,

$$\int_{\mathbb{S}^{d-1}} \int_{\mathbb{S}^{d-1}} \langle x, y \rangle d\mu(x) d\mu(y) = 0 \quad \text{and} \quad \int_{\mathbb{S}^{d-1}} \int_{\mathbb{S}^{d-1}} \langle x, y \rangle^2 d\mu(x) d\mu(y) = \frac{1}{d}.$$

This means (5.1) is sharp for μ and $P(\langle x, y \rangle) = 0$ for any $x, y \in \text{supp}(\mu)$. In particular, $\langle x, y \rangle$ must be $-\frac{1}{d}$ for any distinct x and y in the support of μ . Therefore, $\text{supp}(\mu)$ is a subset of the $d+1$ vertices of a regular simplex in \mathbb{S}^{d-1} . Among linear combinations of these vertices, only those with equal coefficients are 0. Given that

μ is balanced, it must be a uniform distribution over the set of vertices of a regular simplex. \square

Remark 5.6. Theorem 5.5 implies that the spherical diameter of the support of a balanced isotropic measure on the unit sphere is at least $\arccos(-\frac{1}{d})$. Moreover, if the diameter is precisely $\arccos(-\frac{1}{d})$, the measure is necessarily a uniform distribution over a regular simplex.

We now have all the ingredients necessary for the proof of Theorem 5.1.

Proof of Theorem 5.1. For the first step, we use Theorem 5.2 and, as described above, find $z \in \mathbb{S}^{d-1}$ such that $\langle x, z \rangle \geq \frac{1}{\sqrt{d}}$ for all $x \in \text{supp}(\mu)$. By Equality (2.4),

$$\int_{\mathbb{S}^{d-1}} \langle x, z \rangle^2 d\mu(x) = \frac{1}{d},$$

so $\langle x, z \rangle = \frac{1}{\sqrt{d}}$ for all $x \in \text{supp}(\mu)$.

This means μ is a lifted measure for $\mu' \in \mathcal{P}(\mathbb{S}^{d-2})$. By Lemma 5.3, μ' is balanced and isotropic. If $x, y \in \text{supp}(\mu)$, then $x = \sqrt{\frac{d-1}{d}}x' + \frac{1}{\sqrt{d}}z$ and $y = \sqrt{\frac{d-1}{d}}y' + \frac{1}{\sqrt{d}}z$, where $x', y' \in \text{supp}(\mu')$. The condition $\langle x, y \rangle \geq 0$ is equivalent to $\langle x', y' \rangle \geq -\frac{1}{d-1}$. Then μ' satisfies Theorem 5.5 and must be a uniform distribution over a regular simplex in \mathbb{S}^{d-2} . Lifting the simplex to \mathbb{S}^{d-1} we get that μ must be a uniform distribution over an orthonormal basis. \square

Remark 5.7. Similarly to Remark 5.6, Theorem 5.1 implies that the spherical diameter of the support of an isotropic measure on the unit sphere is at least $\frac{\pi}{2}$ and, if the diameter is precisely $\frac{\pi}{2}$, the measure is necessarily a uniform distribution over an orthonormal basis.

6. SUPPORT OF A MINIMIZER FOR 1-FRAME ENERGY HAS DIMENSION NOT GREATER THAN $d - 2$

One of the main results of [BGMPV2] states that the support of a minimizer of a two-point p -frame energy has empty interior, when p is not an even integer. Here we prove a similar result for the three-point 1-frame energy.

Lemma 6.1. *If μ is a minimizer of the three-point 1-frame energy among all measures in $\mathcal{P}(\mathbb{S}^{d-1})$, then for any $x \in \text{supp}(\mu)$ there is $y \in \text{supp}(\mu)$ such that $\langle x, y \rangle = 0$.*

Proof. Assume there is x in the support of μ such that none of the other points in $\text{supp}(\mu)$ are orthogonal to x . Rearrange the measure so that all points in its support form an acute angle with x by switching points to their opposites if necessary. Now if $y, z \in \text{supp}(\mu)$, both $\langle x, y \rangle$ and $\langle x, z \rangle$ are positive. As was established in Corollary 4.3, $\langle x, y \rangle \langle x, z \rangle \langle y, z \rangle \geq 0$ for any triple of points in the support of μ , hence, also $\langle y, z \rangle \geq 0$. The rearranged measure satisfies Theorem 5.1 so it must be a uniform distribution over an orthonormal basis. Subsequently, the initial assumption is false. \square

Theorem 6.2. *If μ is a minimizer of the three-point 1-frame energy among all measures in $\mathcal{P}(\mathbb{S}^{d-1})$, then the Hausdorff dimension of $\text{supp}(\mu)$ is no greater than $d - 2$.*

Proof. Fix $0 < \vartheta < \pi/4$. We consider an arbitrary spherical cap C of radius ϑ . It is sufficient to show that the $(d-2)$ -dimensional Hausdorff content of $C \cap \text{supp}(\mu)$ is finite. Indeed, since the sphere can be covered by a finite number of caps with radius ϑ , the total $(d-2)$ -dimensional Hausdorff content of $\text{supp}(\mu)$ is finite and the dimension of $\text{supp}(\mu)$ is no greater than $d-2$.

Take an arbitrary x from $C \cap \text{supp}(\mu)$. By Lemma 6.1, there is $y \in \text{supp}(\mu)$ such that $\langle x, y \rangle = 0$. Assume there are two points x_1, x_2 in $C \cap \text{supp}(\mu)$ such that $\langle x_1, y \rangle > 0$ and $\langle x_2, y \rangle < 0$. The spherical distance between x_1 and x_2 is no greater than $2\vartheta < \pi/2$ so $\langle x_1, x_2 \rangle > 0$. Therefore, we have found three points x_1, x_2, y in $\text{supp}(\mu)$ such that $\langle x_1, y \rangle \langle x_2, y \rangle \langle x_1, x_2 \rangle < 0$. This contradicts Corollary 4.3.

This means all points $z \in C \cap \text{supp}(\mu)$ simultaneously satisfy either $\langle z, y \rangle \geq 0$ or $\langle z, y \rangle \leq 0$. We denote by H_x a halfspace defined either by $\langle z, y \rangle \geq 0$ or $\langle z, y \rangle \leq 0$ such that $C \cap \text{supp}(\mu) \subset H_x$. Then $C \cap \text{supp}(\mu)$ is a subset of the convex set H defined as the intersection of H_x taken for all $x \in C \cap \text{supp}(\mu)$. Moreover, each point $x \in C \cap \text{supp}(\mu)$ is a boundary point of H because it belongs to a hyperplane $\langle x, y \rangle = 0$ defining H_x . Since $C \cap H$ is convex, Corollary 7.2 tells us its boundary has finite $(d-2)$ -dimensional Hausdorff content so the content of $C \cap \text{supp}(\mu)$ is finite as well. \square

7. APPENDIX

For $s > 0$, we denote the s -dimensional Hausdorff measure by \mathcal{H}_s . The fact that the Hausdorff dimension of the boundary of a convex d -dimensional set cannot be larger than $d-1$ seems to be well-known. However, we were not able to find a concrete reference for this statement in the literature so we suggest a short proof here.

Proposition 7.1. *If $K \subset \mathbb{R}^d$ is a non-empty, compact, convex set, then $\mathcal{H}_{d-1}(\partial K)$ is finite.*

Proof. If there is a set $d+1$ points in K that is not contained in an affine hyperplane, K contains the simplex with these points as vertices, so K has an interior point. If no such subset exists, then K is contained by a hyperplane H of dimension $d-1$, so, since K is compact, $\mathcal{H}_{d-1}(\partial K) \leq \mathcal{H}_{d-1}(K) < \infty$.

Now, without loss of generality, assume $0 \in K^\circ$, let $p : \mathbb{S}^{d-1} \rightarrow \partial K$ be the central projection from the unit sphere onto the boundary of K , i.e. $p(x) \in x\mathbb{R}_+ \cap \partial K$. Note that this function is a bijection.

For any $x, y \in \mathbb{S}^{d-1}$, we have that

$$\begin{aligned} \frac{\|p(x) - p(y)\|}{\|x - y\|} &= \frac{\left| \|p(x)\|x - \|p(y)\|x + \|p(y)\|x - \|p(y)\|y \right|}{\|x - y\|} \\ &\leq \frac{\left| \|p(x)\| - \|p(y)\| \right| \|x\| + \|p(y)\| \|x - y\|}{\|x - y\|} \\ &= \frac{\left| \|p(x)\| - \|p(y)\| \right|}{\|x - y\|} + \|p(y)\|. \end{aligned}$$

The radial function of K , $r : \mathbb{S}^{d-1} \rightarrow [0, \infty)$ with $r(x) = \|p(x)\|$, is Lipschitz (see, e.g., [T, Theorem 1]), and $\|p(y)\|$ is bounded, so we can see that the projection p is

indeed Lipschitz (in fact, one can show it is bi-Lipschitz), and so $\mathcal{H}_{d-1}(\partial K) < \infty$ (see, e.g., [Ma, Theorem 7.5]). \square

If C is a spherical cap of radius $\vartheta \in (0, \frac{\pi}{2})$ centered at x , and T is the hyperplane tangent to the sphere at x , then the central projection from C to T is injective, preserves geodesics (and therefore convexity) and is continuously differentiable, with a continuously differentiable inverse on its image (and therefore is bi-Lipschitz). We then have Corollary 7.2:

Corollary 7.2. *If $K \subset \mathbb{S}^{d-1}$ is geodesically convex and contained by a spherical cap of radius $\vartheta \in (0, \frac{\pi}{2})$, then $\mathcal{H}_{d-2}(\partial K)$ is finite.*

ACKNOWLEDGMENTS

We would like to thank David de Laat and Alexandr Polyanskii for useful discussions. All of the authors express gratitude to ICERM for hospitality and support during the Collaborate@ICERM program in 2021.

REFERENCES

- [A] J. Aczél, *Solution to problem 35, I* (Hungarian). Mat. Lapok **3** (1952), 94–95.
- [AlSz] Noga Alon and Mario Szegedy, *Large sets of nearly orthogonal vectors*, Graphs Combin. **15** (1999), no. 1, 1–4. MR1684496
- [BE] C. Bachoc and M. Ehler, *Tight p -fusion frames*, Appl. Comput. Harmon. Anal. **35** (2013), no. 1, 1–15, DOI 10.1016/j.acha.2012.07.001. MR3053743
- [BV] Christine Bachoc and Frank Vallentin, *New upper bounds for kissing numbers from semidefinite programming*, J. Amer. Math. Soc. **21** (2008), no. 3, 909–924, DOI 10.1090/S0894-0347-07-00589-9. MR2393433
- [Ball1] Keith Ball, *Volume ratios and a reverse isoperimetric inequality*, J. London Math. Soc. (2) **44** (1991), no. 2, 351–359, DOI 10.1112/jlms/s2-44.2.351. MR1136445
- [BGOY] Alexander Barg, Alexey Glazyrin, Kasso A. Okoudjou, and Wei-Hsuan Yu, *Finite two-distance tight frames*, Linear Algebra Appl. **475** (2015), 163–175, DOI 10.1016/j.laa.2015.02.020. MR3325226
- [BCGKO] R. Ben-Av, X. Chen, A. Goldberger, S. Kang, and K. A. Okoudjou, *Phase transitions for the minimizers of the p^{th} frame potentials in \mathbb{R}^2* , Preprint, [arXiv:2212.04444](https://arxiv.org/abs/2212.04444), 2022.
- [BFGMPV] Dmitriy Bilyk, Damir Ferizović, Alexey Glazyrin, Ryan W. Matzke, Josiah Park, and Oleksandr Vlsiuk, *Potential theory with multivariate kernels*, Math. Z. **301** (2022), no. 3, 2907–2935, DOI 10.1007/s00209-022-03000-z. MR4437344
- [BFGMPV1] D. Bilyk, D. Ferizović, A. Glazyrin, R. Matzke, J. Park, and O. Vlsiuk, *Optimal measures for multivariate geometric potentials*, Indiana Univ. Math. J. (to appear 2024), <https://www.iumj.indiana.edu/IUMJ/Preprints/60289.pdf>, last accessed 7/21/2024.
- [BGMPV1] Dmitriy Bilyk, Alexey Glazyrin, Ryan Matzke, Josiah Park, and Oleksandr Vlsiuk, *Optimal measures for p -frame energies on spheres*, Rev. Mat. Iberoam. **38** (2022), no. 4, 1129–1160, DOI 10.4171/rmi/1329. MR4445910
- [BGMPV2] Dmitriy Bilyk, Alexey Glazyrin, Ryan Matzke, Josiah Park, and Oleksandr Vlsiuk, *Energy on spheres and discreteness of minimizing measures*, J. Funct. Anal. **280** (2021), no. 11, Paper No. 108995, 28, DOI 10.1016/j.jfa.2021.108995. MR4233399
- [BF] John J. Benedetto and Matthew Fickus, *Finite normalized tight frames*, Adv. Comput. Math. **18** (2003), no. 2-4, 357–385, DOI 10.1023/A:1021323312367. MR1968126
- [BHS] Sergiy V. Borodachov, Douglas P. Hardin, and Edward B. Saff, *Discrete energy on rectifiable sets*, Springer Monographs in Mathematics, Springer, New York, [2019] ©2019, DOI 10.1007/978-0-387-84808-2. MR3970999

- [CGGKO] X. Chen, V. Gonzalez, E. Goodman, S. Kang, and K. A. Okoudjou, *Universal optimal configurations for the p -frame potentials*, Adv. Comput. Math. **46** (2020), no. 1, Paper No. 4, 22, DOI 10.1007/s10444-020-09745-7. MR4057629
- [CK] Henry Cohn and Abhinav Kumar, *Universally optimal distribution of points on spheres*, J. Amer. Math. Soc. **20** (2007), no. 1, 99–148, DOI 10.1090/S0894-0347-06-00546-7. MR2257398
- [DH] H. Davenport and Gy. Hajós, *Problem 35* (Hungarian), Mat. Lapok **2** (1951), 68.
- [Dea] Louis Deaett, *The minimum semidefinite rank of a triangle-free graph*, Linear Algebra Appl. **434** (2011), no. 8, 1945–1955, DOI 10.1016/j.laa.2010.11.052. MR2775783
- [D] B. V. Dekster, *The Jung theorem for spherical and hyperbolic spaces*, Acta Math. Hungar. **67** (1995), no. 4, 315–331, DOI 10.1007/BF01874495. MR1315815
- [Del] P. Delsarte, *An algebraic approach to the association schemes of coding theory*, Philips Res. Rep. Suppl. **10** (1973), vi+97. MR384310
- [EO] M. Ehler and K. A. Okoudjou, *Minimization of the probabilistic p -frame potential*, J. Statist. Plann. Inference **142** (2012), no. 3, 645–659, DOI 10.1016/j.jspi.2011.09.001. MR2853573
- [GI] A. Glazyrin, *Moments of isotropic measures and optimal projective codes*, Preprint, arXiv:1904.11159, 2019.
- [GP] Alexey Glazyrin and Josiah Park, *Repeated minimizers of p -frame energies*, SIAM J. Discrete Math. **34** (2020), no. 4, 2411–2423, DOI 10.1137/19M1282702. MR4175830
- [H1] Masatake Hirao, *On p -frame potentials of determinantal point processes on the sphere*, JSIAM Lett. **13** (2021), 21–24, DOI 10.14495/jsiaml.13.21. MR4252568
- [H2] Masatake Hirao, *On p -frame potentials of the Beltrán and Etayo point processes on the sphere*, JSIAM Lett. **15** (2023), 1–4. MR4570003
- [Jung] Heinrich Jung, *Ueber die kleinste Kugel, die eine räumliche Figur einschliesst* (German), J. Reine Angew. Math. **123** (1901), 241–257, DOI 10.1515/crll.1901.123.241. MR1580570
- [KL] G. A. Kabatjanskiĭ and V. I. Levenšteĭn, *Bounds for packings on the sphere and in space* (Russian), Problemy Peredači Informacii **14** (1978), no. 1, 3–25. MR514023
- [Lev] V. I. Levenšteĭn, *Boundaries for packings in n -dimensional Euclidean space* (Russian), Dokl. Akad. Nauk SSSR **245** (1979), no. 6, 1299–1303. MR529659
- [Ma] Pertti Mattila, *Geometry of sets and measures in Euclidean spaces*, Cambridge Studies in Advanced Mathematics, vol. 44, Cambridge University Press, Cambridge, 1995. Fractals and rectifiability, DOI 10.1017/CBO9780511623813. MR1333890
- [Mus] Oleg R. Musin, *The kissing number in four dimensions*, Ann. of Math. (2) **168** (2008), no. 1, 1–32, DOI 10.4007/annals.2008.168.1. MR2415397
- [OS] A. M. Odlyzko and N. J. A. Sloane, *New bounds on the number of unit spheres that can touch a unit sphere in n dimensions*, J. Combin. Theory Ser. A **26** (1979), no. 2, 210–214, DOI 10.1016/0097-3165(79)90074-8. MR530296
- [Pol] A. Polyanskii, *On almost-equidistant sets*, Linear Algebra Appl. **563** (2019), 220–230, DOI 10.1016/j.laa.2018.11.005. MR3873956
- [Pud] Pavel Pudlák, *Cycles of nonzero elements in low rank matrices*, Combinatorica **22** (2002), no. 2, 321–334, DOI 10.1007/s004930200015. Special issue: Paul Erdős and his mathematics. MR1909089
- [Ran] R. A. Rankin, *The closest packing of spherical caps in n dimensions*, Proc. Glasgow Math. Assoc. **2** (1955), 139–144. MR74013
- [Ros] Moshe Rosenfeld, *Almost orthogonal lines in E^d* , Applied geometry and discrete mathematics, DIMACS Ser. Discrete Math. Theoret. Comput. Sci., vol. 4, Amer. Math. Soc., Providence, RI, 1991, pp. 489–492, DOI 10.1090/dimacs/004/38. MR1116372
- [SvdW] K. Schütte and B. L. van der Waerden, *Das Problem der dreizehn Kugeln* (German), Math. Ann. **125** (1953), 325–334, DOI 10.1007/BF01343127. MR53537
- [Sid] V. M. Sidel’nikov, *New estimates for the closest packing of spheres in n -dimensional Euclidean space*, Mat. Sb. **24** (1974), 148–158.
- [Sz] T. Szele, *Solution to problem 35, II* (Hungarian), Mat. Lapok **3** (1952), 95.
- [T] F. A. Toranzos, *Radial functions of convex and star-shaped bodies*, Amer. Math. Monthly **74** (1967), 278–280, DOI 10.2307/2316022. MR208469

- [W] L. Welch, *Lower bounds on the maximum cross correlation of signals*, IEEE Trans. Inform. Theory **20** (2006), 397–399.
- [WO] Clare Wickman and Kasso A. Okoudjou, *Gradient flows for probabilistic frame potentials in the Wasserstein space*, SIAM J. Math. Anal. **55** (2023), no. 3, 2324–2346, DOI 10.1137/21M1425633. MR4603681
- [XX] Zhiqiang Xu and Zili Xu, *The minimizers of the p -frame potential*, Appl. Comput. Harmon. Anal. **52** (2021), 366–379, DOI 10.1016/j.acha.2020.04.003. MR4218427
- [Y] V. A. Yudin, *Minimum potential energy of a point system of charges* (Russian, with Russian summary), Diskret. Mat. **4** (1992), no. 2, 115–121, DOI 10.1515/dma.1993.3.1.75; English transl., Discrete Math. Appl. **3** (1993), no. 1, 75–81. MR1181534

SCHOOL OF MATHEMATICS, UNIVERSITY OF MINNESOTA, MINNEAPOLIS, MINNESOTA 55455
Email address: `dbilyk@math.umn.edu`

DEPARTMENT OF MATHEMATICS, KATHOLIEKE UNIVERSITEIT LEUVEN, LEUVEN, BELGIUM
Email address: `damir.ferizovic@kuleuven.be`

SCHOOL OF MATHEMATICAL & STATISTICAL SCIENCES, THE UNIVERSITY OF TEXAS RIO GRANDE VALLEY, BROWNSVILLE, TEXAS 78520
Email address: `alexey.glazyrin@utrgv.edu`

DEPARTMENT OF MATHEMATICS, VANDERBILT UNIVERSITY, NASHVILLE, TENNESSEE 37240
Email address: `ryan.w.matzke@vanderbilt.edu`

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GEORGIA 30332
Email address: `j.park@gatech.edu`

DEPARTMENT OF MATHEMATICS, VANDERBILT UNIVERSITY, NASHVILLE, TENNESSEE 37240
Email address: `oleksandr.vlasiuk@gmail.com`