

# CLASSIFYING CONTACT STRUCTURES ON THE SPHERE

## INTRODUCTION

Contact geometry is the study of odd-dimensional smooth manifolds equipped with contact structures, i.e. hyperplane distributions  $\xi = \ker \alpha$  satisfying the contact condition

$$\alpha \wedge (d\alpha)^n \neq 0.$$

While they originally arise in the study of ODEs and in classical mechanics, the topological study of contact manifolds is a more recent and very active field of research.

A manifold can have multiple different contact structures, which can be either rigid (in which case one speaks of a "tight" manifold) or flexible (in the sense that they satisfy an h-principle). The latter contact manifolds are then called overtwisted. A foundational result of Eliashberg [Eli89] and Borman–Eliashberg–Murphy [BEM15], roughly speaking, states that overtwisted contact manifolds exist in abundance, namely whenever the manifold admits the topological version of a contact structure (an *almost* contact structure), which is a first obvious obstruction. In dimension three, an almost contact structure is simply an oriented 2-plane field.

To illustrate this dichotomy, consider the sphere  $S^3$ . By a result of Eliashberg, it has precisely one tight contact structure. On the other hand, it has infinitely many overtwisted contact structures, corresponding to the infinitely many homotopy classes of 2-plane fields on the 3-sphere. There are other examples where there are infinitely many or no tight contact structures on a contact manifold.

A further interesting property of contact manifolds comes from the fact that contact geometry is the odd-dimensional counterpart to symplectic geometry. Often, it is possible to view a contact manifold as the boundary of a symplectic manifold. Manifolds that are in this sense "fillable" are always tight [Gro85, Eli91]. The contrary, however, doesn't need to hold and one can ask the question under which conditions such tight, but non-fillable manifolds exist. The first examples of tight and non-fillable contact manifolds were constructed by Etnyre–Honda [EH02] in dimension three, and by Massot–Niederkrueger–Wendl [MNW13] in higher dimensions.

In the following, I will start with a brief overview of the results and methods of my Master's thesis (for more details see the summary of the thesis). Then, I give a brief overview of the current research status in the area and finally state and explain some open questions that I plan to work on in the dissertation.

## TIGHT NON-FILLABLE CONTACT STRUCTURES ON THE SPHERE

Recently, Bowden–Gironella–Moreno–Zhou [BGMZ22] have proved the existence of homotopically standard, non-fillable but tight contact structures on all spheres  $S^{2n+1}$  with  $n \geq 2$ . One fundamental ingredient of the work is the Giroux correspondence [Gir02]. There is a one to one correspondence between oriented contact

structures up to isotopy and open book decompositions up to positive stabilization. Starting with a specific open book decomposition of  $S^{2n-1}$ , one can, as an instance of the Giroux correspondence, construct a contact form on this manifold using a well-known construction by Thurston–Winkelnkemper. Then, according to Bourgeois, this contact structure can be extended to a contact structure on  $S^{2n-1} \times T^2$  that is  $T^2$ -invariant. More recent results by [BGM22] and [AZ24] show that this contact structure is tight. Applying subcritical surgery (preserving the tightness), one can kill the topology of the  $T^2$ -factor and obtain a tight contact structure on  $S^{2n+1}$ . The symplectic fillability fails for homological reasons as can be shown via holomorphic curves.

### FILLABILITY HIERARCHY

As explained above, the notion of fillability is crucial in understanding tight contact structures. However, there are actually a lot of different notions of fillability in symplectic geometry.

**Definition 1** (weak and strong filling). *Let  $(M, \xi)$  be a contact manifold and  $(W, \omega)$  a symplectic manifold s.t.  $\partial W = M$  (as oriented manifolds).*

- $(W, \omega)$  is a weak filling of  $(M, \xi)$  iff  $\omega|_\xi > 0$
- $(W, \omega)$  is a strong filling of  $(M, \xi)$  iff  $\omega = d\alpha$  near the boundary.

**Definition 2** (Liouville filling). *A Liouville cobordism is a compact oriented symplectic manifold where the globally defined Liouville vector field  $X$  is negatively transverse along  $\partial W_-$  (concave side) and positively transverse along  $\partial W_+$  (convex side). If  $\partial W_- = \emptyset$ , we call it a Liouville domain. A Liouville filling of a contact manifold  $(M, \xi = \ker \alpha)$  is a Liouville domain  $(W, \lambda)$  such that  $(\partial W, \ker \lambda|_{\partial W})$  is contactomorphic to  $(M, \ker \alpha)$ .*

**Definition 3** (Weinstein filling). *A Weinstein filling is a Liouville filling such that the Liouville vector field  $X$  is gradient-like for a Morse function which is locally constant on the boundary.*

In general, we have the following sequence of strict inclusions (we abbreviate e.g. "Weinstein" for "Weinstein fillable contact manifold")

$$\{\text{Weinstein}\} \stackrel{1}{\subsetneq} \{\text{Liouville}\} \stackrel{2}{\subsetneq} \{\text{strong}\} \stackrel{3}{\subsetneq} \{\text{weak}\} \stackrel{4}{\subsetneq} \{\text{tight}\}$$

In dimension  $\geq 5$ , 1 was proved by [BCS14, Theorem 1.5], 2 by [Zho21], 3 by [BGM22] and 4 by [MNW13] (in dimension 3, they have been known for a while). All of the above examples where such an inclusion is strict rely on special geometric constructions. Now, due to [BGMZ22], the last inclusion is known to be strict for all contact manifolds (of course only if there are tight contact structures at all). In the same spirit, they prove that for any  $n \geq 3$ , there exist homotopically standard Liouville fillable contact structures on  $S^{2n+1}$  that are not Weinstein fillable.

### OPEN QUESTIONS

**Question 1.** *For being able to further classify contact structures on the sphere, it is certainly interesting to better understand the underlying almost contact structures. Using the fact that  $S^{2n+1}$  is stably parallelizable, we can show that an almost*

contact structure on  $S^{2n+1}$  corresponds to a map that associates an almost complex structure on  $R^{2n+2}$  to every point of the sphere. Almost complex structures on  $R^{2n+2}$  are given by  $\mathrm{SO}(2n+2)/\mathrm{U}(n+1)$ , so we have the following correspondence

$$\mathrm{Acont}(S^{2n+1}) = \pi_{2n+1}(\mathrm{SO}(2n+2)/\mathrm{U}(n+1)).$$

This fundamental group has been computed by Harris [Har63] and we obtain

$$\mathrm{Acont}(S^{2n+1}) = \pi_{2n+1}(\mathrm{SO}(2n+2)/\mathrm{U}(n+1)) = \begin{cases} \mathbb{Z}_{n!} & n \equiv 0 \pmod{4} \\ \mathbb{Z} & n \equiv 1 \pmod{4} \\ \mathbb{Z}_{n!/2} & n \equiv 2 \pmod{4} \\ \mathbb{Z} \oplus \mathbb{Z}_2 & n \equiv 3 \pmod{4} \end{cases}.$$

It is possible to choose the first identification in such a way that  $\xi_{\mathrm{std}}$  corresponds to 0. Taking the negative stabilization of that open book, we obtain a contact structure  $\xi_{\mathrm{neg}}$ . Which element does  $\xi_{\mathrm{neg}}$  correspond to?

**Question 2.** Is there a Liouville but not Weinstein fillable contact structure on  $S^5$ ?

**Question 3.** Is there a strong but not Liouville fillable structure on  $S^{2n+1}$ ,  $n \geq 2$ ?

## REFERENCES

- [AZ24] Russell Avdek and Zhengyi Zhou. Bourgeois' contact manifolds are tight. Technical report, April 2024. arXiv:2404.16311 [math] type: article.
- [BCS14] Jonathan Bowden, Diarmuid Crowley, and András I. Stipsicz. The topology of Stein fillable manifolds in high dimensions I. *Proceedings of the London Mathematical Society*, 109(6):1363–1401, December 2014.
- [BEM15] Matthew Strom Borman, Yakov Eliashberg, and Emmy Murphy. Existence and classification of overtwisted contact structures in all dimensions. *Acta Mathematica*, 215(2):281–361, January 2015.
- [BGM22] Jonathan Bowden, Fabio Gironella, and Agustin Moreno. Bourgeois contact structures: Tightness, fillability and applications. *Inventiones mathematicae*, 230(2):713–765, November 2022.
- [BGMZ22] Jonathan Bowden, Fabio Gironella, Agustin Moreno, and Zhengyi Zhou. Tight contact structures without symplectic fillings are everywhere. 2022.
- [BO11] Mohan Bhupal and Burak Ozbagci. Milnor open books of links of some rational surface singularities. *Pacific Journal of Mathematics*, 254(1):47–65, December 2011.
- [Bou02] Frédéric Bourgeois. Odd dimensional tori are contact manifolds. *International Mathematics Research Notices*, 2002(30):1571–1574, January 2002.
- [CMP19] Roger Casals, Emmy Murphy, and Francisco Presas. Geometric criteria for overtwistedness. *Journal of the American Mathematical Society*, 32(2):563–604, April 2019.
- [EH02] John B. Etnyre and Ko Honda. Tight contact structures with no symplectic fillings. *Inventiones mathematicae*, 148(3):609–626, June 2002.
- [Eli89] Y. Eliashberg. Classification of overtwisted contact structures on 3-manifolds. *Inventiones Mathematicae*, 98(3):623–637, October 1989.
- [Eli91] Yakov Eliashberg. Filling by holomorphic discs and its applications. volume 2 of *London Mathematical Society Lecture Note Series*, pages 45–68. Cambridge University Press, Cambridge, 1991.
- [Gei08] Hansjörg Geiges. *An Introduction to Contact Topology*. Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, March 2008.
- [Gir02] Emmanuel Giroux. Geometrie de contact: de la dimension trois vers les dimensions superieures. In Ta Tsien et al. Li, editor, *Proceedings of the international congress of mathematicians, ICM 2002*, 3 volumes. ISBN 7-04-008690-5, pages 405–414, Beijing, China, 2002. Beijing: Higher Education Press.
- [Gro85] M. Gromov. Pseudo holomorphic curves in symplectic manifolds. *Inventiones mathematicae*, 82(2):307–347, June 1985.
- [Har63] Bruno Harris. Some Calculations of Homotopy Groups of Symmetric Spaces. *Transactions of the American Mathematical Society*, 106(1):174–184, 1963.
- [HH23] Ko Honda and Yang Huang. Convex hypersurface theory in contact topology. Technical report, June 2023. arXiv:1907.06025 [math] type: article.
- [Hua17] Yang Huang. On plastikstufe, bordered Legendrian open book and overtwisted contact structures. *Journal of Topology*, 10(3):720–743, June 2017.
- [MNW13] Patrick Massot, Klaus Niederkrüger, and Chris Wendl. Weak and strong fillability of higher dimensional contact manifolds. *Inventiones mathematicae*, 192(2):287–373, May 2013.
- [Nie06] Klaus Niederkrüger. The plastikstufe – a generalization of the overtwisted disk to higher dimensions. *Algebraic & Geometric Topology*, 6(5):2473–2508, December 2006.
- [Zho21] Zhengyi Zhou.  $(\mathbb{RP}^{2n-1}, \xi_{\text{std}})$  is not exactly fillable for  $n \neq 2^k$ . *Geometry & Topology*, 25(6):3013–3052, November 2021. arXiv:2001.09718 [math].