

# Homotopically Standard Tight Non-fillable Contact Structures on the Sphere

Josua Kugler  
results by Bowden, Gironella, Moreno and Zhou

Heidelberg University

# Background

# Contact topology

## Contact manifold:

- odd-dimensional smooth manifold
- codim-1 hyperplane distribution  $\ker \alpha$
- contact condition:

$$\alpha \wedge (d\alpha)^n \neq 0$$

# Contact topology

**Contact topology:** The study of contact manifolds, up to isotopy.

# Contact topology

**Contact topology:** The study of contact manifolds, up to isotopy.

**Fillability:** *fillable* contact mflds are boundaries of symplectic mflds.

# Contact topology

**Contact topology:** The study of contact manifolds, up to isotopy.

**Fillability:** *fillable* contact mflds are boundaries of symplectic mflds.

## Fillability question

Which contact manifolds are **fillable**?

# Contact topology

**Contact topology:** The study of contact manifolds, up to isotopy.

**Fillability:** *fillable* contact mflds are boundaries of symplectic mflds.

## Fillability question

Which contact manifolds are **fillable**?

Eliashberg, Borman–Eliashberg–Murphy:

**Dichotomy:** Rigidity vs. Flexibility.

- **tight** (*rigid/geometric*);
- **overtwisted** (*flexible/topological*).

# Contact topology

**Contact topology:** The study of contact manifolds, up to isotopy.

**Fillability:** *fillable* contact mflds are boundaries of symplectic mflds.

## Fillability question

Which contact manifolds are **fillable**?

Eliashberg, Borman–Eliashberg–Murphy:

**Dichotomy:** Rigidity vs. Flexibility.

- **tight** (*rigid/geometric*);
- **overtwisted** (*flexible/topological*).

## Theorem (Eliashberg–Gromov)

*Fillable contact manifolds are tight.*

Converse is false (Etnyre–Honda, Massot–Niederkrueger–Wendl).



# Existence and classification

*Topological* obstruction: *almost* contact structure, i.e. reduction of structure group to  $U(n) \times \mathbb{1}$ .

Theorem (Lutz–Martinet (dim 3), Casals–Pancholi–Presas (dim 5), Borman–Eliashberg–Murphy (any dim))

*Almost contact manifolds are contact, where the contact structure is overtwisted.*

# Existence and classification

*Topological obstruction: almost contact structure, i.e. reduction of structure group to  $U(n) \times \mathbb{1}$ .*

Theorem (Lutz–Martinet (dim 3), Casals–Pancholi–Presas (dim 5), Borman–Eliashberg–Murphy (any dim))

*Almost contact manifolds are contact, where the contact structure is overtwisted.*

## Tight manifolds

How can we understand **tight** contact manifolds?

# Contact topology: fillability

## Hierarchy of fillability:

$$\begin{array}{ccccccc} \{Stein\} & \overset{\textcircled{1}}{=} & \{Weinstein\} & \overset{\textcircled{2}}{\subsetneq} & \{Liouville\} & \overset{\textcircled{3}}{\subsetneq} & \{strong\} \\ & & & & & & \\ & & \overset{\textcircled{4}}{\subsetneq} & \{weak\} & \overset{\textcircled{5}}{\subsetneq} & \{tight\} & \end{array}$$

- $dim = 3$ :  $\textcircled{1}$  Cieliebak–Eliashberg,  $\textcircled{2}$  Bowden,  $\textcircled{3}$  Ghiggini,  $\textcircled{4}$  Eliashberg,  $\textcircled{5}$  Etnyre–Honda.
- $dim \geq 5$ :  $\textcircled{1}$  Cieliebak–Eliashberg,  
 $\textcircled{2}$  Bowden–Crowley–Stipsicz,  $\textcircled{3}$  Zhou,  
 $\textcircled{4}$  Bowden–Gironella–Moreno,  $\textcircled{5}$  Massot–Niederkrüger–Wendl.

# Contact structures on spheres

**First step in classification:** contact structures on spheres.

## Standard contact structure

The standard contact structure is  $(S^{2n-1}, \xi) = \partial(B^{2n}, \omega_{std})$ .

# Contact structures on spheres

**First step in classification:** contact structures on spheres.

## Standard contact structure

The standard contact structure is  $(S^{2n-1}, \xi) = \partial(B^{2n}, \omega_{std})$ .

## Theorem (Eliashberg, '91)

*On  $S^3$ , it is the unique tight contact structure.*

In particular, no tight and non-fillable contact structures on  $S^3$ .

# Exotic spheres

Theorem (Bowden–Gironella–Moreno–Zhou '22-'24)

*For every  $n \geq 2$ , the sphere  $\mathbb{S}^{2n+1}$  admits a tight, non-fillable contact structure that is homotopically standard.*

# Tight and non-fillable structures in $\dim \geq 5$

$(M, \xi)$  tight contact manifold.

# Tight and non-fillable structures in $\dim \geq 5$

$(M, \xi)$  tight contact manifold.

Then connected sum with an “exotic” sphere yields:

**Theorem (Bowden–Gironella–Moreno–Zhou '22-'24)**

*In  $\dim \geq 7$ ,  $M$  admits a tight and non strongly-fillable contact structure  $\xi_{\text{exotic}}$ , in the same almost contact class as  $\xi$ .*



# Tight and non-fillable structures in $\dim \geq 5$

$(M, \xi)$  tight contact manifold.

Then connected sum with an “exotic” sphere yields:

## Theorem (Bowden–Gironella–Moreno–Zhou '22-'24)

*In  $\dim \geq 7$ ,  $M$  admits a tight and non strongly-fillable contact structure  $\xi_{\text{exotic}}$ , in the same almost contact class as  $\xi$ .*

*In  $\dim = 5$ , the same holds, if the first Chern class vanishes.*

# General remarks


- This is a novel and strictly higher-dimensional phenomenon (false in dim 3).

# General remarks

- This is a novel and strictly higher-dimensional phenomenon (false in dim 3).
- Suggests that higher-dimensional contact phenomena should occur independently of underlying smooth topology.

# **Tight and non-fillable spheres**

# Open books



# Giroux correspondence

**Giroux:** Contact structures are *supported* by open books.




Figure: Supported contact structure.

# Bourgeois contact structures

## Theorem (Bourgeois '02)

*Open book supporting  $(M, \xi) \rightsquigarrow$  contact structure on  $M \times \mathbb{T}^2$ .*

The resulting contact structure is  $\mathbb{T}^2$ -equivariant.

# Geometric construction

We now construct **one** tight and non-fillable contact structure on  $\mathbb{S}^{2n+1}$ .



# Geometric construction

We now construct **one** tight and non-fillable contact structure on  $\mathbb{S}^{2n+1}$ .

- Milnor open book on  $\mathbb{S}^{2n-1}$

# Geometric construction

We now construct **one** tight and non-fillable contact structure on  $\mathbb{S}^{2n+1}$ .

- Milnor open book on  $\mathbb{S}^{2n-1}$
- $\rightsquigarrow$  Bourgeois manifold  $\mathbb{S}^{2n-1} \times \mathbb{T}^2$

# Geometric construction

We now construct **one** tight and non-fillable contact structure on  $\mathbb{S}^{2n+1}$ .

- Milnor open book on  $\mathbb{S}^{2n-1}$
- $\rightsquigarrow$  Bourgeois manifold  $\mathbb{S}^{2n-1} \times \mathbb{T}^2$
- $\rightsquigarrow$  two 1-surgeries =  $\mathbb{S}^{2n-1} \times \mathbb{S}^2 \rightsquigarrow$  one 2-surgery =  $\mathbb{S}^{2n+1}$ .

# Geometric construction

We now construct **one** tight and non-fillable contact structure on  $\mathbb{S}^{2n+1}$ .

- Milnor open book on  $\mathbb{S}^{2n-1}$
- $\rightsquigarrow$  Bourgeois manifold  $\mathbb{S}^{2n-1} \times \mathbb{T}^2$
- $\rightsquigarrow$  two 1-surgeries =  $\mathbb{S}^{2n-1} \times \mathbb{S}^2 \rightsquigarrow$  one 2-surgery =  $\mathbb{S}^{2n+1}$ .
- If  $n \geq 3$ , surgeries are *subcritical*  $\rightsquigarrow$  by 'Eliashberg's' h-ppl, Weinstein cobordism  $\rightsquigarrow$  contact manifold  $(\mathbb{S}^{2n+1}, \xi_{ex})$ .

# Geometric construction

We now construct **one** tight and non-fillable contact structure on  $\mathbb{S}^{2n+1}$ .

- Milnor open book on  $\mathbb{S}^{2n-1}$
- $\rightsquigarrow$  Bourgeois manifold  $\mathbb{S}^{2n-1} \times \mathbb{T}^2$
- $\rightsquigarrow$  two 1-surgeries =  $\mathbb{S}^{2n-1} \times \mathbb{S}^2 \rightsquigarrow$  one 2-surgery =  $\mathbb{S}^{2n+1}$ .
- If  $n \geq 3$ , surgeries are *subcritical*  $\rightsquigarrow$  by 'Eliashberg's' h-pple, Weinstein cobordism  $\rightsquigarrow$  contact manifold  $(\mathbb{S}^{2n+1}, \xi_{ex})$ .

**Claim:**  $(\mathbb{S}^{2n+1}, \xi_{ex})$  is homotopically standard, tight and non-fillable.

# Tightness

## Facts:

- ① Milnor open book  $\Rightarrow$  algebraically tight Bourgeois manifold.

# Tightness

## Facts:

- ① Milnor open book  $\Rightarrow$  algebraically tight Bourgeois manifold.
  - Algebraic tightness is non-vanishing of a certain contact homology algebra.

# Tightness

## Facts:

- ① Milnor open book  $\Rightarrow$  algebraically tight Bourgeois manifold.
  - Algebraic tightness is non-vanishing of a certain contact homology algebra.
- ② Algebraic tightness is preserved under surgeries.



# Tightness

## Facts:

- ① Milnor open book  $\Rightarrow$  algebraically tight Bourgeois manifold.
  - Algebraic tightness is non-vanishing of a certain contact homology algebra.
- ② Algebraic tightness is preserved under surgeries.
- ③ Algebraically tight  $\implies$  tight.

# Tightness

## Facts:

- 1 Milnor open book  $\Rightarrow$  algebraically tight Bourgeois manifold.
  - Algebraic tightness is non-vanishing of a certain contact homology algebra.
- 2 Algebraic tightness is preserved under surgeries.
- 3 Algebraically tight  $\implies$  tight.

Milnor open book  $\Rightarrow (\mathbb{S}^{2n+1}, \xi_{ex})$  is *tight*.

# Non-fillability

**Non-fillability** of  $(\mathbb{S}^{2n+1}, \xi_{ex})$  can be proven via:

- 1 Homological obstruction and cobordisms as in [Bowden–Gironella–Moreno], building on [Massot–Niederkrüger–Wendl].
- 2 Symplectic cohomology computations as in [Zhou].

# Convex Decomposition and Capping Cobordism

## Observations:

- Bourgeois manifolds have convex decomposition

$$M \times \mathbb{T}^2 = (M \times \mathbb{S}^1) \times \mathbb{S}^1 = V_+ \times \mathbb{S}^1 \cup_{\phi} \overline{V}_- \times \mathbb{S}^1,$$

with  $V_{\pm} = \Sigma \times D^*\mathbb{S}^1$ ,  $\Sigma$  = page of the open book,  $\phi$  = monodromy.

# Convex Decomposition and Capping Cobordism

## Observations:


- Bourgeois manifolds have convex decomposition

$$M \times \mathbb{T}^2 = (M \times \mathbb{S}^1) \times \mathbb{S}^1 = V_+ \times \mathbb{S}^1 \cup_{\phi} \overline{V}_- \times \mathbb{S}^1,$$

with  $V_{\pm} = \Sigma \times D^*\mathbb{S}^1$ ,  $\Sigma$  = page of the open book,  $\phi$  = monodromy.

- Capping cobordism from  $M \times T^2$  to  $\Gamma \times \mathbb{S}^2$ , via handles  $V_{\pm} \times D^2$  with co-core  $C_{\pm} \simeq V_{\pm}$ .

# Capping Cobordism: Example



# Homological obstructions

## Theorem (Bowden–Gironella–Moreno)

$M \times T^2 = V \times \mathbb{S}^1 = V_+ \times \mathbb{S}^1 \cup_{\phi} \overline{V_-} \times \mathbb{S}^1$  with convex decomposition,  $\Gamma = \partial V_{\pm}$  dividing set. If  $W$  is a symplectic filling of  $M \times T^2$ , then

$$H_*(\Gamma) \rightarrow H_*(V_{\pm}) \rightarrow H_*(W),$$

induced by inclusion. Then second map is injective on image of the first.

Namely, if a homology class in  $\Gamma$  survives in  $V_{\pm}$ , then it survives in the filling.

# Idea of proof


- Capping cobordism from  $M \times T^2$  to  $\Gamma \times \mathbb{S}^2$ , via handles  $V_{\pm} \times D^2$  with co-core  $C_{\pm} \simeq V_{\pm}$ .



# Idea of proof

- Capping cobordism from  $M \times T^2$  to  $\Gamma \times \mathbb{S}^2$ , via handles  $V_{\pm} \times D^2$  with co-core  $C_{\pm} \simeq V_{\pm}$ .
- Second factor gives moduli space of spheres  $\mathcal{M}_*$  with evaluation map  $ev : \mathcal{M}_* \rightarrow W$ .


# Idea of proof



# Idea of proof

- Capping cobordism from  $M \times T^2$  to  $\Gamma \times \mathbb{S}^2$ , via handles  $V_{\pm} \times D^2$  with co-core  $C_{\pm} \simeq V_{\pm}$ .
- Second factor gives moduli space of spheres  $\mathcal{M}_*$  with evaluation map  $ev : \mathcal{M}_* \rightarrow W$ .
- Spheres intersect  $C_{\pm}$  precisely once  $\rightsquigarrow$  intersection map  $\mathcal{I}_{\pm} : \mathcal{M}_* \rightarrow C_{\pm}$ .

# Idea of proof



# Idea of proof

- Capping cobordism from  $M \times T^2$  to  $\Gamma \times \mathbb{S}^2$ , via handles  $V_{\pm} \times D^2$  with co-core  $C_{\pm} \simeq V_{\pm}$ .
- Second factor gives moduli space of spheres  $\mathcal{M}_*$  with evaluation map  $ev : \mathcal{M}_* \rightarrow W$ .
- Spheres intersect  $C_{\pm}$  precisely once  $\rightsquigarrow$  intersection map  $\mathcal{I}_{\pm} : \mathcal{M}_* \rightarrow C_{\pm}$ .
- If  $\sigma \subset W$  satisfies  $\partial\sigma = z$  with  $z$  cycle in  $\Gamma$ , then  $z$  also bounds  $b = \mathcal{I}_{\pm} ev^{-1}(\sigma) \subset V_{\pm}$ . □

# Homological obstructions

## Fact:

- ① If  $\dim \geq 7$ , subcritical surgeries on  $\mathbb{S}^{2n-1} \times \mathbb{T}^2$  can be pushed away from dividing set to  $V_+$ .

$\Rightarrow (\mathbb{S}^{2n+1}, \xi_{ex})$  still has a dividing set  $\Gamma$ ,

with  $H_n(\Gamma) \neq 0$ .

# Homological obstructions

## Fact:

- 1 If  $\dim \geq 7$ , subcritical surgeries on  $\mathbb{S}^{2n-1} \times \mathbb{T}^2$  can be pushed away from dividing set to  $V_+$ .

$\Rightarrow (\mathbb{S}^{2n+1}, \xi_{ex})$  still has a dividing set  $\Gamma$ ,

with  $H_n(\Gamma) \neq 0$ .

- 2 Homological obstruction theorem persists under surgery away from dividing set (capping cobordisms).




Figure: Capping cobordism.

**End of the proof:** Assume  $W$  filling of  $(\mathbb{S}^{2n+1}, \xi_{ex})$ .

- $\exists$  homology class  $z \subset \Gamma$  s.t.  $z$  survives in  $V_{\pm}$ .



**End of the proof:** Assume  $W$  filling of  $(\mathbb{S}^{2n+1}, \xi_{ex})$ .

- $\exists$  homology class  $z \in \Gamma$  s.t.  $z$  survives in  $V_{\pm}$ .
- Homological obstruction theorem:  $H_n(\Gamma) \rightarrow H_n(W) \neq 0$ .

**End of the proof:** Assume  $W$  filling of  $(\mathbb{S}^{2n+1}, \xi_{ex})$ .

- $\exists$  homology class  $z \in \Gamma$  s.t.  $z$  survives in  $V_{\pm}$ .
- Homological obstruction theorem:  $H_n(\Gamma) \rightarrow H_n(W) \neq 0$ .
- However, this factors as

$$0 \neq H_n(\Gamma) \rightarrow H_n(\mathbb{S}^{2n+1}) = 0 \rightarrow H_n(W),$$

contradiction.

Thank you!