

ELEC 4700 Winter 2020-2021
Assignment 1
Monte-Carlo Modeling of Electron Transport
Submitted by: Joud Alhuwaitat (101074617)
Due date: February 07, 2021
Lecture Section: A

1. Electron Modelling

In this section of the report, we will model the electrons in the silicon as particles with effective mass using a simplistic Monte-Carlo model. The code used for this step is shown below as well as the deliverables set out in the assignment manual afterwards.

```
% Part 1: Electron Modelling
% To calculate the thermal energy, we use Maxwell's principle of
equipartition
% of energy,
%
%  $\overline{KE} = \frac{1}{2}kT = 2(\frac{1}{2}m\overline{v^2}) \rightarrow \overline{v^2} = \frac{2kT}{m}$ 

clear all
close all
m0 = 9.10938356e-31;
m = 0.26*m0;
T = 300;
k = 1.38064852e-23;
vth = sqrt(2*k*T/m)

%%
% Or 187 km/s. The mean free path,  $l$ , is simply

l = vth*0.2e-12

%%
% Thus the mean free path is about 37.4 nm.
%
% Below are some variables to control the simulations:

height = 100e-9;
length = 200e-9;
population_size = 3000;
plot_population = 10;
time_step = height/vth/100;
iterations = 1000;
% Set to 1 to watch the movies,
% or to 0 to just see the final plots
show_movie = 0;

%%
% The arrays below will be used to hold the information about the
% state of the system, including the positions, velocities, and
% temperatures.

% Each row corresponds to an electron with the positions and velocities
% [x y vx vy]
state = zeros(population_size, 4);
```

```

trajectories = zeros(iterations, plot_population*2);
temperature = zeros(iterations,1);

%%
% Generate an initial population with constant speeds
for i = 1:population_size
    angle = rand*2*pi;
    state(i,:) = [length*rand height*rand vth*cos(angle) vth*sin(angle)];
end

%%
% Iterate over time and update the positions, while plotting the state.
for i = 1:iterations
    state(:,1:2) = state(:,1:2) + time_step.*state(:,3:4);

    % Look for collisions with the boundaries
    j = state(:,1) > length;
    state(j,1) = state(j,1) - length;

    j = state(:,1) < 0;
    state(j,1) = state(j,1) + length;

    j = state(:,2) > height;
    state(j,2) = 2*height - state(j,2);
    state(j,4) = -state(j,4);

    j = state(:,2) < 0;
    state(j,2) = -state(j,2);
    state(j,4) = -state(j,4);

    temperature(i) = (sum(state(:,3).^2) +
sum(state(:,4).^2))*m/k/2/population_size;

    % Record the trajectories
    for j=1:plot_population
        trajectories(i, (2*j):(2*j+1)) = state(j, 1:2);
    end

    % Update the movie every 5 iterations
    if show_movie && mod(i,5) == 0
        figure(1);
        subplot(2,1,1);
        hold off;
        plot(state(1:plot_population,1)./1e-9,
state(1:plot_population,2)./1e-9, 'o');
        axis([0 length/1e-9 0 height/1e-9]);
        title(sprintf('Trajectories for %d of %d Electrons with Fixed
Velocity (Part 1)',...
        plot_population, population_size));
        xlabel('x (nm)');
        ylabel('y (nm)');
        if i > 1
            subplot(2,1,2);
            hold off;

```

```

        plot(time_step*(0:i-1), temperature(1:i));
        axis([0 time_step*iterations min(temperature)*0.98
max(temperature)*1.02]);
        title('Semiconductor Temperature');
        xlabel('Time (s)');
        ylabel('Temperature (K)');
    end
    pause(0.05);
end
end


% Show trajectories after the movie is over
figure(1);
subplot(2,1,1);
title(sprintf('Electron Trajectories for %d of %d Electrons with Fixed
Velocity (Part 1)',...
    plot_population, population_size));
xlabel('x (nm)');
ylabel('y (nm)');
axis([0 length/1e-9 0 height/1e-9]);
hold on;
for i=1:plot_population
    plot(trajectories(:,i*2)./1e-9, trajectories(:,i*2+1)./1e-9, '.');
end

if(~show_movie)
    subplot(2,1,2);
    hold off;
    plot(time_step*(0:iterations-1), temperature);
    axis([0 time_step*iterations min(temperature)*0.98
max(temperature)*1.02]);
    title('Semiconductor Temperature');
    xlabel('Time (s)');
    ylabel('Temperature (K)');
end


```

- a) $v_{th} = \sqrt{2*k*T/m}$
- $v_{th} = 1.8702*10^5$
- b) The mean free path $l =$
- $l = v_{th} * 0.2*10^{-12}$
- $l = 3.7404 * 10^{-8}$

c)



d)



2. Collisions with Mean Free Path (MFP)

In this section of the report, we will assign a random velocity to each of the particles and then Model the scattering of the electrons using an exponential scattering probability: $P_{\text{scat}} = 1 - e^{-dt/\tau_{\text{mn}}}$. We will then measure the actual mean free path as well as the mean time between collisions. The code used for this step is shown below as well as the deliverables set out in the assignment manual afterwards.

```
%% Part 2: Collisions with Mean Free Path
% For the second simulation, the initial velocities are assigned based on a
% Maxwell-Boltzmann distribution, and scattering is modelled.
% First, we must calculate the scattering probability in one time step:
p_sc = 1 - exp(-time_step/0.2e-12)
```

```

%%
% The distribution of velocities in x and y is Gaussian, with a standard
% deviation of  $\sqrt{kT/m}$ . This results in an overall Maxwell-Boltzmann
% velocity distribution at temperature T
v_pdf = makedist('Normal', 'mu', 0, 'sigma', sqrt(k*T/m));

%%
% The initial population is generated using this distribution:
for i = 1:population_size
    angle = rand*2*pi;
    state(i,:) = [length*rand height*rand random(v_pdf) random(v_pdf)];
end

%%
% The average velocity should be calculated to be correct:
avg_v = sqrt(sum(state(:,3).^2)/population_size + ...
    sum(state(:,4).^2)/population_size)

%%
% This returns a velocity of about 187 km/s, which is correct.

%%
% The second simulation loop follows:
for i = 1:iterations
    %Update positions
    state(:,1:2) = state(:,1:2) + time_step.*state(:,3:4);

    j = state(:,1) > length;
    state(j,1) = state(j,1) - length;

    j = state(:,1) < 0;
    state(j,1) = state(j,1) + length;

    j = state(:,2) > height;
    state(j,2) = 2*height - state(j,2);
    state(j,4) = -state(j,4);

    j = state(:,2) < 0;
    state(j,2) = -state(j,2);
    state(j,4) = -state(j,4);

    % Scatter particles
    j = rand(population_size, 1) < p_scatter;
    state(j,3:4) = random(v_pdf, [sum(j),2]);

    % Record the temperature
    temperature(i) = (sum(state(:,3).^2) +
    sum(state(:,4).^2))*m/k/2/population_size;

    % Record positions for subset of particles that will be graphed
    for j=1:plot_population
        trajectories(i, (2*j):(2*j+1)) = state(j, 1:2);
    end
end

```

```

    % Update the movie every 5 iterations
    if show_movie && mod(i,5) == 0
        figure(2);
        subplot(3,1,1);
        hold off;
        plot(state(1:plot_population,1)./1e-9,
state(1:plot_population,2)./1e-9, 'o');
        axis([0 length/1e-9 0 height/1e-9]);
        title(sprintf('Trajectories for %d of %d Electrons (Part 2)',...
plot_population, population_size));
        xlabel('x (nm)');
        ylabel('y (nm)');
        if i > 1
            subplot(3,1,2);
            hold off;
            plot(time_step*(0:i-1), temperature(1:i));
            axis([0 time_step*iterations min(temperature)*0.98
max(temperature)*1.02]);
            title('Semiconductor Temperature');
            xlabel('Time (s)');
            ylabel('Temperature (K)');
        end

        % Show histogram of speeds
        subplot(3,1,3);
        v = sqrt(state(:,3).^2 + state(:,4).^2);
        title('Histogram of Electron Speeds');
        histogram(v);
        xlabel('Speed (m/s)');
        ylabel('Number of particles');

        pause(0.05);
    end
end

% Show trajectories after the movie is over
figure(2);
subplot(3,1,1);
title(sprintf('Trajectories for %d of %d Electrons (Part 2)',...
plot_population, population_size));
xlabel('x (nm)');
ylabel('y (nm)');
axis([0 length/1e-9 0 height/1e-9]);
hold on;
for i=1:plot_population
    plot(trajectories(:,i*2)./1e-9, trajectories(:,i*2+1)./1e-9, '.');
end

% Show temperature plot over time
if(~show_movie)
    subplot(3,1,2);
    hold off;
    plot(time_step*(0:iterations-1), temperature);
    axis([0 time_step*iterations min(temperature)*0.98
max(temperature)*1.02]);

```

```


        title('Semiconductor Temperature');
        xlabel('Time (s)');
        ylabel('Temperature (K)');
    end

% Show speed histogram
subplot(3,1,3);
v = sqrt(state(:,3).^2 + state(:,4).^2);
title('Histogram of Electron Speeds');
histogram(v);
xlabel('Speed (m/s)');
ylabel('Number of particles');


%%
% This second simulation shows that the average temperature fluctuates
% over time due to the scattering, but it has an average of 300K over time.

```


a)



b)



c)



3. Enhancements

In this section of the report, we will add an inner rectangle boundary known as a bottle neck to the simulation with the correct boundaries. We will calculate the electron density map from the final electron positions and a temperature map displayed with colors. The code used for this step is shown below as well as the deliverables set out in the assignment manual afterwards.

```
% Part 3: Enhancements
% Here, the boundaries can be set to be either specular or diffusive. If they
% are diffusive, the electrons will bounce off at a random angle rather than
% one that is symmetrical about the normal with the boundary.
%
% The non-periodic top and bottom boundaries can be set to be either
% specular (1) or diffusive (0) with the following parameters:
top_specular = 0;
bottom_specular = 0;

%%
% This simulation also includes boxes (obstacles)
% Each box can separately be set to be specular (1) or diffusive (0)

% The values are [xmin xmax ymin ymax] for each box
boxes = 1e-9.*[80 120 0 40; 80 120 60 100];
boxes_specular = [0 1];

% Generate an initial population
for i = 1:population_size
    angle = rand*2*pi;
    state(i,:) = [length*rand height*rand random(v_pdf) random(v_pdf)];

    % Make sure no particles start in a box
    while(in_box(state(i,1:2), boxes))
        state(i,1:2) = [length*rand height*rand];
    end
end
```

```

end
end

%%
% Run through the third simulation:
for i = 1:iterations
    state(:,1:2) = state(:,1:2) + time_step.*state(:,3:4);

    j = state(:,1) > length;
    state(j,1) = state(j,1) - length;

    j = state(:,1) < 0;
    state(j,1) = state(j,1) + length;

    j = state(:,2) > height;

    if(top_specular)
        state(j,2) = 2*height - state(j,2);
        state(j,4) = -state(j,4);
    else % Diffusive
        % The electron bounces off at a random angle
        state(j,2) = height;
        v = sqrt(state(j,3).^2 + state(j,4).^2);
        angle = rand([sum(j),1])*2*pi;
        state(j,3) = v.*cos(angle);
        state(j,4) = -abs(v.*sin(angle));
    end

    j = state(:,2) < 0;

    if(bottom_specular)
        state(j,2) = -state(j,2);
        state(j,4) = -state(j,4);
    else % Diffusive
        % The electron bounces off at a random angle
        state(j,2) = 0;
        v = sqrt(state(j,3).^2 + state(j,4).^2);
        angle = rand([sum(j),1])*2*pi;
        state(j,3) = v.*cos(angle);
        state(j,4) = abs(v.*sin(angle));
    end

    % Look for particles that have "entered" a box and move them to
    % where they should be.
    for j=1:population_size
        box_num = in_box(state(j,1:2), boxes);
        while(box_num ~= 0)
            % To see which side the electron collided with,
            % find which one it's closer to
            x_dist = 0;
            new_x = 0;
            if(state(j,3) > 0)
                x_dist = state(j,1) - boxes(box_num,1);
                new_x = boxes(box_num,1);
            else

```

```

        x_dist = boxes(box_num,2) - state(j,1);
        new_x = boxes(box_num,2);
    end

    y_dist = 0;
    new_y = 0;
    if(state(j,4) > 0)
        y_dist = state(j,2) - boxes(box_num, 3);
        new_y = boxes(box_num, 3);
    else
        y_dist = boxes(box_num, 4) - state(j,2);
        new_y = boxes(box_num, 4);
    end

    if(x_dist < y_dist)
        state(j,1) = new_x;
        if(~boxes_specular(box_num))
            sgn = -sign(state(j,3));
            v = sqrt(state(j,3).^2 + state(j,4).^2);
            angle = rand()*2*pi;
            state(j,3) = sgn.*abs(v.*cos(angle));
            state(j,4) = v.*sin(angle);
        else % Specular
            state(j,3) = -state(j,3);
        end
    else
        state(j,2) = new_y;
        if(~boxes_specular(box_num))
            sgn = -sign(state(j,4));
            v = sqrt(state(j,3).^2 + state(j,4).^2);
            angle = rand()*2*pi;
            state(j,3) = v.*cos(angle);
            state(j,4) = sgn.*abs(v.*sin(angle));
        else % Specular
            state(j,4) = -state(j,4);
        end
    end

    box_num = in_box(state(j,1:2), boxes);
end

% Scatter particles
j = rand(population_size, 1) < p_scatter;
state(j,3:4) = random(v_pdf, [sum(j),2]);

% Record the temperature
temperature(i) = (sum(state(:,3).^2) +
sum(state(:,4).^2))*m/k/2/population_size;

% Record positions for subset of particles that will be graphed
for j=1:plot_population
    trajectories(i, (2*j):(2*j+1)) = state(j, 1:2);
end

```

```

% Update the movie every 5 iterations
if show_movie && mod(i,5) == 0
    figure(3);
    subplot(3,1,1);
    hold off;
    plot(state(1:plot_population,1)./1e-9,
state(1:plot_population,2)./1e-9, 'o');
    hold on;

    % Plot the boxes
    for j=1:size(boxes,1)
        plot([boxes(j, 1) boxes(j, 1) boxes(j, 2) boxes(j, 2) boxes(j,
1)]./1e-9, ...
            [boxes(j, 3) boxes(j, 4) boxes(j, 4) boxes(j, 3) boxes(j,
3)]./1e-9, 'k-');
    end

    axis([0 length/1e-9 0 height/1e-9]);
    title(sprintf('Trajectories for %d of %d Electrons (Part 3)',...
plot_population, population_size));
    xlabel('x (nm)');
    ylabel('y (nm)');
    if i > 1
        subplot(3,1,2);
        hold off;
        plot(time_step*(0:i-1), temperature(1:i));
        axis([0 time_step*iterations min(temperature(1:i))*0.98
max(temperature)*1.02]);
        title('Semiconductor Temperature');
        xlabel('Time (s)');
        ylabel('Temperature (K)');
    end

    subplot(3,1,3);
    v = sqrt(state(:,3).^2 + state(:,4).^2);
    title('Histogram of Electron Speeds');
    histogram(v);
    xlabel('Speed (m/s)');
    ylabel('Number of particles');

    pause(0.05);
end
end

% Show trajectories after the movie is over
figure(3);
subplot(3,1,1);
title(sprintf('Electron Trajectories for %d of %d Electrons (Part 3)',...
plot_population, population_size));
xlabel('x (nm)');
ylabel('y (nm)');
axis([0 length/1e-9 0 height/1e-9]);
hold on;
for i=1:plot_population
    plot(trajectories(:,i*2)./1e-9, trajectories(:,i*2+1)./1e-9, '.');

```

```

end

% Plot the boxes
for j=1:size(boxes,1)
    plot([boxes(j, 1) boxes(j, 1) boxes(j, 2) boxes(j, 2) boxes(j, 1)]./1e-
9,...
        [boxes(j, 3) boxes(j, 4) boxes(j, 4) boxes(j, 3) boxes(j, 3)]./1e-9,
        'k-');
end

% Plot temperature
if(~show_movie)
    subplot(3,1,2);
    hold off;
    plot(time_step*(0:iterations-1), temperature);
    axis([0 time_step*iterations min(temperature)*0.98
max(temperature)*1.02]);
    title('Semiconductor Temperature');
    xlabel('Time (s)');
    ylabel('Temperature (K)');
end

subplot(3,1,3);
v = sqrt(state(:,3).^2 + state(:,4).^2);
title('Histogram of Electron Speeds');
histogram(v);
xlabel('Speed (m/s)');
ylabel('Number of particles');

%%
% For the final simulation, an electron density map is created, by creating
% a 2D histogram over space:

density = hist3(state(:,1:2),[200 100]);

% Smooth out the electron density map
N = 20;
sigma = 3;
[x y]=meshgrid(round(-N/2):round(N/2), round(-N/2):round(N/2));
f=exp(-x.^2/(2*sigma^2)-y.^2/(2*sigma^2));
f=f./sum(f(:));
figure(4);
imagesc(conv2(density,f,'same'));
set(gca,'YDir','normal');
title('Electron Density');
xlabel('x (nm)');
ylabel('y (nm)');

%%
% The temperature map is created using a similar procedure. The electron
% velocities are put into bins over space to calculate the temperature at
% different points:
temp_sum_x = zeros(ceil(length/1e-9),ceil(height/1e-9));
temp_sum_y = zeros(ceil(length/1e-9),ceil(height/1e-9));

```

```

temp_num = zeros(ceil(length/1e-9),ceil(height/1e-9));

% Look at velocities of all the particles
for i=1:population_size
    % Find which "bin" it belongs in:
    x = floor(state(i,1)/1e-9);
    y = floor(state(i,2)/1e-9);
    if(x==0)
        x = 1;
    end
    if(y==0)
        y= 1;
    end

    % Add its velocity components to the cumulative count:
    temp_sum_y(x,y) = temp_sum_y(x,y) + state(i,3)^2;
    temp_sum_x(x,y) = temp_sum_x(x,y) + state(i,4)^2;
    temp_num(x,y) = temp_num(x,y) + 1;
end

%%
% Now, with the velocities added up, calculate the temperatures:
temp = (temp_sum_x + temp_sum_y).*/k./2./temp_num;
temp(isnan(temp)) = 0;
temp = temp';

%%
% Like with the density map, perform some smoothing:
N = 20;
sigma = 3;
[x y]=meshgrid(round(-N/2):round(N/2), round(-N/2):round(N/2));
f=exp(-x.^2/(2*sigma^2)-y.^2/(2*sigma^2));
f=f./sum(f(:));
figure(5);
imagesc(conv2(temp,f,'same'));
set(gca,'YDir','normal');
title('Temperature Map');
xlabel('x (nm)');
ylabel('y (nm)');

function box_num = in_box(pos, boxes)


    box_num = 0;
    for i=1:size(boxes,1)
        if(pos(1) > boxes(i,1) && pos(1) < boxes(i,2) && pos(2) > boxes(i,3)
        && pos(2) < boxes(i,4))
            box_num = i;
            return;
        end
    end
end

%%
% The relationship between the temperature map and the electron density map


```

% is very similar as can be seen. However, some the of the electrons have considerably
% higher speeds, and this can be seen on the temperature map.

a)



c)



d)

