



# CS201 DISCRETE MATHEMATICS FOR COMPUTER SCIENCE

Dr. QI WANG

Department of Computer Science and Engineering

Office: Room413, CoE South Tower

Email: [wangqi@sustech.edu.cn](mailto:wangqi@sustech.edu.cn)

# Euclidean Algorithm

- The Euclidean algorithm in pseudocode

## ALGORITHM 1 The Euclidean Algorithm.

```
procedure gcd(a, b: positive integers)
  x := a
  y := b
  while y ≠ 0
    r := x mod y
    x := y
    y := r
  return x{gcd(a, b) is x}
```

The number of **divisions** required to find  $\text{gcd}(a, b)$  is  $O(\log b)$ , where  $a \geq b$ . (this will be proved later.)



# Euclidean Algorithm

- The Euclidean algorithm in pseudocode

## ALGORITHM 1 The Euclidean Algorithm.

```
procedure gcd( $a, b$ : positive integers)
 $x := a$ 
 $y := b$ 
while  $y \neq 0$ 
     $r := x \bmod y$ 
     $x := y$ 
     $y := r$ 
return  $x$  {gcd( $a, b$ ) is  $x$ }
```

The number of **divisions** required to find  $\text{gcd}(a, b)$  is  $O(\log b)$ , where  $a \geq b$ . (this will be proved later.)

**Why ?**



# Euclidean Algorithm

- Key steps in the Euclidean algorithm

$$r_0 = r_1 q_1 + r_2 \quad 0 \leq r_2 < r_1,$$

$$r_1 = r_2 q_2 + r_3 \quad 0 \leq r_3 < r_2,$$

.

.

.

$$r_{n-2} = r_{n-1} q_{n-1} + r_n \quad 0 \leq r_n < r_{n-1},$$

$$r_{n-1} = r_n q_n .$$

# Euclidean Algorithm

- Key steps in the Euclidean algorithm

$$\begin{aligned}r_0 &= r_1 q_1 + r_2 & 0 \leq r_2 < r_1, \\r_1 &= r_2 q_2 + r_3 & 0 \leq r_3 < r_2, \\&\cdot \\&\cdot \\&\cdot \\r_{n-2} &= r_{n-1} q_{n-1} + r_n & 0 \leq r_n < r_{n-1}, \\r_{n-1} &= r_n q_n.\end{aligned}$$

**Observation:**

$$r_{i+2} = r_i \bmod r_{i+1}$$

# Euclidean Algorithm

- Key steps in the Euclidean algorithm

$$r_0 = r_1 q_1 + r_2 \quad 0 \leq r_2 < r_1,$$

$$r_1 = r_2 q_2 + r_3 \quad 0 \leq r_3 < r_2,$$

.

.

.

$$r_{n-2} = r_{n-1} q_{n-1} + r_n \quad 0 \leq r_n < r_{n-1},$$

$$r_{n-1} = r_n q_n.$$

**Observation:**

$$r_{i+2} = r_i \bmod r_{i+1}$$

We claim that  $r_{i+2} < \frac{1}{2} r_i$

# Euclidean Algorithm

- Key steps in the Euclidean algorithm

$$\begin{aligned}r_0 &= r_1 q_1 + r_2 & 0 \leq r_2 < r_1, \\r_1 &= r_2 q_2 + r_3 & 0 \leq r_3 < r_2, \\&\vdots \\&\vdots \\&\vdots \\r_{n-2} &= r_{n-1} q_{n-1} + r_n & 0 \leq r_n < r_{n-1}, \\r_{n-1} &= r_n q_n.\end{aligned}$$

**Observation:**

$$r_{i+2} = r_i \bmod r_{i+1}$$

We claim that  $r_{i+2} < \frac{1}{2} r_i$

Case (i):  $r_{i+1} \leq \frac{1}{2} r_i$ :  $r_{i+2} < r_{i+1} \leq \frac{1}{2} r_i$ .

Case (ii):  $r_{i+1} > \frac{1}{2} r_i$ :  $r_{i+2} = r_i \bmod r_{i+1} = r_i - r_{i+1} < \frac{1}{2} r_i$ .

# Euclidean Algorithm

- Key steps in the Euclidean algorithm

$$r_0 = r_1 q_1 + r_2 \quad 0 \leq r_2 < r_1,$$

$$r_1 = r_2 q_2 + r_3 \quad 0 \leq r_3 < r_2,$$

.

.

.

$$r_{n-2} = r_{n-1} q_{n-1} + r_n \quad 0 \leq r_n < r_{n-1},$$

$$r_{n-1} = r_n q_n.$$

See [Theorem 1 p. 347].

**Observation:**

$$r_{i+2} = r_i \bmod r_{i+1}$$

We claim that  $r_{i+2} < \frac{1}{2} r_i$

Case (i):  $r_{i+1} \leq \frac{1}{2} r_i$ :  $r_{i+2} < r_{i+1} \leq \frac{1}{2} r_i$ .

Case (ii):  $r_{i+1} > \frac{1}{2} r_i$ :  $r_{i+2} = r_i \bmod r_{i+1} = r_i - r_{i+1} < \frac{1}{2} r_i$ .



# Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .



# Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .

- ◇ **linear**: it is a linear combination of previous terms



# Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .

- ◇ **linear**: it is a linear combination of previous terms
- ◇ **homogeneous**: all terms are multiples of  $a_j$ 's



# Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .

- ◇ **linear**: it is a linear combination of previous terms
- ◇ **homogeneous**: all terms are multiples of  $a_j$ 's
- ◇ **degree  $k$** :  $a_n$  is expressed by the previous  $k$  terms



# Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .

- ◇ **linear**: it is a linear combination of previous terms
- ◇ **homogeneous**: all terms are multiples of  $a_j$ 's
- ◇ **degree  $k$** :  $a_n$  is expressed by the previous  $k$  terms
- ◇ **constant coefficients**: coefficients are constants



# Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .

- ◇ **linear**: it is a linear combination of previous terms
- ◇ **homogeneous**: all terms are multiples of  $a_j$ 's
- ◇ **degree  $k$** :  $a_n$  is expressed by the previous  $k$  terms
- ◇ **constant coefficients**: coefficients are constants

By induction, such a recurrence relation is **uniquely** determined by this recurrence relation, and  **$k$  initial conditions**  $a_0, a_1, \dots, a_{k-1}$ .



# Examples of Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .

## Examples

$$P_n = (1.11)P_{n-1}$$

$$f_n = f_{n-1} + f_{n-2}$$

$$a_n = a_{n-1} + a_{n-2}^2$$

$$H_n = 2H_{n-1} + 1$$

$$B_n = nB_{n-1}$$



# Examples of Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .

## Examples

$$P_n = (1.11)P_{n-1} \quad \text{linear homogeneous recurrence relation of degree 1}$$

$$f_n = f_{n-1} + f_{n-2}$$

$$a_n = a_{n-1} + a_{n-2}^2$$

$$H_n = 2H_{n-1} + 1$$

$$B_n = nB_{n-1}$$





# Examples of Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .

## Examples

$$P_n = (1.11)P_{n-1} \quad \text{linear homogeneous recurrence relation of degree 1}$$

$$f_n = f_{n-1} + f_{n-2} \quad \text{linear homogeneous recurrence relation of degree 2}$$

$$a_n = a_{n-1} + a_{n-2}^2$$

$$H_n = 2H_{n-1} + 1$$

$$B_n = nB_{n-1}$$



# Examples of Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .

## Examples

$P_n = (1.11)P_{n-1}$  linear homogeneous recurrence relation of degree 1

$f_n = f_{n-1} + f_{n-2}$  linear homogeneous recurrence relation of degree 2

$a_n = a_{n-1} + a_{n-2}^2$  NOT linear

$$H_n = 2H_{n-1} + 1$$

$$B_n = nB_{n-1}$$



# Examples of Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .

## Examples

$P_n = (1.11)P_{n-1}$  linear homogeneous recurrence relation of degree 1

$f_n = f_{n-1} + f_{n-2}$  linear homogeneous recurrence relation of degree 2

$a_n = a_{n-1} + a_{n-2}^2$  NOT linear

$H_n = 2H_{n-1} + 1$  NOT homogeneous

$B_n = nB_{n-1}$



# Examples of Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .

## Examples

$P_n = (1.11)P_{n-1}$  linear homogeneous recurrence relation of degree 1

$f_n = f_{n-1} + f_{n-2}$  linear homogeneous recurrence relation of degree 2

$a_n = a_{n-1} + a_{n-2}^2$  NOT linear

$H_n = 2H_{n-1} + 1$  NOT homogeneous

$B_n = nB_{n-1}$  coefficients are not constants



# Solving Linear Recurrence Relations

- **Example** Consider the recurrence relation

$$a_n = 2a_{n-1} - a_{n-2},$$

Which of the following are solutions?

◇  $a_n = 3n$ :

◇  $a_n = 2^n$ :

◇  $a_n = 5$ :



# Solving Linear Recurrence Relations

- **Example** Consider the recurrence relation

$$a_n = 2a_{n-1} - a_{n-2},$$

Which of the following are solutions?

◇  $a_n = 3n$ : YES

◇  $a_n = 2^n$ : NO

◇  $a_n = 5$ : YES



# Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .



# Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .

**Basic idea:** Look for solutions of the form  $a_n = r^n$ , where  $r$  is a constant.





# Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .

**Basic idea:** Look for solutions of the form  $a_n = r^n$ , where  $r$  is a constant.

- ◇ Bring  $a_n = r^n$  back to the recurrence relation:

$$\begin{aligned} & r^n = c_1 r^{n-1} + c_2 r^{n-2} + \cdots + c_k r^{n-k}, \\ \text{i.e.,} \quad & r^{n-k} (r^k - c_1 r^{k-1} - \cdots - c_k) = 0 \end{aligned}$$



# Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .

**Basic idea:** Look for solutions of the form  $a_n = r^n$ , where  $r$  is a constant.

- ◇ Bring  $a_n = r^n$  back to the recurrence relation:

$$\begin{aligned} & r^n = c_1 r^{n-1} + c_2 r^{n-2} + \cdots + c_k r^{n-k}, \\ \text{i.e.,} \quad & r^{n-k} (r^k - c_1 r^{k-1} - \cdots - c_k) = 0 \end{aligned}$$

- ◇ The solutions to the *characteristic equation* can yield an explicit formula for the sequence.

$$(r^k - c_1 r^{k-1} - \cdots - c_k) = 0$$



# Recall: Problem IV

## ■ Fibonacci number

$$F_0 = 0, F_1 = 1, F_n = F_{n-1} + F_{n-2} \text{ for } n \geq 2$$



# Recall: Problem IV

## ■ Fibonacci number

$$F_0 = 0, F_1 = 1, F_n = F_{n-1} + F_{n-2} \text{ for } n \geq 2$$

◇ What is the closed-form expression of  $F_n$ ?



# Recall: Problem IV

## ■ Fibonacci number

$$F_0 = 0, F_1 = 1, F_n = F_{n-1} + F_{n-2} \text{ for } n \geq 2$$

◇ What is the closed-form expression of  $F_n$ ?

Consider  $x^n = x^{n-1} + x^{n-2}$ , with  $x \neq 0$ . There are two different roots

$$\phi = \frac{1 + \sqrt{5}}{2}, \quad \psi = \frac{1 - \sqrt{5}}{2}$$

Then  $F_n$  can be the form of  $a\phi^n + b\psi^n$ . By  $F_0 = 0$  and  $F_1 = 1$ , we have  $a + b = 0$  and  $\phi a + \psi b = 1$ , leading to  $a = \frac{1}{\sqrt{5}}$ ,  $b = -a$ . Therefore,

$$F_n = \frac{\phi^n - \psi^n}{\sqrt{5}}$$



# Solving Linear Recurrence Relations of degree 2

- Consider an arbitrary linear homogeneous relation of degree 2 with constant coefficients:

$$a_n = c_1 a_{n-1} + c_2 a_{n-2}.$$



# Solving Linear Recurrence Relations of degree 2

- Consider an arbitrary linear homogeneous relation of degree 2 with constant coefficients:

$$a_n = c_1 a_{n-1} + c_2 a_{n-2}.$$

The characteristic equation (CE) is:

$$r^2 - c_1 r - c_2 = 0.$$



# Solving Linear Recurrence Relations of degree 2

- Consider an arbitrary linear homogeneous relation of degree 2 with constant coefficients:

$$a_n = c_1 a_{n-1} + c_2 a_{n-2}.$$

The characteristic equation (CE) is:

$$r^2 - c_1 r - c_2 = 0.$$

**Theorem** If this CE has 2 roots  $r_1 \neq r_2$ , then the sequence  $\{a_n\}$  is a solution of the recurrence relation **if and only if**  $a_n = \alpha_1 r_1^n + \alpha_2 r_2^n$  for  $n \geq 0$  and constants  $\alpha_1, \alpha_2$ .





# Solving Linear Recurrence Relations of degree 2

- Consider an arbitrary linear homogeneous relation of degree 2 with constant coefficients:

$$a_n = c_1 a_{n-1} + c_2 a_{n-2}.$$

The characteristic equation (CE) is:

$$r^2 - c_1 r - c_2 = 0.$$

**Theorem** If this CE has 2 roots  $r_1 \neq r_2$ , then the sequence  $\{a_n\}$  is a solution of the recurrence relation **if and only if**  $a_n = \alpha_1 r_1^n + \alpha_2 r_2^n$  for  $n \geq 0$  and constants  $\alpha_1, \alpha_2$ .

**Proof?**



# Solving Linear Recurrence Relations of degree 2

- Consider an arbitrary linear homogeneous relation of degree 2 with constant coefficients:

$$a_n = c_1 a_{n-1} + c_2 a_{n-2}.$$

The characteristic equation (CE) is:

$$r^2 - c_1 r - c_2 = 0.$$

**Theorem** If this CE has 2 roots  $r_1 \neq r_2$ , then the sequence  $\{a_n\}$  is a solution of the recurrence relation **if and only if**  $a_n = \alpha_1 r_1^n + \alpha_2 r_2^n$  for  $n \geq 0$  and constants  $\alpha_1, \alpha_2$ .

**Proof?**

See [Theorem 1 p. 515].



# Solving Linear Recurrence Relations of degree 2

- **Example 1**  $a_n = a_{n-1} + 2a_{n-2}$ , with  $a_0 = 2$ ,  $a_1 = 7$



# Solving Linear Recurrence Relations of degree 2

- **Example 1**  $a_n = a_{n-1} + 2a_{n-2}$ , with  $a_0 = 2$ ,  $a_1 = 7$

The *characteristic equation* is

$$r^2 - r - 2 = 0.$$



# Solving Linear Recurrence Relations of degree 2

- **Example 1**  $a_n = a_{n-1} + 2a_{n-2}$ , with  $a_0 = 2$ ,  $a_1 = 7$

The *characteristic equation* is

$$r^2 - r - 2 = 0.$$

Two roots are 2 and  $-1$ . So, assume that

$$a_n = \alpha_1 2^n + \alpha_2 (-1)^n.$$



# Solving Linear Recurrence Relations of degree 2

- **Example 1**  $a_n = a_{n-1} + 2a_{n-2}$ , with  $a_0 = 2$ ,  $a_1 = 7$

The *characteristic equation* is

$$r^2 - r - 2 = 0.$$

Two roots are 2 and  $-1$ . So, assume that

$$a_n = \alpha_1 2^n + \alpha_2 (-1)^n.$$

By the two initial conditions, we have

$$a_0 = \alpha_1 + \alpha_2 = 2$$

$$a_1 = 2\alpha_1 - \alpha_2 = 7$$



# Solving Linear Recurrence Relations of degree 2

- **Example 1**  $a_n = a_{n-1} + 2a_{n-2}$ , with  $a_0 = 2$ ,  $a_1 = 7$

The *characteristic equation* is

$$r^2 - r - 2 = 0.$$

Two roots are 2 and  $-1$ . So, assume that

$$a_n = \alpha_1 2^n + \alpha_2 (-1)^n.$$

By the two initial conditions, we have

$$a_0 = \alpha_1 + \alpha_2 = 2$$

$$a_1 = 2\alpha_1 - \alpha_2 = 7$$

We get  $\alpha_1 = 3$  and  $\alpha_2 = -1$ . Thus,

$$a_n = 3 \cdot 2^n - (-1)^n$$



# Solving Linear Recurrence Relations of degree 2

- **Example 2**  $a_n = 7a_{n-1} - 10a_{n-2}$ , with  $a_0 = 2$ ,  $a_1 = 1$





# Solving Linear Recurrence Relations of degree 2

- **Example 2**  $a_n = 7a_{n-1} - 10a_{n-2}$ , with  $a_0 = 2$ ,  $a_1 = 1$

The *characteristic equation* is

$$r^2 - 7r + 10 = 0.$$



# Solving Linear Recurrence Relations of degree 2

- **Example 2**  $a_n = 7a_{n-1} - 10a_{n-2}$ , with  $a_0 = 2$ ,  $a_1 = 1$

The *characteristic equation* is

$$r^2 - 7r + 10 = 0.$$

Two roots are 2 and 5. So, assume that

$$a_n = \alpha_1 2^n + \alpha_2 5^n.$$



# Solving Linear Recurrence Relations of degree 2

- **Example 2**  $a_n = 7a_{n-1} - 10a_{n-2}$ , with  $a_0 = 2$ ,  $a_1 = 1$

The *characteristic equation* is

$$r^2 - 7r + 10 = 0.$$

Two roots are 2 and 5. So, assume that

$$a_n = \alpha_1 2^n + \alpha_2 5^n.$$

By the two initial conditions, we have

$$a_0 = \alpha_1 + \alpha_2 = 2$$

$$a_1 = 2\alpha_1 + 5\alpha_2 = 1$$



# Solving Linear Recurrence Relations of degree 2

- **Example 2**  $a_n = 7a_{n-1} - 10a_{n-2}$ , with  $a_0 = 2$ ,  $a_1 = 1$

The *characteristic equation* is

$$r^2 - 7r + 10 = 0.$$

Two roots are 2 and 5. So, assume that

$$a_n = \alpha_1 2^n + \alpha_2 5^n.$$

By the two initial conditions, we have

$$a_0 = \alpha_1 + \alpha_2 = 2$$

$$a_1 = 2\alpha_1 + 5\alpha_2 = 1$$

We get  $\alpha_1 = 3$  and  $\alpha_2 = -1$ . Thus,

$$a_n = 3 \cdot 2^n - 5^n$$



# Solving Linear Recurrence Relations of degree $k$

- Consider an arbitrary linear homogeneous relation of degree  $k$  with constant coefficients:

$$a_n = \sum_{i=1}^k c_i a_{n-i}.$$



# Solving Linear Recurrence Relations of degree $k$

- Consider an arbitrary linear homogeneous relation of degree  $k$  with constant coefficients:

$$a_n = \sum_{i=1}^k c_i a_{n-i}.$$

The characteristic equation (CE) is:

$$r^k - \sum_{i=1}^k c_i r^{k-i} = 0.$$



# Solving Linear Recurrence Relations of degree $k$

- Consider an arbitrary linear homogeneous relation of degree  $k$  with constant coefficients:

$$a_n = \sum_{i=1}^k c_i a_{n-i}.$$

The characteristic equation (CE) is:

$$r^k - \sum_{i=1}^k c_i r^{k-i} = 0.$$

**Theorem** If this CE has  $k$  distinct roots  $r_i$ , then the solutions to the recurrence are of the form

$$a_n = \sum_{i=1}^k \alpha_i r_i^n$$

for all  $n \geq 0$ , where the  $\alpha_i$ 's are constants.



# Solving Linear Recurrence Relations of degree $k$

- Consider an arbitrary linear homogeneous relation of degree  $k$  with constant coefficients:

$$a_n = \sum_{i=1}^k c_i a_{n-i}.$$

The characteristic equation (CE) is:

$$r^k - \sum_{i=1}^k c_i r^{k-i} = 0.$$

**Theorem** If this CE has  $k$  distinct roots  $r_i$ , then the solutions to the recurrence are of the form

$$a_n = \sum_{i=1}^k \alpha_i r_i^n$$

for all  $n \geq 0$ , where the  $\alpha_i$ 's are constants.

**Example**  $a_n = 2a_{n-1} + 5a_{n-2} - 6a_{n-3}$





# The Case of Degenerate Roots

- **Theorem** If the CE  $r^2 - c_1r - c_2 = 0$  has **only 1** root  $r_0$ , then

$$a_n = \alpha_1 r_0^n + \alpha_2 n r_0^n,$$

for all  $n \geq 0$  and two constants  $\alpha_1$  and  $\alpha_2$ .



# The Case of Degenerate Roots

- **Theorem** If the CE  $r^2 - c_1 r - c_2 = 0$  has **only 1** root  $r_0$ , then

$$a_n = \alpha_1 r_0^n + \alpha_2 n r_0^n,$$

for all  $n \geq 0$  and two constants  $\alpha_1$  and  $\alpha_2$ .

**Proof?**



# The Case of Degenerate Roots

- **Theorem** If the CE  $r^2 - c_1 r - c_2 = 0$  has **only 1** root  $r_0$ , then

$$a_n = \alpha_1 r_0^n + \alpha_2 n r_0^n,$$

for all  $n \geq 0$  and two constants  $\alpha_1$  and  $\alpha_2$ .

**Proof?**

Exercise.



# The Case of Degenerate Roots

- **Example**  $a_n = 4a_{n-1} - 4a_{n-2}$ , with  $a_0 = 1$ ,  $a_1 = 0$



# The Case of Degenerate Roots

- **Example**  $a_n = 4a_{n-1} - 4a_{n-2}$ , with  $a_0 = 1$ ,  $a_1 = 0$

The *characteristic equation* is

$$r^2 - 4r + 4 = 0.$$



# The Case of Degenerate Roots

- **Example**  $a_n = 4a_{n-1} - 4a_{n-2}$ , with  $a_0 = 1$ ,  $a_1 = 0$

The *characteristic equation* is

$$r^2 - 4r + 4 = 0.$$

The only root is 2. So, assume that

$$a_n = \alpha_1 2^n + \alpha_2 n 2^n.$$



# The Case of Degenerate Roots

- **Example**  $a_n = 4a_{n-1} - 4a_{n-2}$ , with  $a_0 = 1$ ,  $a_1 = 0$

The *characteristic equation* is

$$r^2 - 4r + 4 = 0.$$

The only root is 2. So, assume that

$$a_n = \alpha_1 2^n + \alpha_2 n 2^n.$$

By the two initial conditions, we have

$$a_0 = \alpha_1 = 1$$

$$a_1 = 2\alpha_1 + 2\alpha_2 = 0$$



# The Case of Degenerate Roots

- **Example**  $a_n = 4a_{n-1} - 4a_{n-2}$ , with  $a_0 = 1$ ,  $a_1 = 0$

The *characteristic equation* is

$$r^2 - 4r + 4 = 0.$$

The only root is 2. So, assume that

$$a_n = \alpha_1 2^n + \alpha_2 n 2^n.$$

By the two initial conditions, we have

$$a_0 = \alpha_1 = 1$$

$$a_1 = 2\alpha_1 + 2\alpha_2 = 0$$

We get  $\alpha_1 = 1$  and  $\alpha_2 = -1$ . Thus,

$$a_n = 2^n - n2^n$$





# The Case of Degenerate Roots in General

- **Theorem** [Theorem 4, p.519] Suppose that there are  $t$  roots  $r_1, \dots, r_t$  with multiplicities  $m_1, \dots, m_t$ . Then

$$a_n = \sum_{i=1}^t \left( \sum_{j=0}^{m_i-1} \alpha_{i,j} n^j \right) r_i^n,$$

for all  $n \geq 0$  and constants  $\alpha_{i,j}$ .



# The Case of Degenerate Roots in General

- **Theorem** [Theorem 4, p.519] Suppose that there are  $t$  roots  $r_1, \dots, r_t$  with multiplicities  $m_1, \dots, m_t$ . Then

$$a_n = \sum_{i=1}^t \left( \sum_{j=0}^{m_i-1} \alpha_{i,j} n^j \right) r_i^n,$$

for all  $n \geq 0$  and constants  $\alpha_{i,j}$ .

## Example

$$a_n = -3a_{n-1} - 3a_{n-2} - a_{n-3} \text{ with } a_0 = 1, a_1 = -2, \\ a_2 = -1$$



# Linear Nonhomogeneous Recurrence Relations

- **Definition** A *linear nonhomogeneous relation* with constant coefficients may contain some terms  $F(n)$  that depend only on  $n$

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k} + F(n).$$

The recurrence relation

$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k}$  is called the *associated homogeneous recurrence relation*.



# Linear Nonhomogeneous Recurrence Relations

- **Theorem** If  $a_n = p(n)$  is any particular solution to the linear nonhomogeneous relation with constant coefficients,

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k} + F(n),$$

Then all its solutions are of the form

$$a_n = p(n) + h(n),$$

where  $a_n = h(n)$  is any solution to the associated homogeneous recurrence relation

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k}.$$



# Solving Linear Nonhomogeneous Recurrence Relations

- **Example**  $a_n = 3a_{n-1} + 2n$ . Which solution has  $a_1 = 3$ ?



# Solving Linear Nonhomogeneous Recurrence Relations

- **Example**  $a_n = 3a_{n-1} + 2n$ . Which solution has  $a_1 = 3$ ?

The *characteristic equation* of the associated linear homogeneous recurrence relation is  $r^2 - 3r = 0$ . Thus, the solution to the original problem are all of the form

$$a_n = \alpha 3^n + p(n) .$$



# Solving Linear Nonhomogeneous Recurrence Relations

- **Example**  $a_n = 3a_{n-1} + 2n$ . Which solution has  $a_1 = 3$ ?

The *characteristic equation* of the associated linear homogeneous recurrence relation is  $r^2 - 3r = 0$ . Thus, the solution to the original problem are all of the form

$$a_n = \alpha 3^n + p(n) .$$

We try a degree- $t$  polynomial as the particular solution  $p(n)$ .



# Solving Linear Nonhomogeneous Recurrence Relations

- **Example**  $a_n = 3a_{n-1} + 2n$ . Which solution has  $a_1 = 3$ ?

The *characteristic equation* of the associated linear homogeneous recurrence relation is  $r^2 - 3r = 0$ . Thus, the solution to the original problem are all of the form  $a_n = \alpha 3^n + p(n)$ .

We try a degree- $t$  polynomial as the particular solution  $p(n)$ .

Let  $p(n) = cn + d$ , then

$$cn + d = 3(c(n-1) + d) + 2n, \text{ which means } (2c + 2)n + (2d - 3c) = 0.$$





# Solving Linear Nonhomogeneous Recurrence Relations

- **Example**  $a_n = 3a_{n-1} + 2n$ . Which solution has  $a_1 = 3$ ?

The *characteristic equation* of the associated linear homogeneous recurrence relation is  $r^2 - 3r = 0$ . Thus, the solution to the original problem are all of the form  $a_n = \alpha 3^n + p(n)$ .

We try a degree- $t$  polynomial as the particular solution  $p(n)$ .

Let  $p(n) = cn + d$ , then

$$cn + d = 3(c(n-1) + d) + 2n, \text{ which means } (2c + 2)n + (2d - 3c) = 0.$$

We get  $c = -1$  and  $d = -3/2$ . Thus,

$$p(n) = -n - 3/2$$



# Generating Functions

- We may use *generating functions* to **characterize** sequences.



# Generating Functions

- We may use *generating functions* to **characterize** sequences.
  - ◇ The sequence  $\{a_k\}$  with  $a_k = 3$



# Generating Functions

- We may use *generating functions* to **characterize** sequences.

◇ The sequence  $\{a_k\}$  with  $a_k = 3$

$$\sum_{k=0}^{\infty} 3x^k$$



# Generating Functions

- We may use *generating functions* to **characterize** sequences.

- ◇ The sequence  $\{a_k\}$  with  $a_k = 3$

$$\sum_{k=0}^{\infty} 3x^k$$

- ◇ The sequence  $\{a_k\}$  with  $a_k = 2^k$



# Generating Functions

- We may use *generating functions* to **characterize** sequences.

◇ The sequence  $\{a_k\}$  with  $a_k = 3$

$$\sum_{k=0}^{\infty} 3x^k$$

◇ The sequence  $\{a_k\}$  with  $a_k = 2^k$

$$\sum_{k=0}^{\infty} 2^k x^k$$



# Generating Functions

- We may use *generating functions* to **characterize** sequences.

- ◇ The sequence  $\{a_k\}$  with  $a_k = 3$  
$$\sum_{k=0}^{\infty} 3x^k$$

- ◇ The sequence  $\{a_k\}$  with  $a_k = 2^k$  
$$\sum_{k=0}^{\infty} 2^k x^k$$

**Definition** The *generating function* for the sequence  $a_0, a_1, \dots, a_k, \dots$  of real numbers is the infinite series

$$G(x) = a_0 + a_1x + \cdots + a_kx^k$$



# Generating Functions for Finite Sequences

- A finite sequence  $a_0, a_1, \dots, a_n$  can be easily extended by setting  $a_{n+1} = a_{n+2} = \dots = 0$





# Generating Functions for Finite Sequences

- A finite sequence  $a_0, a_1, \dots, a_n$  can be easily extended by setting  $a_{n+1} = a_{n+2} = \dots = 0$

The *generating function*  $G(x)$  of this infinite sequence  $\{a_n\}$  is a polynomial of degree  $n$ , i.e.,

$$G(x) = a_0 + a_1x + \dots + a_nx^n$$



# Generating Functions for Finite Sequences

- A finite sequence  $a_0, a_1, \dots, a_n$  can be easily extended by setting  $a_{n+1} = a_{n+2} = \dots = 0$

The *generating function*  $G(x)$  of this infinite sequence  $\{a_n\}$  is a polynomial of degree  $n$ , i.e.,

$$G(x) = a_0 + a_1x + \dots + a_nx^n$$

- ◇ What is the generating function for the sequence  $a_0, a_1, \dots, a_m$ , with  $a_k = C(m, k)$ ?



# Generating Functions for Finite Sequences

- A finite sequence  $a_0, a_1, \dots, a_n$  can be easily extended by setting  $a_{n+1} = a_{n+2} = \dots = 0$

The *generating function*  $G(x)$  of this infinite sequence  $\{a_n\}$  is a polynomial of degree  $n$ , i.e.,

$$G(x) = a_0 + a_1x + \dots + a_nx^n$$

- ◇ What is the generating function for the sequence  $a_0, a_1, \dots, a_m$ , with  $a_k = C(m, k)$ ?

$$G(x) = C(m, 0) + \dots + C(m,$$



# Examples

◇  $G(x) = 1/(1 - x)$  for  $|x| < 1$



# Examples

◇  $G(x) = 1/(1 - x)$  for  $|x| < 1$

$1, 1, 1, 1, 1, \dots$



# Examples

◇  $G(x) = 1/(1 - x)$  for  $|x| < 1$

$1, 1, 1, 1, 1, \dots$

◇  $G(x) = 1/(1 - ax)$  for  $|ax| < 1$



# Examples

◇  $G(x) = 1/(1 - x)$  for  $|x| < 1$

$1, 1, 1, 1, 1, \dots$

◇  $G(x) = 1/(1 - ax)$  for  $|ax| < 1$

$1, a, a^2, a^3, a^4, \dots$



# Examples

◇  $G(x) = 1/(1 - x)$  for  $|x| < 1$

$1, 1, 1, 1, 1, \dots$

◇  $G(x) = 1/(1 - ax)$  for  $|ax| < 1$

$1, a, a^2, a^3, a^4, \dots$

◇  $G(x) = 1/(1 - x)^2$  for  $|x| < 1$





# Examples

◇  $G(x) = 1/(1 - x)$  for  $|x| < 1$

$$1, 1, 1, 1, 1, \dots$$

◇  $G(x) = 1/(1 - ax)$  for  $|ax| < 1$

$$1, a, a^2, a^3, a^4, \dots$$

◇  $G(x) = 1/(1 - x)^2$  for  $|x| < 1$

$$1, 2, 3, 4, 5, \dots$$



# Operations of Generating Functions

- **Theorem** Let  $f(x) = \sum_{k=0}^{\infty} a_k x^k$ , and  $g(x) = \sum_{k=0}^{\infty} b_k x^k$ .  
Then

$$f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$$

$$f(x)g(x) = \sum_{k=0}^{\infty} \left( \sum_{j=0}^k a_j b_{k-j} \right) x^k$$



# Operations of Generating Functions

■ **Theorem** Let  $f(x) = \sum_{k=0}^{\infty} a_k x^k$ , and  $g(x) = \sum_{k=0}^{\infty} b_k x^k$ .  
Then

$$f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$$

$$f(x)g(x) = \sum_{k=0}^{\infty} \left( \sum_{j=0}^k a_j b_{k-j} \right) x^k$$

◇  $G(x) = 1/(1-x)^2$  for  $|x| < 1$



# Operations of Generating Functions

■ **Theorem** Let  $f(x) = \sum_{k=0}^{\infty} a_k x^k$ , and  $g(x) = \sum_{k=0}^{\infty} b_k x^k$ .  
Then

$$f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$$

$$f(x)g(x) = \sum_{k=0}^{\infty} \left( \sum_{j=0}^k a_j b_{k-j} \right) x^k$$

◇  $G(x) = 1/(1-x)^2$  for  $|x| < 1$

$$f(x) = 1/(1-x), \quad g(x) = 1/(1-x)$$

$$G(x) = f(x)g(x) = \sum_{k=0}^{\infty} \left( \sum_{j=0}^k 1 \right) x^k = \sum_{k=0}^{\infty} (k+1) x^k$$



# Operations of Generating Functions

■ **Theorem** Let  $f(x) = \sum_{k=0}^{\infty} a_k x^k$ , and  $g(x) = \sum_{k=0}^{\infty} b_k x^k$ .  
Then

$$f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$$

$$f(x)g(x) = \sum_{k=0}^{\infty} \left( \sum_{j=0}^k a_j b_{k-j} \right) x^k$$

◇  $G(x) = 1/(1 - ax)^2$  for  $|x| < 1$



# Operations of Generating Functions

■ **Theorem** Let  $f(x) = \sum_{k=0}^{\infty} a_k x^k$ , and  $g(x) = \sum_{k=0}^{\infty} b_k x^k$ .  
Then

$$f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$$

$$f(x)g(x) = \sum_{k=0}^{\infty} \left( \sum_{j=0}^k a_j b_{k-j} \right) x^k$$

◇  $G(x) = 1/(1 - ax)^2$  for  $|x| < 1$

$$f(x) = 1/(1 - ax), \quad g(x) = 1/(1 - ax)$$

$$G(x) = f(x)g(x) = \sum_{k=0}^{\infty} \left( \sum_{j=0}^k a^j a^{k-j} \right) x^k = \sum_{k=0}^{\infty} (k+1) a^k x^k$$



# Useful Generating Functions

$$(1 + x)^n = \sum_{k=0}^n C(n, k) x^k$$

$$(1 + ax)^n = \sum_{k=0}^n C(n, k) a^k x^k$$

$$(1 + x^r)^n = \sum_{k=0}^n C(n, k) x^{rk}$$



# Useful Generating Functions

$$(1 + x)^n = \sum_{k=0}^n C(n, k)x^k$$

$$(1 + ax)^n = \sum_{k=0}^n C(n, k)a^k x^k$$

$$(1 + x^r)^n = \sum_{k=0}^n C(n, k)x^{rk}$$

$$\frac{1-x^{n+1}}{1-x} = \sum_{k=0}^n x^k = 1 + x + x^2 + \dots + x^n$$

$$\frac{1}{1-x} = \sum_{k=0}^{\infty} x^k = 1 + x + x^2 + \dots$$

$$\frac{1}{1-ax} = \sum_{k=0}^{\infty} a^k x^k = 1 + ax + a^2 x^2 + \dots$$

$$\frac{1}{1-x^r} = \sum_{k=0}^{\infty} x^{rk} = 1 + x^r + x^{2r} + \dots$$





# Useful Generating Functions

$$(1 + x)^n = \sum_{k=0}^n C(n, k)x^k$$

$$(1 + ax)^n = \sum_{k=0}^n C(n, k)a^k x^k$$

$$(1 + x^r)^n = \sum_{k=0}^n C(n, k)x^{rk}$$

$$\frac{1-x^{n+1}}{1-x} = \sum_{k=0}^n x^k = 1 + x + x^2 + \dots + x^n$$

$$\frac{1}{1-x} = \sum_{k=0}^{\infty} x^k = 1 + x + x^2 + \dots$$

$$\frac{1}{1-ax} = \sum_{k=0}^{\infty} a^k x^k = 1 + ax + a^2 x^2 + \dots$$

$$\frac{1}{1-x^r} = \sum_{k=0}^{\infty} x^{rk} = 1 + x^r + x^{2r} + \dots$$

$$\frac{1}{(1-x)^2} = \sum_{k=0}^{\infty} (k+1)x^k = 1 + 2x + 3x^2 + \dots$$



# Useful Generating Functions

$$\frac{1}{(1-x)^n} = \sum_{k=0}^{\infty} C(n+k-1, k)x^k$$

$$\frac{1}{(1+x)^n} = \sum_{k=0}^{\infty} C(n+k-1, k)(-1)^k x^k$$

$$\frac{1}{(1-ax)^n} = \sum_{k=0}^{\infty} C(n+k-1, k)a^k x^k$$



# Useful Generating Functions

$$\frac{1}{(1-x)^n} = \sum_{k=0}^{\infty} C(n+k-1, k)x^k$$

$$\frac{1}{(1+x)^n} = \sum_{k=0}^{\infty} C(n+k-1, k)(-1)^k x^k$$

$$\frac{1}{(1-ax)^n} = \sum_{k=0}^{\infty} C(n+k-1, k)a^k x^k$$

$$e^x = \sum_{k=0}^{\infty} \frac{x^k}{k!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$$

$$\ln(1+x) = \sum_{k=0}^{\infty} \frac{(-1)^{k+1} x^k}{k} = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$$



# Counting and Generating Functions

- **Problem 1** How many solutions are there to the equation

$$x_1 + x_2 + x_3 = 17,$$

where  $x_1, x_2, x_3$  are **nonnegative** integers?



# Counting and Generating Functions

- **Problem 1** How many solutions are there to the equation

$$x_1 + x_2 + x_3 = 17,$$

where  $x_1, x_2, x_3$  are **nonnegative** integers?

This is **equivalent** to the problem of  $r$ -combinations from a set with  $n$  elements when **repetition** is allowed.



# Counting and Generating Functions

- **Problem 1** How many solutions are there to the equation

$$x_1 + x_2 + x_3 = 17,$$

where  $x_1, x_2, x_3$  are **nonnegative** integers?

This is **equivalent** to the problem of  $r$ -combinations from a set with  $n$  elements when **repetition** is allowed.

$$C(n + r - 1, r) = C(19, 17) = C(19, 2)$$



# Counting and Generating Functions

- **Problem 2** Find the number of solutions of

$$x_1 + x_2 + x_3 = 17,$$

where  $x_1, x_2, x_3$  are **nonnegative** integers with  $2 \leq x_1 \leq 5$ ,  
 $3 \leq x_2 \leq 6$ ,  $4 \leq x_3 \leq 7$ .



# Counting and Generating Functions

- **Problem 2** Find the number of solutions of

$$x_1 + x_2 + x_3 = 17,$$

where  $x_1, x_2, x_3$  are **nonnegative** integers with  $2 \leq x_1 \leq 5$ ,  
 $3 \leq x_2 \leq 6$ ,  $4 \leq x_3 \leq 7$ .

Using *generating functions*, the number is the **coefficient** of  $x^{17}$  in the expansion of

$$(x^2 + x^3 + x^4 + x^5)(x^3 + x^4 + x^5 + x^6)(x^4 + x^5 + x^6 + x^7)$$





# Counting and Generating Functions

- **Problem 3** In how many ways can eight identical cookies be distributed among three distinct children if each child receives at least two cookies and no more than four cookies?



# Counting and Generating Functions

- **Problem 3** In how many ways can eight identical cookies be distributed among three distinct children if each child receives at least two cookies and no more than four cookies?

The coefficient of  $x^8$  in the expansion

$$(x^2 + x^3 + x^4)^3$$



# Counting and Generating Functions

- **Problem 4** Use generating functions to find the number of  $k$ -combinations of a set with  $n$  elements,  $C(n, k)$ .



# Counting and Generating Functions

- **Problem 4** Use **generating functions** to find the number of  **$k$ -combinations of a set with  $n$  elements**,  $C(n, k)$ .

Each of the  $n$  elements in the set contributes the term  $(1 + x)$  to the generating function  $f(x) = \sum_{k=0}^n a^k x^k$ .  
Hence,  $f(x) = (1 + x)^n$ .

Then by the **binomial theorem**, we have  $a_k = \binom{n}{k}$ .



# Next Lecture

- relation ...

