



# CS201 DISCRETE MATHEMATICS FOR COMPUTER SCIENCE

Dr. QI WANG

Department of Computer Science and Engineering

Office: Room413, CoE South Tower

Email: [wangqi@sustech.edu.cn](mailto:wangqi@sustech.edu.cn)

# First-Order Linear Recurrences

- A recurrence of the form  $T(n) = f(n)T(n-1) + g(n)$  is called a *first-order linear recurrence*.
  - ◇ **First Order** because it only depends upon going back one step, i.e.,  $T(n-1)$   
If it depends upon  $T(n-2)$ , it would be a **second-order** recurrence, e.g.,  $T(n) = T(n-1) + 2T(n-2)$ .
  - ◇ **Linear** because  $T(n-1)$  only appears to the **first power**.  
Something like  $T(n) = (T(n-1))^2 + 3$  would be a **non-linear** first-order recurrence relation.



# First-Order Linear Recurrences

- $T(n) = f(n)T(n-1) + g(n)$



# First-Order Linear Recurrences

- $T(n) = f(n)T(n-1) + g(n)$

When  $f(n)$  is a constant, say  $r$ , the general solution is almost as easy as we derived before. Iterating the recurrence gives

$$\begin{aligned}T(n) &= rT(n-1) + g(n) \\&= r(rT(n-2) + g(n-1)) + g(n) \\&= r^2 T(n-2) + rg(n-1) + g(n) \\&= r^3 T(n-3) + r^2 g(n-2) + rg(n-1) + g(n) \\&\vdots \\&= r^n T(0) + \sum_{i=0}^{n-1} r^i g(n-i)\end{aligned}$$



# First-Order Linear Recurrences

- **Theorem** For any positive constants  $a$  and  $r$ , and any function  $g$  defined on nonnegative integers, the solution to the first-order linear recurrence

$$T(n) = \begin{cases} rT(n-1) + g(n) & \text{if } n > 0 \\ a & \text{if } n = 0 \end{cases}$$

is

$$T(n) = r^n a + \sum_{i=1}^n r^{n-i} g(i).$$



# First-Order Linear Recurrences

- **Theorem** For any positive constants  $a$  and  $r$ , and any function  $g$  defined on nonnegative integers, the solution to the first-order linear recurrence

$$T(n) = \begin{cases} rT(n-1) + g(n) & \text{if } n > 0 \\ a & \text{if } n = 0 \end{cases}$$

is

$$T(n) = r^n a + \sum_{i=1}^n r^{n-i} g(i).$$

**Proof by induction**



# Examples

- Solve  $T(n) = 4T(n-1) + 2^n$  with  $T(0) = 6$



# Examples

- Solve  $T(n) = 4T(n-1) + 2^n$  with  $T(0) = 6$

$$\begin{aligned}T(n) &= 6 \cdot 4^n + \sum_{i=1}^n 4^{n-i} \cdot 2^i \\&= 6 \cdot 4^n + 4^n \sum_{i=1}^n 4^{-i} \cdot 2^i \\&= 6 \cdot 4^n + 4^n \sum_{i=1}^n \left(\frac{1}{2}\right)^i \\&= 6 \cdot 4^n + \left(1 - \frac{1}{2^n}\right) \cdot 4^n \\&= 7 \cdot 4^n - 2^n.\end{aligned}$$





# Examples

- Solve  $T(n) = 3T(n-1) + n$  with  $T(0) = 10$



# Examples

- Solve  $T(n) = 3T(n-1) + n$  with  $T(0) = 10$

$$\begin{aligned} T(n) &= 10 \cdot 3^n + \sum_{i=1}^n 3^{n-i} \cdot i \\ &= 10 \cdot 3^n + 3^n \sum_{i=1}^n i \cdot 3^{-i} \end{aligned}$$



# Examples

- Solve  $T(n) = 3T(n-1) + n$  with  $T(0) = 10$

$$\begin{aligned} T(n) &= 10 \cdot 3^n + \sum_{i=1}^n 3^{n-i} \cdot i \\ &= 10 \cdot 3^n + 3^n \sum_{i=1}^n i \cdot 3^{-i} \end{aligned}$$

**Theorem.** For any real number  $x \neq 1$ ,

$$\sum_{i=1}^n ix^i = \frac{nx^{n+2} - (n+1)x^{n+1} + x}{(1-x)^2}.$$



# Examples

- Solve  $T(n) = 3T(n-1) + n$  with  $T(0) = 10$

$$T(n) = 10 \cdot 3^n + \sum_{i=1}^n 3^{n-i} \cdot i$$

$$= 10 \cdot 3^n + 3^n \sum_{i=1}^n i \cdot 3^{-i}$$

$$= 10 \cdot 3^n + 3^n \left( -\frac{3}{2}(n+1)3^{-(n+1)} - \frac{3}{4}3^{-(n+1)} + \frac{3}{4} \right)$$

$$= \frac{43}{4}3^n - \frac{n+1}{2} - \frac{1}{4}.$$



# Growth Rates of Solutions to Recurrences

- Divide and conquer algorithms
- Iterating recurrences
- Three different behaviors



# Divide and conquer algorithms

- We just analyzed recurrences of the form

$$T(n) = \begin{cases} b & \text{if } n = 0 \\ r \cdot T(n-1) + a & \text{if } n > 0 \end{cases}$$



# Divide and conquer algorithms

- We just analyzed recurrences of the form

$$T(n) = \begin{cases} b & \text{if } n = 0 \\ r \cdot T(n-1) + a & \text{if } n > 0 \end{cases}$$

These corresponded to the analysis of recursive algorithms in which a problem of size  $n$  is solved by recursively solving a problem of size  $n - 1$ .



# Divide and conquer algorithms

- We just analyzed recurrences of the form

$$T(n) = \begin{cases} b & \text{if } n = 0 \\ r \cdot T(n-1) + a & \text{if } n > 0 \end{cases}$$

These corresponded to the analysis of recursive algorithms in which a problem of size  $n$  is solved by recursively solving a problem of size  $n - 1$ .

$$T(n) = r^n b + a \frac{1 - r^n}{1 - r}$$





# Divide and conquer algorithms

- We just analyzed recurrences of the form

$$T(n) = \begin{cases} b & \text{if } n = 0 \\ r \cdot T(n-1) + a & \text{if } n > 0 \end{cases}$$

These corresponded to the analysis of recursive algorithms in which a problem of size  $n$  is solved by recursively solving a problem of size  $n-1$ .

$$T(n) = r^n b + a \frac{1 - r^n}{1 - r}$$

We will now look at recurrences of the form

$$T(n) = \begin{cases} \text{something given} & \text{if } n \leq n_0 \\ r \cdot T(n/m) + a & \text{if } n > n_0 \end{cases}$$



# Binary Search

- Someone has chosen a number  $x$  between 1 and  $n$ .  
We need to discover  $x$ .



# Binary Search

- Someone has chosen a number  $x$  between 1 and  $n$ . We need to discover  $x$ .

We are only allowed to ask **two types of questions**:



# Binary Search

- Someone has chosen a number  $x$  between 1 and  $n$ . We need to discover  $x$ .

We are only allowed to ask two types of questions:

- ◇ Is  $x$  greater than  $k$ ?
- ◇ Is  $x$  equal to  $k$ ?



# Binary Search

- Someone has chosen a number  $x$  between 1 and  $n$ . We need to discover  $x$ .

We are only allowed to ask two types of questions:

- ◇ Is  $x$  greater than  $k$ ?
- ◇ Is  $x$  equal to  $k$ ?

Our strategy will be to always ask greater than questions, at each step halving our search range, until the range only contains one number, when we ask a final equal to question.



# Binary Search Example

1

32

48

64

11 - 1



# Binary Search Example

1

32


48

64

Is  $x > 32$ ?



# Binary Search Example




Is  $x > 32$ ?

Answer: Yes





# Binary Search Example




Is  $x > 32$ ?

Answer: Yes

Is  $x > 48$ ?



# Binary Search Example



Is  $x > 32$ ?


Answer: Yes

Is  $x > 48$ ?

Answer: No



# Binary Search Example



Is  $x > 32$ ?

Answer: Yes


Is  $x > 48$ ?

Answer: No

Is  $x > 40$ ?



# Binary Search Example



Is  $x > 32$ ?

Answer: Yes

Is  $x > 48$ ?


Answer: No

Is  $x > 40$ ?

Answer: No



# Binary Search Example



Is  $x > 32$ ? Answer: Yes


Is  $x > 48$ ? Answer: No

Is  $x > 40$ ? Answer: No

Is  $x > 36$ ?



# Binary Search Example



Is  $x > 32$ ? Answer: Yes


Is  $x > 48$ ? Answer: No

Is  $x > 40$ ? Answer: No

Is  $x > 36$ ? Answer: No



# Binary Search Example



Is  $x > 32$ ? Answer: Yes

Is  $x > 48$ ? Answer: No


Is  $x > 40$ ? Answer: No

Is  $x > 36$ ? Answer: No

Is  $x > 34$ ?



# Binary Search Example



Is  $x > 32$ ?      Answer: Yes

Is  $x > 48$ ?      Answer: No

Is  $x > 40$ ?      Answer: No


Is  $x > 36$ ?      Answer: No

Is  $x > 34$ ?      Answer: Yes





# Binary Search Example



Is  $x > 32$ ? Answer: Yes

Is  $x > 48$ ? Answer: No

Is  $x > 40$ ? Answer: No


Is  $x > 36$ ? Answer: No

Is  $x > 34$ ? Answer: Yes

Is  $x > 35$ ?



# Binary Search Example



Is  $x > 32$ ? Answer: Yes

Is  $x > 48$ ? Answer: No

Is  $x > 40$ ? Answer: No


Is  $x > 36$ ? Answer: No

Is  $x > 34$ ? Answer: Yes

Is  $x > 35$ ? Answer: No



# Binary Search Example



Is  $x > 32$ ? Answer: Yes

Is  $x > 48$ ? Answer: No

Is  $x > 40$ ? Answer: No

Is  $x > 36$ ? Answer: No


Is  $x > 34$ ? Answer: Yes

Is  $x > 35$ ? Answer: No

Is  $x = 35$ ?



# Binary Search Example



Is  $x > 32$ ? Answer: Yes

Is  $x > 48$ ? Answer: No

Is  $x > 40$ ? Answer: No

Is  $x > 36$ ? Answer: No

Is  $x > 34$ ? Answer: Yes

Is  $x > 35$ ? Answer: No

Is  $x = 35$ ? Answer: BINGO!



# Binary Search Example

- **Method**: Each guess **reduces** the problem to one in which the range is only **half** as big.



# Binary Search Example

- **Method**: Each guess **reduces** the problem to one in which the range is only **half** as big.

This **divides** the original problem into one that is only half as big; we can now (**recursively**) **conquer** this smaller problem.



# Binary Search Example

- **Method:** Each guess **reduces** the problem to one in which the range is only **half** as big.

This **divides** the original problem into one that is only half as big; we can now (**recursively**) **conquer** this smaller problem.

**Note:** When  $n$  is a power of 2,  $T(n)$ , the number of questions in a binary search on  $[1, n]$ , satisfies

$$T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$



# Binary Search Example

- **Method:** Each guess **reduces** the problem to one in which the range is only **half** as big.

This **divides** the original problem into one that is only half as big; we can now (**recursively**) **conquer** this smaller problem.

**Note:** When  $n$  is a power of 2,  $T(n)$ , the number of questions in a binary search on  $[1, n]$ , satisfies

$$T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$

This can also be proved **inductively**, similar to the tower of Hanoi recurrence.





# Binary Search Example

- $T(n)$ : number of questions in a binary search on  $[1, n]$



# Binary Search Example

- $T(n)$ : number of questions in a binary search on  $[1, n]$

Assume:  $n$  is a power of 2.      Give recurrence for  $T(n)$



# Binary Search Example

- $T(n)$ : number of questions in a binary search on  $[1, n]$

Assume:  $n$  is a power of 2.      Give recurrence for  $T(n)$

$$T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$



# Binary Search Example

- $T(n)$ : number of questions in a binary search on  $[1, n]$

Assume:  $n$  is a power of 2.      Give recurrence for  $T(n)$

$$T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$

Number of questions needed for **binary search** on  $n$  items is:



# Binary Search Example

- $T(n)$ : number of questions in a binary search on  $[1, n]$

Assume:  $n$  is a power of 2.      Give recurrence for  $T(n)$

$$T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$

Number of questions needed for **binary search** on  $n$  items is:

**first step**

+

**time to perform binary search on the remaining  $n/2$  items**



# Binary Search Example

- $T(n)$ : number of questions in a binary search on  $[1, n]$

Assume:  $n$  is a power of 2.      Give recurrence for  $T(n)$

$$T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$

Number of questions needed for **binary search** on  $n$  items is:

**first step**

+

**time to perform binary search on the remaining  $n/2$  items**

Base case (1 item):  $T(1) = 1$  to ask: “**Is the number  $k$ ?**”



# Binary Search Example

$$(*) \quad T(n) = \begin{cases} C_1 & \text{if } n = 1 \\ T(\lceil n/2 \rceil) + C_2 & \text{if } n \geq 2 \end{cases}$$

For simplicity, we will (usually) assume that  $n$  is a power of 2 (or sometimes 3 or 4) and also often that constants such as  $C_1, C_2$  are 1. This will let us replace a recurrence such as (\*) by one such as (\*\*).



# Binary Search Example

$$(*) \quad T(n) = \begin{cases} C_1 & \text{if } n = 1 \\ T(\lceil n/2 \rceil) + C_2 & \text{if } n \geq 2 \end{cases}$$

For simplicity, we will (usually) assume that  $n$  is a power of 2 (or sometimes 3 or 4) and also often that constants such as  $C_1, C_2$  are 1. This will let us replace a recurrence such as (\*) by one such as (\*\*).

$$(**) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$





# Binary Search Example

$$(*) \quad T(n) = \begin{cases} C_1 & \text{if } n = 1 \\ T(\lceil n/2 \rceil) + C_2 & \text{if } n \geq 2 \end{cases}$$

For simplicity, we will (usually) assume that  $n$  is a power of 2 (or sometimes 3 or 4) and also often that constants such as  $C_1, C_2$  are 1. This will let us replace a recurrence such as (\*) by one such as (\*\*).

$$(**) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$

In practice, the solution of (\*) will be very close to that of (\*\*) (this can be proved mathematically). Hence, we can restrict attention to (\*\*).



# Growth Rates of Solutions to Recurrences

- Divide and conquer algorithms
- Iterating recurrences
- Three different behaviors



# Iterating Recurrences: Example 1

■

$$(*) \quad T(n) = \begin{cases} T(1) & \text{if } n = 1 \\ 2T(n/2) + n & \text{if } n \geq 2 \end{cases}$$



# Iterating Recurrences: Example 1

■

$$(*) \quad T(n) = \begin{cases} T(1) & \text{if } n = 1 \\ 2T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

This corresponds to solving a problem of size  $n$ , by

- (i) solving 2 subproblems of size  $n/2$  and
- (ii) doing  $n$  units of additional work

or using  $T(1)$  work for “bottom” case of  $n = 1$



# Iterating Recurrences: Example 1

$$(*) \quad T(n) = \begin{cases} T(1) & \text{if } n = 1 \\ 2T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

This corresponds to solving a problem of size  $n$ , by

- (i) solving 2 subproblems of size  $n/2$  and
- (ii) doing  $n$  units of additional work

or using  $T(1)$  work for “bottom” case of  $n = 1$

In the course “Analysis of Algorithms”, this is exactly how **Mergesort** works.



# Iterating Recurrences: Example 1

$$(*) \quad T(n) = \begin{cases} T(1) & \text{if } n = 1 \\ 2T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

This corresponds to solving a problem of size  $n$ , by

- (i) solving 2 subproblems of size  $n/2$  and
- (ii) doing  $n$  units of additional work

or using  $T(1)$  work for “bottom” case of  $n = 1$

In the course “Analysis of Algorithms”, this is exactly how **Mergesort** works.

We now see how to solve  $(*)$  by algebraically iterating the recurrence.



# Iterating Recurrences: Example 1

- Algebraically iterating the recurrence

$$T(n) = 2T\left(\frac{n}{2}\right) + n$$



# Iterating Recurrences: Example 1

- Algebraically iterating the recurrence

Assume that  $n$  is a power of 2

$$T(n) = 2T\left(\frac{n}{2}\right) + n = 2\left(2T\left(\frac{n}{4}\right) + \frac{n}{2}\right) + n$$





# Iterating Recurrences: Example 1

- Algebraically iterating the recurrence

Assume that  $n$  is a power of 2

$$\begin{aligned} T(n) &= 2T\left(\frac{n}{2}\right) + n &= 2\left(2T\left(\frac{n}{4}\right) + \frac{n}{2}\right) + n \\ &= 4T\left(\frac{n}{4}\right) + 2n &= 4\left(2T\left(\frac{n}{8}\right) + \frac{n}{4}\right) + 2n \end{aligned}$$



# Iterating Recurrences: Example 1

- Algebraically iterating the recurrence

Assume that  $n$  is a power of 2

$$\begin{aligned} T(n) &= 2T\left(\frac{n}{2}\right) + n &= 2\left(2T\left(\frac{n}{4}\right) + \frac{n}{2}\right) + n \\ &= 4T\left(\frac{n}{4}\right) + 2n &= 4\left(2T\left(\frac{n}{8}\right) + \frac{n}{4}\right) + 2n \\ &= 8T\left(\frac{n}{8}\right) + 3n \end{aligned}$$



# Iterating Recurrences: Example 1

- Algebraically iterating the recurrence

Assume that  $n$  is a power of 2

$$\begin{aligned}T(n) &= 2T\left(\frac{n}{2}\right) + n &= 2\left(2T\left(\frac{n}{4}\right) + \frac{n}{2}\right) + n \\&= 4T\left(\frac{n}{4}\right) + 2n &= 4\left(2T\left(\frac{n}{8}\right) + \frac{n}{4}\right) + 2n \\&= 8T\left(\frac{n}{8}\right) + 3n \\&\quad \vdots \quad \vdots \\&= 2^i T\left(\frac{n}{2^i}\right) + in\end{aligned}$$



# Iterating Recurrences: Example 1

- Algebraically iterating the recurrence

Assume that  $n$  is a power of 2

$$T(n) = 2T\left(\frac{n}{2}\right) + n = 2\left(2T\left(\frac{n}{4}\right) + \frac{n}{2}\right) + n$$

$$= 4T\left(\frac{n}{4}\right) + 2n = 4\left(2T\left(\frac{n}{8}\right) + \frac{n}{4}\right) + 2n$$

$$= 8T\left(\frac{n}{8}\right) + 3n$$

$$\begin{array}{c} \vdots \\ \vdots \\ = 2^i T\left(\frac{n}{2^i}\right) + in \end{array}$$

$$\begin{array}{c} \vdots \\ \vdots \\ = 2^{\log_2 n} T\left(\frac{n}{2^{\log_2 n}}\right) + (\log_2 n)n \end{array}$$

End when  $i = \log_2 n$



# Iterating Recurrences: Example 1

- Algebraically iterating the recurrence

Assume that  $n$  is a power of 2

$$T(n) = 2T\left(\frac{n}{2}\right) + n = 2\left(2T\left(\frac{n}{4}\right) + \frac{n}{2}\right) + n$$

$$= 4T\left(\frac{n}{4}\right) + 2n = 4\left(2T\left(\frac{n}{8}\right) + \frac{n}{4}\right) + 2n$$

$$= 8T\left(\frac{n}{8}\right) + 3n$$

$$\vdots \quad \vdots$$
$$= 2^i T\left(\frac{n}{2^i}\right) + in$$

$$\vdots \quad \vdots$$
$$= 2^{\log_2 n} T\left(\frac{n}{2^{\log_2 n}}\right) + (\log_2 n)n$$

$$= nT(1) + n\log_2 n$$

End when  $i = \log_2 n$



# Iterating Recurrences: Example 1

- We just iterated the recurrence to derive that the solution to

$$(*) \quad T(n) = \begin{cases} T(1) & \text{if } n = 1 \\ 2T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

is  $nT(1) + n \log_2 n$ .



# Iterating Recurrences: Example 1

- We just iterated the recurrence to derive that the solution to

$$(*) \quad T(n) = \begin{cases} T(1) & \text{if } n = 1 \\ 2T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

is  $nT(1) + n \log_2 n$ .

**Note:** Technically, we still need to use **induction** to prove that our solution is correct. Practically, we **never** explicitly perform this step, since it is obvious how the induction would work.



# Iterating Recurrences: Example 2

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$





# Iterating Recurrences: Example 2

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$

$$T(n) = T\left(\frac{n}{2}\right) + 1$$



# Iterating Recurrences: Example 2



$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$

$$T(n) = T\left(\frac{n}{2}\right) + 1 = \left(T\left(\frac{n}{2^2}\right) + 1\right) + 1$$



# Iterating Recurrences: Example 2



$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$

$$\begin{aligned} T(n) &= T\left(\frac{n}{2}\right) + 1 &= \left(T\left(\frac{n}{2^2}\right) + 1\right) + 1 \\ &= T\left(\frac{n}{2^2}\right) + 2 \end{aligned}$$



# Iterating Recurrences: Example 2



$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$

$$\begin{aligned} T(n) &= T\left(\frac{n}{2}\right) + 1 &= \left(T\left(\frac{n}{2^2}\right) + 1\right) + 1 \\ &= T\left(\frac{n}{2^2}\right) + 2 &= \left(T\left(\frac{n}{2^3}\right) + 1\right) + 2 \end{aligned}$$



# Iterating Recurrences: Example 2



$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$

$$\begin{aligned} T(n) &= T\left(\frac{n}{2}\right) + 1 &= \left(T\left(\frac{n}{2^2}\right) + 1\right) + 1 \\ &= T\left(\frac{n}{2^2}\right) + 2 &= \left(T\left(\frac{n}{2^3}\right) + 1\right) + 2 \\ &= T\left(\frac{n}{2^3}\right) + 3 \end{aligned}$$



# Iterating Recurrences: Example 2



$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$

$$T(n) = T\left(\frac{n}{2}\right) + 1 = \left(T\left(\frac{n}{2^2}\right) + 1\right) + 1$$

$$= T\left(\frac{n}{2^2}\right) + 2 = \left(T\left(\frac{n}{2^3}\right) + 1\right) + 2$$

$$= T\left(\frac{n}{2^3}\right) + 3$$

$$\vdots \quad \vdots$$

$$= T\left(\frac{n}{2^i}\right) + i$$



# Iterating Recurrences: Example 2

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$

$$T(n) = T\left(\frac{n}{2}\right) + 1 = \left(T\left(\frac{n}{2^2}\right) + 1\right) + 1$$

$$= T\left(\frac{n}{2^2}\right) + 2 = \left(T\left(\frac{n}{2^3}\right) + 1\right) + 2$$

$$= T\left(\frac{n}{2^3}\right) + 3$$

$$\begin{array}{c} \vdots \\ \vdots \\ = T\left(\frac{n}{2^i}\right) + i \end{array}$$

$$\begin{array}{c} \vdots \\ \vdots \\ = T\left(\frac{n}{2^{\log_2 n}}\right) + \log_2 n \end{array}$$



# Iterating Recurrences: Example 2

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$

$$T(n) = T\left(\frac{n}{2}\right) + 1 = \left(T\left(\frac{n}{2^2}\right) + 1\right) + 1$$

$$= T\left(\frac{n}{2^2}\right) + 2 = \left(T\left(\frac{n}{2^3}\right) + 1\right) + 2$$

$$= T\left(\frac{n}{2^3}\right) + 3$$

$$\begin{array}{c} \vdots \\ \vdots \\ = T\left(\frac{n}{2^i}\right) + i \end{array}$$

$$\begin{array}{c} \vdots \\ \vdots \\ = T\left(\frac{n}{2^{\log_2 n}}\right) + \log_2 n = 1 + \log_2 n \end{array}$$





# Iterating Recurrences: Example 3

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + n & \text{if } n \geq 2 \end{cases}$$



# Iterating Recurrences: Example 3



$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$$\begin{aligned} T(n) &= T\left(\frac{n}{2}\right) + n \\ &= T\left(\frac{n}{2^2}\right) + \frac{n}{2} + n \end{aligned}$$



# Iterating Recurrences: Example 3



$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$$T(n) = T\left(\frac{n}{2}\right) + n$$

$$= T\left(\frac{n}{2^2}\right) + \frac{n}{2} + n$$

$$= T\left(\frac{n}{2^3}\right) + \frac{n}{2^2} + \frac{n}{2} + n$$



# Iterating Recurrences: Example 3

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$$\begin{aligned} T(n) &= T\left(\frac{n}{2}\right) + n \\ &= T\left(\frac{n}{2^2}\right) + \frac{n}{2} + n \\ &= T\left(\frac{n}{2^3}\right) + \frac{n}{2^2} + \frac{n}{2} + n \\ &\quad \vdots \\ &= T\left(\frac{n}{2^i}\right) + \frac{n}{2^{i-1}} + \cdots + \frac{n}{2^2} + \frac{n}{2} + n \end{aligned}$$



# Iterating Recurrences: Example 3

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$$T(n) = T\left(\frac{n}{2}\right) + n$$

$$= T\left(\frac{n}{2^2}\right) + \frac{n}{2} + n$$

$$= T\left(\frac{n}{2^3}\right) + \frac{n}{2^2} + \frac{n}{2} + n$$

$$\vdots \quad \vdots$$

$$= T\left(\frac{n}{2^i}\right) + \frac{n}{2^{i-1}} + \cdots + \frac{n}{2^2} + \frac{n}{2} + n$$

$$\vdots \quad \vdots$$

$$= T\left(\frac{n}{2^{\log_2 n}}\right) + \frac{n}{2^{\log_2 n - 1}} + \cdots + \frac{n}{2^2} + \frac{n}{2} + n$$



# Iterating Recurrences: Example 3

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$$T(n) = T\left(\frac{n}{2}\right) + n$$

$$= T\left(\frac{n}{2^2}\right) + \frac{n}{2} + n$$

$$= T\left(\frac{n}{2^3}\right) + \frac{n}{2^2} + \frac{n}{2} + n$$

$$\vdots \quad \vdots$$

$$= T\left(\frac{n}{2^i}\right) + \frac{n}{2^{i-1}} + \cdots + \frac{n}{2^2} + \frac{n}{2} + n$$

$$\vdots \quad \vdots$$

$$= T\left(\frac{n}{2^{\log_2 n}}\right) + \frac{n}{2^{\log_2 n - 1}} + \cdots + \frac{n}{2^2} + \frac{n}{2} + n$$

$$= 1 + 2 + 2^2 + \cdots + \frac{n}{2^2} + \frac{n}{2} + n$$



# Iterating Recurrences: Example 3

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$$T(n) = T\left(\frac{n}{2}\right) + n$$

$$= T\left(\frac{n}{2^2}\right) + \frac{n}{2} + n$$

$$= T\left(\frac{n}{2^3}\right) + \frac{n}{2^2} + \frac{n}{2} + n$$

$$\vdots \quad \vdots$$

$$= T\left(\frac{n}{2^i}\right) + \frac{n}{2^{i-1}} + \cdots + \frac{n}{2^2} + \frac{n}{2} + n$$

$$\vdots \quad \vdots$$

$$= T\left(\frac{n}{2^{\log_2 n}}\right) + \frac{n}{2^{\log_2 n - 1}} + \cdots + \frac{n}{2^2} + \frac{n}{2} + n$$

$$= 1 + 2 + 2^2 + \cdots + \frac{n}{2^2} + \frac{n}{2} + n = \Theta(n)$$



# Iterating Recurrences: Example 4

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n < 3 \\ 3T(n/3) + n & \text{if } n \geq 3 \end{cases}$$





# Iterating Recurrences: Example 4

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n < 3 \\ 3T(n/3) + n & \text{if } n \geq 3 \end{cases}$$

$$T(n) = 3T\left(\frac{n}{3}\right) + n$$



# Iterating Recurrences: Example 4

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n < 3 \\ 3T(n/3) + n & \text{if } n \geq 3 \end{cases}$$

$$T(n) = 3T\left(\frac{n}{3}\right) + n = 3\left(3T\left(\frac{n}{3^2}\right) + \frac{n}{3}\right) + n$$



# Iterating Recurrences: Example 4

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n < 3 \\ 3T(n/3) + n & \text{if } n \geq 3 \end{cases}$$

$$\begin{aligned} T(n) &= 3T\left(\frac{n}{3}\right) + n &= 3\left(3T\left(\frac{n}{3^2}\right) + \frac{n}{3}\right) + n \\ &= 3^2 T\left(\frac{n}{3^2}\right) + 2n \end{aligned}$$



# Iterating Recurrences: Example 4

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n < 3 \\ 3T(n/3) + n & \text{if } n \geq 3 \end{cases}$$

$$\begin{aligned} T(n) &= 3T\left(\frac{n}{3}\right) + n &= 3\left(3T\left(\frac{n}{3^2}\right) + \frac{n}{3}\right) + n \\ &= 3^2 T\left(\frac{n}{3^2}\right) + 2n &= 3^2\left(3T\left(\frac{n}{3^3}\right) + \frac{n}{3^2}\right) + 2n \end{aligned}$$



# Iterating Recurrences: Example 4

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n < 3 \\ 3T(n/3) + n & \text{if } n \geq 3 \end{cases}$$

$$\begin{aligned} T(n) &= 3T\left(\frac{n}{3}\right) + n &= 3\left(3T\left(\frac{n}{3^2}\right) + \frac{n}{3}\right) + n \\ &= 3^2 T\left(\frac{n}{3^2}\right) + 2n &= 3^2\left(3T\left(\frac{n}{3^3}\right) + \frac{n}{3^2}\right) + 2n \\ &= 3^3 T\left(\frac{n}{3^3}\right) + 3n \end{aligned}$$



# Iterating Recurrences: Example 4

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n < 3 \\ 3T(n/3) + n & \text{if } n \geq 3 \end{cases}$$

$$\begin{aligned} T(n) &= 3T\left(\frac{n}{3}\right) + n &= 3\left(3T\left(\frac{n}{3^2}\right) + \frac{n}{3}\right) + n \\ &= 3^2 T\left(\frac{n}{3^2}\right) + 2n &= 3^2\left(3T\left(\frac{n}{3^3}\right) + \frac{n}{3^2}\right) + 2n \\ &= 3^3 T\left(\frac{n}{3^3}\right) + 3n \\ &\quad \vdots \quad \vdots \\ &= 3^i T\left(\frac{n}{3^i}\right) + in \end{aligned}$$



# Iterating Recurrences: Example 4

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n < 3 \\ 3T(n/3) + n & \text{if } n \geq 3 \end{cases}$$

$$\begin{aligned} T(n) &= 3T\left(\frac{n}{3}\right) + n &= 3\left(3T\left(\frac{n}{3^2}\right) + \frac{n}{3}\right) + n \\ &= 3^2 T\left(\frac{n}{3^2}\right) + 2n &= 3^2\left(3T\left(\frac{n}{3^3}\right) + \frac{n}{3^2}\right) + 2n \\ &= 3^3 T\left(\frac{n}{3^3}\right) + 3n \\ &\quad \vdots \quad \vdots \\ &= 3^i T\left(\frac{n}{3^i}\right) + in \\ &\quad \vdots \quad \vdots \\ &= 3^{\log_3 n} T\left(\frac{n}{3^{\log_3 n}}\right) + n \log_3 n \end{aligned}$$



# Iterating Recurrences: Example 4

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n < 3 \\ 3T(n/3) + n & \text{if } n \geq 3 \end{cases}$$

$$\begin{aligned} T(n) &= 3T\left(\frac{n}{3}\right) + n &= 3\left(3T\left(\frac{n}{3^2}\right) + \frac{n}{3}\right) + n \\ &= 3^2 T\left(\frac{n}{3^2}\right) + 2n &= 3^2\left(3T\left(\frac{n}{3^3}\right) + \frac{n}{3^2}\right) + 2n \\ &= 3^3 T\left(\frac{n}{3^3}\right) + 3n \\ &\quad \vdots \quad \vdots \\ &= 3^i T\left(\frac{n}{3^i}\right) + in \\ &\quad \vdots \quad \vdots \\ &= 3^{\log_3 n} T\left(\frac{n}{3^{\log_3 n}}\right) + n \log_3 n = n + n \log_3 n \end{aligned}$$





# Iterating Recurrences: Example 5

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ 4T(n/2) + n & \text{if } n \geq 2 \end{cases}$$



# Iterating Recurrences: Example 5

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ 4T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$$T(n) = 4T\left(\frac{n}{2}\right) + n$$



# Iterating Recurrences: Example 5

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ 4T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$$T(n) = 4T\left(\frac{n}{2}\right) + n = 4\left(4T\left(\frac{n}{2^2}\right) + \frac{n}{2}\right) + n$$



# Iterating Recurrences: Example 5

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ 4T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$$\begin{aligned} T(n) &= 4T\left(\frac{n}{2}\right) + n &= 4\left(4T\left(\frac{n}{2^2}\right) + \frac{n}{2}\right) + n \\ &= 4^2 T\left(\frac{n}{2^2}\right) + \frac{4}{2}n + n \end{aligned}$$



# Iterating Recurrences: Example 5

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ 4T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$$\begin{aligned} T(n) &= 4T\left(\frac{n}{2}\right) + n &= 4\left(4T\left(\frac{n}{2^2}\right) + \frac{n}{2}\right) + n \\ &= 4^2 T\left(\frac{n}{2^2}\right) + \frac{4}{2}n + n &= 4^2\left(4T\left(\frac{n}{2^3}\right) + \frac{n}{2^2}\right) + \frac{4}{2}n + n \end{aligned}$$



# Iterating Recurrences: Example 5

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ 4T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$$\begin{aligned} T(n) &= 4T\left(\frac{n}{2}\right) + n &&= 4\left(4T\left(\frac{n}{2^2}\right) + \frac{n}{2}\right) + n \\ &= 4^2 T\left(\frac{n}{2^2}\right) + \frac{4}{2}n + n &&= 4^2\left(4T\left(\frac{n}{2^3}\right) + \frac{n}{2^2}\right) + \frac{4}{2}n + n \\ &= 4^3 T\left(\frac{n}{2^3}\right) + \frac{4^2}{2^2}n + \frac{4}{2}n + n \end{aligned}$$



# Iterating Recurrences: Example 5

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ 4T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$$\begin{aligned} T(n) &= 4T\left(\frac{n}{2}\right) + n &&= 4\left(4T\left(\frac{n}{2^2}\right) + \frac{n}{2}\right) + n \\ &= 4^2 T\left(\frac{n}{2^2}\right) + \frac{4}{2}n + n &&= 4^2\left(4T\left(\frac{n}{2^3}\right) + \frac{n}{2^2}\right) + \frac{4}{2}n + n \\ &= 4^3 T\left(\frac{n}{2^3}\right) + \frac{4^2}{2^2}n + \frac{4}{2}n + n \\ &\quad \vdots \quad \vdots \\ &= 4^i T\left(\frac{n}{2^i}\right) + \frac{4^{i-1}}{2^{i-1}}n + \cdots + \frac{4^2}{2^2}n + n \end{aligned}$$



# Iterating Recurrences: Example 5

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ 4T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$$\begin{aligned} T(n) &= 4T\left(\frac{n}{2}\right) + n &&= 4\left(4T\left(\frac{n}{2^2}\right) + \frac{n}{2}\right) + n \\ &= 4^2 T\left(\frac{n}{2^2}\right) + \frac{4}{2}n + n &&= 4^2\left(4T\left(\frac{n}{2^3}\right) + \frac{n}{2^2}\right) + \frac{4}{2}n + n \\ &= 4^3 T\left(\frac{n}{2^3}\right) + \frac{4^2}{2^2}n + \frac{4}{2}n + n \\ &\quad \vdots \quad \vdots \\ &= 4^i T\left(\frac{n}{2^i}\right) + \frac{4^{i-1}}{2^{i-1}}n + \cdots + \frac{4^2}{2^2}n + n \\ &\quad \vdots \quad \vdots \\ &= 4^{\log_2 n} T\left(\frac{n}{2^{\log_2 n}}\right) + \frac{4^{\log_2 n - 1}}{2^{\log_2 n - 1}}n + \cdots + \frac{4}{2}n + n \end{aligned}$$





# Iterating Recurrences: Example 5

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ 4T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$$\begin{aligned} T(n) &= 4T\left(\frac{n}{2}\right) + n &&= 4\left(4T\left(\frac{n}{2^2}\right) + \frac{n}{2}\right) + n \\ &= 4^2 T\left(\frac{n}{2^2}\right) + \frac{4}{2}n + n &&= 4^2\left(4T\left(\frac{n}{2^3}\right) + \frac{n}{2^2}\right) + \frac{4}{2}n + n \\ &= 4^3 T\left(\frac{n}{2^3}\right) + \frac{4^2}{2^2}n + \frac{4}{2}n + n \\ &\quad \vdots \quad \vdots \\ &= 4^i T\left(\frac{n}{2^i}\right) + \frac{4^{i-1}}{2^{i-1}}n + \cdots + \frac{4^2}{2^2}n + n \\ &\quad \vdots \quad \vdots \\ &= 4^{\log_2 n} T\left(\frac{n}{2^{\log_2 n}}\right) + \frac{4^{\log_2 n - 1}}{2^{\log_2 n - 1}}n + \cdots + \frac{4}{2}n + n \\ &= 2n^2 - n \end{aligned}$$



# Growth Rates of Solutions to Recurrences

- Divide and conquer algorithms
- Iteration recurrences
- Three different behaviors



# Three Different Behaviors

- Compare the iteration for the recurrences

$$T(n) = 2T(n/2) + n$$

$$T(n) = T(n/2) + n$$

$$T(n) = 4T(n/2) + n$$



# Three Different Behaviors

- Compare the iteration for the recurrences

$$T(n) = 2T(n/2) + n$$

$$T(n) = T(n/2) + n$$

$$T(n) = 4T(n/2) + n$$

- ◇ all three recurrences iterate  $\log_2 n$  times
- ◇ in each case, size of subproblem in next iteration is **half** the size in the preceding iteration level



# Three Different Behaviors

- **Theorem** Suppose that we have a recurrence of the form

$$T(n) = aT(n/2) + n,$$

where  $a$  is a positive integer and  $T(1)$  is nonnegative. Then we have the following **big  $\Theta$**  bounds on the solution:

1. If  $a < 2$ , then  $T(n) = \Theta(n)$ .
2. If  $a = 2$ , then  $T(n) = \Theta(n \log n)$ .
3. If  $a > 2$ , then  $T(n) = \Theta(n^{\log_2 a})$



# Three Different Behaviors

- **Theorem** Suppose that we have a recurrence of the form

$$T(n) = aT(n/2) + n,$$

where  $a$  is a positive integer and  $T(1)$  is nonnegative. Then we have the following **big  $\Theta$**  bounds on the solution:

1. If  $a < 2$ , then  $T(n) = \Theta(n)$ .
2. If  $a = 2$ , then  $T(n) = \Theta(n \log n)$ .
3. If  $a > 2$ , then  $T(n) = \Theta(n^{\log_2 a})$

## Proof

We already proved Case 1 when  $a = 1$  in Example 3.

(will not prove it for  $1 < a < 2$ )

We already proved Case 2 in Example 1.

We will now prove Case 3.



# Iterating Recurrences

- $T(n) = aT(n/2) + n$ , where  $a > 2$ . Assume that  $n = 2^i$ .



# Iterating Recurrences

- $T(n) = aT(n/2) + n$ , where  $a > 2$ . Assume that  $n = 2^i$ .

Iterating as in Example 5 gives

$$T(n) = a^i T\left(\frac{n}{2^i}\right) + \left(\frac{a^{i-1}}{2^{i-1}} + \frac{a^{i-2}}{2^{i-2}} + \cdots + \frac{a}{2} + 1\right) n$$





# Iterating Recurrences

- $T(n) = aT(n/2) + n$ , where  $a > 2$ . Assume that  $n = 2^i$ .

Iterating as in Example 5 gives

$$T(n) = a^i T\left(\frac{n}{2^i}\right) + \left(\frac{a^{i-1}}{2^{i-1}} + \frac{a^{i-2}}{2^{i-2}} + \cdots + \frac{a}{2} + 1\right) n$$

$$T(n) = a^{\log_2 n} T(1) + n \sum_{i=0}^{\log_2 n - 1} \left(\frac{a}{2}\right)^i$$

Work at  
“bottom”

Iterated  
Work



# Total work

- The total work is

$$T(n) = a^{\log_2 n} T(1) + n \sum_{i=0}^{\log_2 n - 1} \left(\frac{a}{2}\right)^i$$



# Total work

- The total work is

$$T(n) = a^{\log_2 n} T(1) + n \sum_{i=0}^{\log_2 n - 1} \left(\frac{a}{2}\right)^i$$

Since  $a > 2$ , the geometric series is  $\Theta$  of the largest term.

$$n \sum_{i=0}^{\log_2 n - 1} \left(\frac{a}{2}\right)^i = n \frac{1 - (a/2)^{\log_2 n}}{1 - a/2} = n\Theta((a/2)^{\log_2 n - 1})$$



# Total work

- $n$  times the largest term in the geometric series is

$$n \left(\frac{a}{2}\right)^{\log_2 n - 1} = \frac{2}{a} \cdot \frac{n \cdot a^{\log_2 n}}{2^{\log_2 n}} = \frac{2}{a} \cdot \frac{n \cdot a^{\log_2 n}}{n} = \frac{2}{a} \cdot a^{\log_2 n}$$



# Total work

- $n$  times the largest term in the geometric series is

$$n \left(\frac{a}{2}\right)^{\log_2 n - 1} = \frac{2}{a} \cdot \frac{n \cdot a^{\log_2 n}}{2^{\log_2 n}} = \frac{2}{a} \cdot \frac{n \cdot a^{\log_2 n}}{n} = \frac{2}{a} \cdot a^{\log_2 n}$$

Notice that

$$a^{\log_2 n} = (2^{\log_2 a})^{\log_2 n} = (2^{\log_2 n})^{\log_2 a} = n^{\log_2 a}$$



# Total work

- $n$  times the largest term in the geometric series is

$$n \left(\frac{a}{2}\right)^{\log_2 n - 1} = \frac{2}{a} \cdot \frac{n \cdot a^{\log_2 n}}{2^{\log_2 n}} = \frac{2}{a} \cdot \frac{n \cdot a^{\log_2 n}}{n} = \frac{2}{a} \cdot a^{\log_2 n}$$

Notice that

$$a^{\log_2 n} = \left(2^{\log_2 a}\right)^{\log_2 n} = \left(2^{\log_2 n}\right)^{\log_2 a} = n^{\log_2 a}$$

So the total work is

$$a^{\log_2 n} T(1) + n \sum_{i=0}^{\log_2 n - 1} \left(\frac{a}{2}\right)^i$$



# Total work

- $n$  times the largest term in the geometric series is

$$n \left(\frac{a}{2}\right)^{\log_2 n - 1} = \frac{2}{a} \cdot \frac{n \cdot a^{\log_2 n}}{2^{\log_2 n}} = \frac{2}{a} \cdot \frac{n \cdot a^{\log_2 n}}{n} = \frac{2}{a} \cdot a^{\log_2 n}$$

Notice that

$$a^{\log_2 n} = \left(2^{\log_2 a}\right)^{\log_2 n} = \left(2^{\log_2 n}\right)^{\log_2 a} = n^{\log_2 a}$$

So the total work is

$$a^{\log_2 n} T(1) + n \sum_{i=0}^{\log_2 n - 1} \left(\frac{a}{2}\right)^i$$

$$\Theta\left(n^{\log_2 a}\right)$$

$$\Theta\left(n^{\log_2 a}\right)$$



# Example 5 Recap

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ 4T(n/2) + n & \text{if } n \geq 2 \end{cases}$$





## Example 5 Recap

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ 4T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$a = 4$ , so the Theorem says that

$$T(n) = \Theta(n^{\log_2 a}) = \Theta(n^{\log_2 4}) = \Theta(n^2)$$



## Example 5 Recap

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ 4T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$a = 4$ , so the Theorem says that

$$T(n) = \Theta(n^{\log_2 a}) = \Theta(n^{\log_2 4}) = \Theta(n^2)$$

This matches with the exact answer of  $2n^2 - n$ .



# Three Different Behaviors

- **Theorem** Suppose that we have a recurrence of the form

$$T(n) = aT(n/2) + n,$$

where  $a$  is a positive integer and  $T(1)$  is nonnegative. Then we have the following big  $\Theta$  bounds on the solution:

1. If  $a < 2$ , then  $T(n) = \Theta(n)$ .
2. If  $a = 2$ , then  $T(n) = \Theta(n \log n)$ .
3. If  $a > 2$ , then  $T(n) = \Theta(n^{\log_2 a})$



# The Master Theorem

- **Theorem** Suppose that we have a recurrence of the form

$$T(n) = aT(n/b) + cn^d,$$

where  $a$  is a positive integer,  $b \geq 1$ ,  $c, d$  are real numbers with  $c$  positive and  $d$  nonnegative, and  $T(1)$  is nonnegative. Then we have the following **big  $\Theta$**  bounds on the solution:

1. If  $a < b^d$ , then  $T(n) = \Theta(n^d)$ .
2. If  $a = b^d$ , then  $T(n) = \Theta(n^d \log n)$ .
3. If  $a > b^d$ , then  $T(n) = \Theta(n^{\log_b a})$



# Counting

- Assume we have a set of objects with certain properties



# Counting

- Assume we have a set of objects with certain properties  
*Counting* is used to determine the number of these objects.

# Counting

- Assume we have a set of objects with certain properties  
*Counting* is used to determine the number of these objects.


How many different ways are there to choose 2 balls from



# Counting

- Assume we have a set of objects with certain properties  
*Counting* is used to determine the number of these objects.

How many different ways are there to choose 2 balls from






# Counting

- Assume we have a set of objects with certain properties  
*Counting* is used to determine the number of these objects.

How many different ways are there to choose 2 balls from



What about when order counts?


# Counting

- Assume we have a set of objects with certain properties  
*Counting* is used to determine the number of these objects.

How many different ways are there to choose 2 balls from



What about when order counts?



# Counting

- Assume we have a set of objects with certain properties  
*Counting* is used to determine the number of these objects.



# Counting

- Assume we have a set of objects with certain properties  
*Counting* is used to determine the number of these objects.

## Examples

- ◇ the number of steps in a computer program
- ◇ the number of passwords between 6 – 10 characters
- ◇ the number of telephone numbers with 8 digits



# Counting

- Assume we have a set of objects with certain properties  
*Counting* is used to determine the number of these objects.

## Examples

- ◇ the number of steps in a computer program
- ◇ the number of passwords between 6 – 10 characters
- ◇ the number of telephone numbers with 8 digits

Counting may be very hard, not trivial.



# Counting

- Assume we have a set of objects with certain properties  
*Counting* is used to determine the number of these objects.

## Examples

- ◇ the number of steps in a computer program
- ◇ the number of passwords between 6 – 10 characters
- ◇ the number of telephone numbers with 8 digits

Counting may be very hard, not trivial.

- simplify the solution by decomposing the problem



# Basic Counting Rules

- *the Product Rule*

- *the Sum Rule*



# Basic Counting Rules

## ■ *the Product Rule*

- ◇ A count decomposes into a sequence of **dependent** counts  
(each element in the first count is associated with all elements of the second count)

## ■ *the Sum Rule*

- ◇ A count decomposes into a set of **independent** counts  
(elements of counts are alternatives)





# The Product Rule

- A count decomposes into a sequence of **dependent** counts  
(each element in the first count is associated with all elements of the second count)



# The Product Rule

- A count decomposes into a sequence of **dependent** counts (each element in the first count is associated with all elements of the second count)

## Example

In an auditorium, the seats are labeled by a letter and numbers in between 1 to 50 (e.g., A23). What is the total number of seats?



# The Product Rule

- A count decomposes into a sequence of **dependent** counts (each element in the first count is associated with all elements of the second count)

## Example

In an auditorium, the seats are labeled by a letter and numbers in between 1 to 50 (e.g., A23). What is the total number of seats?

We may either list all or use the product rule.

$$26 \times 50 = 1300$$



# The Product Rule

- **Product Rule:** If a count of elements can be broken down into a **sequence of dependent counts** where the first count yields  $n_1$  elements, the second  $n_2$  elements, and  $k$ th count  $n_k$  elements, then the total number of elements is

$$n = n_1 \cdot n_2 \cdot \cdots \cdot n_k$$



# The Product Rule

- **Product Rule:** If a count of elements can be broken down into a **sequence of dependent counts** where the first count yields  $n_1$  elements, the second  $n_2$  elements, and  $k$ th count  $n_k$  elements, then the total number of elements is

$$n = n_1 \cdot n_2 \cdot \dots \cdot n_k$$

## Example

How many different bit strings of length 7 are there?



# The Product Rule

- **Product Rule:** If a count of elements can be broken down into a **sequence of dependent counts** where the first count yields  $n_1$  elements, the second  $n_2$  elements, and  $k$ th count  $n_k$  elements, then the total number of elements is

$$n = n_1 \cdot n_2 \cdot \dots \cdot n_k$$

## Example

How many different bit strings of length 7 are there?

How many different functions are there from a set with  $m$  elements to a set with  $n$  elements?



# The Product Rule

- **Product Rule:** If a count of elements can be broken down into a **sequence of dependent counts** where the first count yields  $n_1$  elements, the second  $n_2$  elements, and  $k$ th count  $n_k$  elements, then the total number of elements is

$$n = n_1 \cdot n_2 \cdot \cdots \cdot n_k$$

## Example

How many different bit strings of length 7 are there?

How many different functions are there from a set with  $m$  elements to a set with  $n$  elements?

How many **one-to-one** functions are there from a set with  $m$  elements to a set with  $n$  elements?



# The Product Rule

- **Product Rule:** If a count of elements can be broken down into a **sequence of dependent counts** where the first count yields  $n_1$  elements, the second  $n_2$  elements, and  $k$ th count  $n_k$  elements, then the total number of elements is

$$n = n_1 \cdot n_2 \cdot \cdots \cdot n_k$$

## Example

How many different bit strings of length 7 are there?

How many different functions are there from a set with  $m$  elements to a set with  $n$  elements?

How many **one-to-one** functions are there from a set with  $m$  elements to a set with  $n$  elements?

How many **onto** functions?





# The Product Rule

- The following loop is a part of program computing the product of two matrices.

```
(1) for i = 1 to r
(2)   for j = 1 to m
(3)     S = 0
(4)     for k = 1 to n
(5)       S = S + A[i,k] * B[k,j]
(6)     C[i,j] = S
```



# The Product Rule

- The following loop is a part of program computing the product of two matrices.

```
(1) for i = 1 to r
(2)   for j = 1 to m
(3)     S = 0
(4)     for k = 1 to n
(5)       S = S + A[i,k] * B[k,j]
(6)     C[i,j] = S
```

How many multiplications (in terms of  $r, m, n$ ) does this program carry out in total among all iterations of line 5?



# The Sum Rule

- A count decomposes into a set of **independent** counts  
(elements of counts are alternatives)



# The Sum Rule

- A count decomposes into a set of **independent** counts  
(elements of counts are alternatives)

## Example

You need to travel from city A to B. You may either fly, take a train, or a bus. There are 12 different flights, 5 different trains and 10 buses. **How many options do you have to get from A to B?**



# The Sum Rule

- A count decomposes into a set of **independent** counts  
(elements of counts are alternatives)

## Example

You need to travel from city A to B. You may either fly, take a train, or a bus. There are 12 different flights, 5 different trains and 10 buses. **How many options do you have to get from A to B?**

We may **use the sum rule.**

$$12 + 5 + 10$$



# The Sum Rule

- **Sum Rule:** If a count of elements can be broken down into a **set of independent counts** where the first count yields  $n_1$  elements, the second  $n_2$  elements, and  $k$ th count  $n_k$  elements, then the total number of elements is

$$n = n_1 + n_2 + \cdots + n_k$$



# The Sum Rule

- The following loop is from [selection sort](#).

```
(1) for i = 1 to n-1
(2)     for j = i+1 to n
(3)         if (A[i] > A[j])
(4)             exchange A[i] and A[j]
```



# The Sum Rule

- The following loop is from [selection sort](#).

```
(1) for i = 1 to n-1
(2)   for j = i+1 to n
(3)     if (A[i] > A[j])
(4)       exchange A[i] and A[j]
```

How many [comparisons](#) (in terms of  $n$ ) does this program carry out in total among all iterations of line 3?





# More Complex Counting

- Typically requires a combination of the sum and product rules.



# More Complex Counting

- Typically requires a **combination** of the sum and product rules.

## Example

Each password is **6 to 8 characters** long, where each character is an lowercase letter or a digit. Each password must contain **at least one digit**. How many possible passwords are there?



# More Complex Counting

- Typically requires a combination of the sum and product rules.

## Example

Each password is 6 to 8 characters long, where each character is an lowercase letter or a digit. Each password must contain at least one digit. How many possible passwords are there?

$$P = P_6 + P_7 + P_8$$



# Tree Diagrams

- A *tree* is a structure that consists of a *root*, *branches* and *leaves*.



# Tree Diagrams

- A *tree* is a structure that consists of a *root*, *branches* and *leaves*.

Can be useful to represent a counting problem and record the choices we made for alternatives. *The count appears on the leaves.*



# Tree Diagrams

- A *tree* is a structure that consists of a **root**, **branches** and **leaves**.

Can be useful to represent a counting problem and record the choices we made for alternatives. **The count appears on the leaves.**

## Example

What is the number of bit strings of length 4 that **do not have two consecutive 1's**?



# Tree Diagrams


- A *tree* is a structure that consists of a *root*, *branches* and *leaves*.

Can be useful to record the choices with the leaves.

Problem and record count appears on

## Example

What is the probability of having two consecutive 1s in a 4-bit sequence?



h 4 that do not



# Tree Diagram


- How many different ways can a “best 3 of 5” playoff occur?





# Tree Diagram

- How many different ways can a “best 3 of 5” playoff occur?



# Pigeonhole Principle

- Assume that there are a set of objects and a set of bins to store them.



# Pigeonhole Principle

- Assume that there are a set of objects and a set of bins to store them.

*The pigeonhole principle* states that if there are more objects than bins then there is at least one bin with more than one object.




# Pigeonhole Principle

- Assume that there are a set of objects and a set of bins to store them.

*The pigeonhole principle* states that if there are more objects than bins then there is at least one bin with more than one object.

**Example:** 7 balls and 5 bins to store them




# Pigeonhole Principle

- Assume that there are a set of objects and a set of bins to store them.

*The pigeonhole principle* states that if there are more objects than bins then there is at least one bin with more than one object.

**Example:** 7 balls and 5 bins to store them



# Pigeonhole Principle

- **Theorem** If there are  $k + 1$  objects and  $k$  bins, then there is at least one bin with two or more objects.



# Pigeonhole Principle

- **Theorem** If there are  $k + 1$  objects and  $k$  bins, then there is at least one bin with two or more objects.

**Proof by contradiction**



# Pigeonhole Principle

- **Theorem** If there are  $k + 1$  objects and  $k$  bins, then there is at least one bin with two or more objects.

## Proof by contradiction

### Example

Assume that there are 367 students. Are there any two people who have the same birthday?

There are 5 bins and 12 objects. Then there must be a bin with at least 3 objects. Why?





# Generalized Pigeonhole Principle

- If  $N$  objects are placed into  $k$  bins, then there is at least one bin containing at least  $\lceil N/k \rceil$  objects.



# Generalized Pigeonhole Principle

- If  $N$  objects are placed into  $k$  bins, then there is at least one bin containing at least  $\lceil N/k \rceil$  objects.

## Example

Assume there are 100 students. How many of them were born in the same month?



# Bijections and Permutations

- A function that is **both one-to-one and onto** is called a *bijection*, or a *one-to-one correspondence*.



# Bijections and Permutations

- A function that is **both one-to-one and onto** is called a *bijection*, or a *one-to-one correspondence*.

How many **bijections** are there?




# Bijections and Permutations

- A function that is **both one-to-one and onto** is called a *bijection*, or a *one-to-one correspondence*.

How many **bijections** are there?

$f : \{a, b, c\} \rightarrow \{1, 2, 3\}$  defined by  $f(a) = 3, f(b) = 2, f(c) = 1$  is a bijection.




# Bijections and Permutations

- A function that is **both one-to-one and onto** is called a *bijection*, or a *one-to-one correspondence*.

How many **bijections** are there?

$f : \{a, b, c\} \rightarrow \{1, 2, 3\}$  defined by  $f(a) = 3, f(b) = 2, f(c) = 1$  is a bijection.



A **bijection** from a set **onto itself** is called a *permutation*.




# Bijections and Permutations

- A function that is **both one-to-one and onto** is called a *bijection*, or a *one-to-one correspondence*.


How many **bijections** are there?

$f : \{a, b, c\} \rightarrow \{1, 2, 3\}$  defined by  $f(a) = 3, f(b) = 2, f(c) = 1$  is a bijection.



A **bijection** from a set **onto itself** is called a *permutation*.

$f : \{1, 2, 3\} \rightarrow \{1, 2, 3\}$  defined by  $f(1) = 3, f(2) = 2, f(3) = 1$  is a bijection.





# Bijections and Permutations

- A function that is **both one-to-one and onto** is called a *bijection*, or a *one-to-one correspondence*.

A **bijection** from a set **onto itself** is called a *permutation*.

In a *bijection*,

**exactly one arrow leaves each item on the left and exactly one arrow arrives at each item on the right.**





# Bijections and Permutations

- A function that is **both one-to-one and onto** is called a *bijection*, or a *one-to-one correspondence*.



A **bijection** from a set **onto itself** is called a *permutation*.

In a *bijection*,

**exactly one arrow leaves each item on the left and exactly one arrow arrives at each item on the right.**

Thus,

**the left and right sides must have the same size.**



# The Bijection Principle

- The following loop is a part of program to determine the number of triangles formed by  $n$  points in the plane.

```
(1) trianglecount = 0
(2)   for i = 1 to n
(3)     for j = i+1 to n
(4)       for k = j+1 to n
(5)         if points i, j, k are not collinear
(6)           trianglecount = trianglecount + 1
```



# The Bijection Principle

- The following loop is a part of program to determine the number of triangles formed by  $n$  points in the plane.

```
(1) trianglecount = 0
```

```
(2)   for  $i = 1$  to  $n$ 
```

```
(3)       for  $j = i+1$  to  $n$ 
```

```
(4)           for  $k = j+1$  to  $n$ 
```

```
(5)               if points  $i, j, k$  are not collinear
```

```
(6)                   trianglecount = trianglecount + 1
```

Among all iterations of line 5, what is the total number of times this line checks three points to see if they are collinear?




# Counting Triangles

- 3 points form a triangle if and only if they are non collinear



# Counting Triangles

- 3 points form a triangle if and only if they are non collinear




50 - 2



# Counting Triangles

- 3 points form a triangle if and only if they are non collinear




1 – 2 – 7: yes

50 - 3



# Counting Triangles

- 3 points form a triangle if and only if they are non collinear




1 – 2 – 7: yes

1 – 2 – 5: yes



# Counting Triangles

- 3 points form a **triangle** if and only if **they are non collinear**



1 – 2 – 7: yes

1 – 2 – 5: yes


1 – 2 – 3: **no**





# Counting Triangles

- 3 points form a triangle if and only if they are non collinear



1 – 2 – 7: yes

1 – 2 – 5: yes


1 – 2 – 3: no

1 – 5 – 6: yes



# Counting Triangles

- 3 points form a triangle if and only if they are non collinear



1 – 2 – 7: yes

1 – 2 – 5: yes

1 – 2 – 3: no


1 – 5 – 6: yes

3 – 4 – 7: yes



# Counting Triangles

- 3 points form a **triangle** if and only if **they are non collinear**



1 – 2 – 7: yes

1 – 2 – 5: yes

1 – 2 – 3: **no**

1 – 5 – 6: yes


3 – 4 – 7: yes

4 – 5 – 6: **no**



# Counting Triangles

- 3 points form a triangle if and only if they are non collinear



1 – 2 – 7: yes

1 – 2 – 5: yes

1 – 2 – 3: no

1 – 5 – 6: yes

3 – 4 – 7: yes

4 – 5 – 6: no

number of triangles: 33



# Counting Triangles

```
(1) trianglecount = 0
(2)   for i = 1 to n
(3)       for j = i+1 to n
(4)           for k = j+1 to n
(5)               if points i, j, k are not collinear
(6)                   trianglecount = trianglecount + 1
```

# Counting Triangles

```
(1) trianglecount = 0
(2)   for i = 1 to n
(3)     for j = i+1 to n
(4)       for k = j+1 to n
(5)         if points i, j, k are not collinear
(6)           trianglecount = trianglecount + 1
```

A loop

# Counting Triangles

```
(1) trianglecount = 0
(2)   for i = 1 to n
(3)     for j = i+1 to n
(4)       for k = j+1 to n
(5)         if points i, j, k are not collinear
(6)           trianglecount = trianglecount + 1
```

A loop embedded in a loop

# Counting Triangles

```
(1) trianglecount = 0
```

```
(2)   for i = 1 to n
```

```
(3)       for j = i+1 to n
```

```
(4)           for k = j+1 to n
```

```
(5)               if points i, j, k are not collinear
```

```
(6)                   trianglecount = trianglecount + 1
```

A loop embedded in a loop embedded in another loop.



# Counting Triangles

```
(1) trianglecount = 0
(2)   for i = 1 to n
(3)     for j = i+1 to n
(4)       for k = j+1 to n
(5)         if points i, j, k are not collinear
(6)           trianglecount = trianglecount + 1
```

A loop embedded in a loop embedded in another loop.

Second loop begins with  $j = i + 1$  and  $j$  increases up to  $n$ .

Third loop begins with  $k = j + 1$  and  $k$  increases up to  $n$ .

# Counting Triangles

```
(1) trianglecount = 0
```

```
(2)   for  $i = 1$  to  $n$ 
```

```
(3)       for  $j = i+1$  to  $n$ 
```

```
(4)           for  $k = j+1$  to  $n$ 
```

```
(5)               if points  $i, j, k$  are not collinear
```

```
(6)                   trianglecount = trianglecount + 1
```

A loop embedded in a loop embedded in another loop.

Second loop begins with  $j = i + 1$  and  $j$  increases up to  $n$ .

Third loop begins with  $k = j + 1$  and  $k$  increases up to  $n$ .

Thus each triple  $i, j, k$  with  $i < j < k$  is examined exactly once.

# Counting Triangles

```
(1) trianglecount = 0
(2)   for i = 1 to n
(3)     for j = i+1 to n
(4)       for k = j+1 to n
(5)         if points i, j, k are not collinear
(6)           trianglecount = trianglecount + 1
```

A loop embedded in a loop embedded in another loop.

Second loop begins with  $j = i + 1$  and  $j$  increases up to  $n$ .

Third loop begins with  $k = j + 1$  and  $k$  increases up to  $n$ .

Thus each triple  $i, j, k$  with  $i < j < k$  is examined exactly once.

For example, if  $n = 4$ , then triples  $(i, j, k)$  used by algorithm are  $(1, 2, 3)$ ,  $(1, 2, 4)$ ,  $(1, 3, 4)$ , and  $(2, 3, 4)$ .

# Counting Triangles

- Want to compute the number of *increasing triples*  $(i, j, k)$  with  $1 \leq i < j < k \leq n$ .

# Counting Triangles

- Want to compute the number of *increasing triples*  $(i, j, k)$  with  $1 \leq i < j < k \leq n$ .

**Claim:** Number of increasing triples is **exactly** the same as number of 3-element subsets from  $\{1, 2, \dots, n\}$

# Counting Triangles

- Want to compute the number of *increasing triples*  $(i, j, k)$  with  $1 \leq i < j < k \leq n$ .

**Claim:** Number of increasing triples is **exactly** the same as number of 3-element subsets from  $\{1, 2, \dots, n\}$

Why? Let  $X =$  set of increasing triples and  
 $Y =$  set of 3-element subsets from  $\{1, 2, \dots, n\}$

# Counting Triangles

- Want to compute the number of *increasing triples*  $(i, j, k)$  with  $1 \leq i < j < k \leq n$ .

**Claim:** Number of increasing triples is **exactly** the same as number of 3-element subsets from  $\{1, 2, \dots, n\}$

Why? Let  $X =$  set of increasing triples and  
 $Y =$  set of 3-element subsets from  $\{1, 2, \dots, n\}$

Define:  $f : X \rightarrow Y$  by  $f((i, j, k)) = \{i, j, k\}$

**Claim:**  $f$  is a **bijection** (why) so  $|X| = |Y|$

# Counting Triangles

- Want to compute the number of *increasing triples*  $(i, j, k)$  with  $1 \leq i < j < k \leq n$ .

**Claim:** Number of increasing triples is **exactly** the same as number of 3-element subsets from  $\{1, 2, \dots, n\}$

Why? Let  $X$  = set of increasing triples and  
 $Y$  = set of 3-element subsets from  $\{1, 2, \dots, n\}$

Define:  $f : X \rightarrow Y$  by  $f((i, j, k)) = \{i, j, k\}$

**Claim:**  $f$  is a **bijection** (why) so  $|X| = |Y|$

$f$  is a bijection because

$f$  is one-to-one

if  $(i, j, k) \neq (i', j', k') \Rightarrow f((i, j, k)) \neq f((i', j', k'))$

$f$  is onto

if  $\gamma$  is a 3-element subset then it can be written as  $\gamma = \{i, j, k\}$

where  $i < j < k$  so  $f((i, j, k)) = \gamma$ .



# Counting Pairs

- The number of  
increasing pairs  $(i, j)$  with  $1 \leq i < j \leq n$   
is the same as the number of  
2-sets from  $\{1, 2, \dots, n\}$



# Counting Pairs

- The number of increasing pairs  $(i, j)$  with  $1 \leq i < j \leq n$  is the same as the number of 2-sets from  $\{1, 2, \dots, n\}$

Define  $f : X \rightarrow Y$  by  $f((i, j)) = \{i, j\}$

**Claim:**  $f$  is a **bijection** so  $|X| = |Y|$



# Counting Pairs

- The number of increasing pairs  $(i, j)$  with  $1 \leq i < j \leq n$  is the same as the number of 2-sets from  $\{1, 2, \dots, n\}$

Define  $f : X \rightarrow Y$  by  $f((i, j)) = \{i, j\}$

**Claim:**  $f$  is a **bijection** so  $|X| = |Y|$

We actually already saw that  $|X| = |Y| = \binom{n}{2}$



# The Bijection Principle

- Two sets have the same size if and only if there is a one-to-one function from one set onto the other.



# The Bijection Principle

- Two sets **have the same size** if and only if there is a **one-to-one function from one set onto the other**.

A standard first step in counting the size of a set is to use a bijection to show that it has the same size as a 2nd set, and then count the 2nd set instead.



# The Bijection Principle

- Two sets **have the same size** if and only if there is a **one-to-one function from one set onto the other**.

A standard first step in counting the size of a set is to use a bijection to show that it has the same size as a 2nd set, and then count the 2nd set instead.

In practice, in real problems we often only *implicitly* use the bijection and don't *explicitly* describe it



# The Bijection Principle

- Two sets **have the same size** if and only if there is a **one-to-one function from one set onto the other**.

A standard first step in counting the size of a set is to use a bijection to show that it has the same size as a 2nd set, and then count the 2nd set instead.

In practice, in real problems we often only *implicitly* use the bijection and don't *explicitly* describe it

Currently, we started with the problem of counting the **# of increasing triples** and changed it to the problem of counting the **# of 3-element sets from  $\{1, 2, \dots, n\}$**



# Next Lecture

- recurrence ...

