



# CS201 DISCRETE MATHEMATICS FOR COMPUTER SCIENCE

Dr. QI WANG

Department of Computer Science and Engineering


Office: Room413, CoE South Tower

Email: [wangqi@sustech.edu.cn](mailto:wangqi@sustech.edu.cn)

# Euler Circuits and Euler Paths

- **Theorem** A connected multigraph with at least two vertices has an *Euler circuit* if and only if **each of its vertices has even degree**.

**Theorem** A connected multigraph has an *Euler path* but not an *Euler circuit* if and only if **it has exactly two vertices of odd degree**.




**No Euler circuit**




# Planar Graphs

- Join three houses to each of three separate utilities.



# Planar Graphs


- Join three houses to each of three separate utilities.



Can this graph be drawn in the plane s.t. no two of its edges cross?

# Planar Graphs

- Join three houses to each of three separate utilities.



Can this graph be drawn in the plane s.t. no two of its edges cross?  $K_{3,3}$



# Planar Graphs


- **Definition** A graph is called *planar* if it can be drawn in the plane **without any edges crossing**. Such a drawing is called a *planar representation* of the graph.



# Planar Graphs

- **Definition** A graph is called *planar* if it can be drawn in the plane **without any edges crossing**. Such a drawing is called a *planar representation* of the graph.



**Example** Is  $K_4$  *planar*?



# Planar Graphs

- **Definition** A graph is called *planar* if it can be drawn in the plane **without any edges crossing**. Such a drawing is called a *planar representation* of the graph.


**Example** Is  $K_4$  *planar*?







# Planar Graphs

## ■ Example






# Planar Graphs

## ■ Example



# Planar Graphs




## ■ Example




(a)

# Planar Graphs

## ■ Example




(a)



(b)


# Planar Graphs

## ■ Example



# Planar Graphs

## ■ Example



## Applications

- ◇ IC design
- ◇ design of road networks



# Euler's Formula

- **Theorem (Euler's Formula)** Let  $G$  be a connected planar simple graph with  $e$  edges and  $v$  vertices. Let  $r$  be the number of regions in a planar representation of  $G$ . Then  $r = e - v + 2$ .

**Proof (by induction )**



# Euler's Formula

- **Theorem (Euler's Formula)** Let  $G$  be a connected planar simple graph with  $e$  edges and  $v$  vertices. Let  $r$  be the number of regions in a planar representation of  $G$ . Then  $r = e - v + 2$ .

**Proof (by induction )**

**Basic step:**






# Euler's Formula

- **Theorem (Euler's Formula)** Let  $G$  be a connected planar simple graph with  $e$  edges and  $v$  vertices. Let  $r$  be the number of regions in a planar representation of  $G$ . Then  $r = e - v + 2$ .

**Proof (by induction )**

**Basic step:**



# Euler's Formula

- **Theorem (Euler's Formula)** Let  $G$  be a connected planar simple graph with  $e$  edges and  $v$  vertices. Let  $r$  be the number of regions in a planar representation of  $G$ . Then  $r = e - v + 2$ .

**Proof (by induction )**

**Basic step:**

**Inductive Hypothesis:**



# Euler's Formula

- **Theorem (Euler's Formula)** Let  $G$  be a connected planar simple graph with  $e$  edges and  $v$  vertices. Let  $r$  be the number of regions in a planar representation of  $G$ . Then  $r = e - v + 2$ .

**Proof (by induction )**

**Basic step:**

**Inductive Hypothesis:**

$$r_k = e_k - v_k + 2$$



# Euler's Formula

- **Theorem (Euler's Formula)** Let  $G$  be a connected planar simple graph with  $e$  edges and  $v$  vertices. Let  $r$  be the number of regions in a planar representation of  $G$ . Then  $r = e - v + 2$ .

**Proof (by induction )**

**Basic step:**

**Inductive Hypothesis:**

**Inductive step:**



# Euler's Formula

- **Theorem (Euler's Formula)** Let  $G$  be a connected planar simple graph with  $e$  edges and  $v$  vertices. Let  $r$  be the number of regions in a planar representation of  $G$ . Then  $r = e - v + 2$ .

**Proof (by induction )**

**Basic step:**

**Inductive Hypothesis:**

**Inductive step:**

Let  $\{a_{k+1}, b_{k+1}\}$  be the edge that is added to  $G_k$  to obtain  $G_{k+1}$ .



# Euler's Formula

- **Theorem (Euler's Formula)** Let  $G$  be a connected planar simple graph with  $e$  edges and  $v$  vertices. Let  $r$  be the number of regions in a planar representation of  $G$ . Then  $r = e - v + 2$ .



**Proof (by induction )**

**Basic step:**

**Inductive Hypothesis:**

**Inductive step:**

Let  $\{a_{k+1}, b_{k+1}\}$  be the edge that is added to  $G_k$  to obtain  $G_{k+1}$ .




# The Degree of Regions

- **Definition** The *degree* of a *region* is defined to be the number of edges on the *boundary of this region*. When an edge occurs *twice* on the boundary, it contributes *two* to the degree.



# The Degree of Regions

- **Definition** The *degree* of a *region* is defined to be the number of edges on the *boundary of this region*. When an edge occurs *twice* on the boundary, it contributes *two* to the degree.





# Corollaries

- **Corollary 1** If  $G$  is a connected planar simple graph with  $e$  edges and  $v$  vertices, where  $v \geq 3$ , then  $e \leq 3v - 6$ .



# Corollaries

- **Corollary 1** If  $G$  is a connected planar simple graph with  $e$  edges and  $v$  vertices, where  $v \geq 3$ , then  $e \leq 3v - 6$ .

**Proof** The degree of every region is at least 3.



# Corollaries

- **Corollary 1** If  $G$  is a connected **planar** simple graph with  $e$  edges and  $v$  vertices, where  $v \geq 3$ , then  $e \leq 3v - 6$ .

**Proof** The degree of every region is **at least 3**.

- ◇  $G$  is **simple**
- ◇  $v \geq 3$



# Corollaries

- **Corollary 1** If  $G$  is a connected **planar** simple graph with  $e$  edges and  $v$  vertices, where  $v \geq 3$ , then  $e \leq 3v - 6$ .

**Proof** The degree of every region is **at least 3**.

- ◇  $G$  is **simple**
- ◇  $v \geq 3$

The sum of the degrees of the regions is **exactly twice** the number of edges in the graph.



# Corollaries

- **Corollary 1** If  $G$  is a connected planar simple graph with  $e$  edges and  $v$  vertices, where  $v \geq 3$ , then  $e \leq 3v - 6$ .

**Proof** The degree of every region is at least 3.

- ◇  $G$  is simple
- ◇  $v \geq 3$

The sum of the degrees of the regions is exactly twice the number of edges in the graph.

$$2e = \sum_{\text{all regions } R} \deg(R) \geq 3r$$



# Corollaries

- **Corollary 1** If  $G$  is a connected planar simple graph with  $e$  edges and  $v$  vertices, where  $v \geq 3$ , then  $e \leq 3v - 6$ .

**Proof** The degree of every region is at least 3.

- ◇  $G$  is simple
- ◇  $v \geq 3$

The sum of the degrees of the regions is exactly twice the number of edges in the graph.

$$2e = \sum_{\text{all regions } R} \deg(R) \geq 3r$$

By Euler's formula, the proof is completed.



# Corollaries

- **Corollary 2** If  $G$  is a connected planar simple graph, then  $G$  has a vertex of degree not exceeding 5.



# Corollaries

- **Corollary 2** If  $G$  is a connected planar simple graph, then  $G$  has a vertex of degree not exceeding 5.

**Proof**





# Corollaries

- **Corollary 2** If  $G$  is a connected planar simple graph, then  $G$  has a vertex of degree not exceeding 5.

## Proof

(By contradiction)

By Corollary 1 and the Handshaking Theorem.



# Corollaries

- **Corollary 2** If  $G$  is a connected planar simple graph, then  $G$  has a vertex of degree not exceeding 5.

## Proof

(By contradiction)

By Corollary 1 and the Handshaking Theorem.

**Corollary 3** In a connected planar simple graph has  $e$  edges and  $v$  vertices with  $v \geq 3$  and no circuits of length three, then  $e \leq 2v - 4$ .



# Corollaries

- **Corollary 2** If  $G$  is a connected planar simple graph, then  $G$  has a vertex of degree not exceeding 5.

## Proof

(By contradiction)

By Corollary 1 and the Handshaking Theorem.

**Corollary 3** In a connected planar simple graph has  $e$  edges and  $v$  vertices with  $v \geq 3$  and no circuits of length three, then  $e \leq 2v - 4$ .

**Proof** similar to that of Corollary 1.



# Examples

- Show that  $K_5$  is nonplanar.



# Examples

- Show that  $K_5$  is nonplanar.

Using Corollary 1



# Examples

- Show that  $K_5$  is nonplanar.

Using Corollary 1

Show that  $K_{3,3}$  is nonplanar.



# Examples

- Show that  $K_5$  is nonplanar.

Using Corollary 1

Show that  $K_{3,3}$  is nonplanar.

Using Corollary 3



# Examples

- Show that  $K_5$  is nonplanar.

Using Corollary 1

Show that  $K_{3,3}$  is nonplanar.



Using Corollary 3

Corollary 2 is used in the proof of Five Color Theorem.






# Only 5 Platonic Solids


				
<p>Tetrahedron <math>\{3, 3\}</math></p>	<p>Cube <math>\{4, 3\}</math></p>	<p>Octahedron <math>\{3, 4\}</math></p>	<p>Dodecahedron <math>\{5, 3\}</math></p>	<p>Icosahedron <math>\{3, 5\}</math></p>

# Only 5 Platonic Solids

				
Tetrahedron $\{3, 3\}$	Cube $\{4, 3\}$	Octahedron $\{3, 4\}$	Dodecahedron $\{5, 3\}$	Icosahedron $\{3, 5\}$



(a)



(b)




# Kuratowski's Theorem

- **Definition** If a graph is planar, so will be **any graph** obtained by **removing an edge  $\{u, v\}$  and adding a new vertex  $w$  together with edges  $\{u, w\}$  and  $\{w, v\}$** . Such an operation is called an *elementary subdivision*. The graphs  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  are called *homomorphic* if they can be obtained from **the same graph** by a sequence of elementary subdivisions.



# Kuratowski's Theorem

- **Definition** If a graph is planar, so will be **any graph** obtained by **removing an edge  $\{u, v\}$**  and adding a new vertex  $w$  together with edges  $\{u, w\}$  and  $\{w, v\}$ . Such an operation is called an *elementary subdivision*. The graphs  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  are called *homomorphic* if they can be obtained from **the same graph** by a sequence of elementary subdivisions.




# Kuratowski's Theorem

- **Definition** If a graph is planar, so will be **any graph** obtained by **removing an edge  $\{u, v\}$  and adding a new vertex  $w$  together with edges  $\{u, w\}$  and  $\{w, v\}$** . Such an operation is called an *elementary subdivision*. The graphs  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  are called *homomorphic* if they can be obtained from **the same graph** by a sequence of elementary subdivisions.



**Theorem** A graph is **nonplanar** if and only if it contains a subgraph **homomorphic to  $K_{3,3}$  or  $K_5$** .






# Examples



# Examples




# Examples





# Graph Coloring

- **Four-color theorem** Given any separation of a plane into contiguous regions, producing a figure called a *map*, no more than four colors are required to color the regions of the map so that no two adjacent regions have the same color.



## ■ Four-color theorem

- ◇ first proposed by Francis Guthrie in 1852
- ◇ his brother Frederick Guthrie told Augustus De Morgan
- ◇ De Morgan wrote to William Hamilton
- ◇ Alfred Kempe proved it **incorrectly** in 1879
- ◇ Percy Heawood found an error in 1890 and proved the *five-color theorem*
- ◇ Finally, Kenneth Appel and Wolfgang Haken proved it with case by case analysis by computer in 1976 (*the first computer-aided proof*)
- ◇ Kempe's incorrect proof serves as a basis



# Graph Coloring

- A *coloring* of a simple graph is the *assignment* of a color to *each vertex* of the graph so that *no two adjacent vertices* are assigned the same color.



# Graph Coloring

- A *coloring* of a simple graph is the *assignment* of a color to *each vertex* of the graph so that *no two adjacent vertices* are assigned the same color.



The *chromatic number* of a graph is the *least number* of colors needed for a coloring of this graph, denoted by  $\chi(G)$ .



# Graph Coloring

- A *coloring* of a simple graph is the *assignment* of a color to *each vertex* of the graph so that *no two adjacent vertices are assigned the same color*.

The *chromatic number* of a graph is the *least number* of colors needed for a coloring of this graph, denoted by  $\chi(G)$ .




# Graph Coloring

- **Theorem** (Four Color Theorem) The chromatic number of a planar graph is no greater than four.





# Graph Coloring

- **Theorem** (Four Color Theorem) The chromatic number of a planar graph is no greater than four.



# Graph Coloring

- **Theorem** (Four Color Theorem) The chromatic number of a planar graph is no greater than four.





# Graph Coloring

- **Theorem** (Six Color Theorem) The chromatic number of a planar graph is no greater than six.



# Graph Coloring

- **Theorem** (Six Color Theorem) The chromatic number of a planar graph is no greater than six.

**Proof** (by induction on the number of vertices)  
w.l.o.g., assume that the graph is connected.



# Graph Coloring

- **Theorem** (Six Color Theorem) The chromatic number of a planar graph is no greater than six.

**Proof** (by induction on the number of vertices)  
w.l.o.g., assume that the graph is connected.

**Basic step:** For one single vertex, pick an arbitrary color.



# Graph Coloring

- **Theorem** (Six Color Theorem) The chromatic number of a planar graph is no greater than six.

**Proof** (by induction on the number of vertices)  
w.l.o.g., assume that the graph is connected.

**Basic step:** For one single vertex, pick an arbitrary color.

**Inductive hypothesis:** Assume that every planar graph with  $k \geq 1$  or fewer vertices can be 6-colored.



# Graph Coloring

- **Theorem** (Six Color Theorem) The chromatic number of a planar graph is no greater than six.

**Proof** (by induction on the number of vertices)  
w.l.o.g., assume that the graph is connected.

**Basic step:** For one single vertex, pick an arbitrary color.

**Inductive hypothesis:** Assume that every planar graph with  $k \geq 1$  or fewer vertices can be 6-colored.

**Inductive step:** Consider a planar graph with  $k + 1$  vertices.



# Graph Coloring

- **Theorem** (Six Color Theorem) The chromatic number of a planar graph is no greater than six.

**Proof** (by induction on the number of vertices)  
w.l.o.g., assume that the graph is connected.

**Basic step:** For one single vertex, pick an arbitrary color.

**Inductive hypothesis:** Assume that every planar graph with  $k \geq 1$  or fewer vertices can be 6-colored.

**Inductive step:** Consider a planar graph with  $k + 1$  vertices. Recall Corollary 2 (the graph has a vertex of degree 5 or fewer). Remove this vertex, by i.h., we can color the remaining graph with 6 colors. Put the vertex back in. Since there are at most 5 colors adjacent, so we have at least one color left.



# Graph Coloring

- **Theorem** (Five Color Theorem) The chromatic number of a planar graph is no greater than five.



# Graph Coloring

- **Theorem** (Five Color Theorem) The chromatic number of a planar graph is no greater than five.

**Proof** (by induction on the number of vertices)  
w.l.o.g., assume that the graph is connected.





# Graph Coloring

- **Theorem** (**Five Color Theorem**) The chromatic number of a planar graph is no greater than five.

**Proof** (by induction on the number of vertices)  
w.l.o.g., assume that the graph is connected.

**Basic step:** For one single vertex, pick an arbitrary color.



# Graph Coloring

- **Theorem** (Five Color Theorem) The chromatic number of a planar graph is no greater than five.

**Proof** (by induction on the number of vertices)  
w.l.o.g., assume that the graph is connected.

**Basic step:** For one single vertex, pick an arbitrary color.

**Inductive hypothesis:** Assume that every planar graph with  $k \geq 1$  or fewer vertices can be 5-colored.



# Graph Coloring

- **Theorem** (Five Color Theorem) The chromatic number of a planar graph is no greater than five.

**Proof** (by induction on the number of vertices)  
w.l.o.g., assume that the graph is connected.

**Basic step:** For one single vertex, pick an arbitrary color.

**Inductive hypothesis:** Assume that every planar graph with  $k \geq 1$  or fewer vertices can be 5-colored.

**Inductive step:** Consider a planar graph with  $k + 1$  vertices.



# Graph Coloring

- **Theorem** (Five Color Theorem) The chromatic number of a planar graph is no greater than five.

**Proof** (by induction on the number of vertices)  
w.l.o.g., assume that the graph is connected.

**Basic step:** For one single vertex, pick an arbitrary color.

**Inductive hypothesis:** Assume that every planar graph with  $k \geq 1$  or fewer vertices can be 5-colored.

**Inductive step:** Consider a planar graph with  $k + 1$  vertices. Recall Corollary 2 (the graph has a vertex of degree 5 or fewer). Remove this vertex, by i.h., we can color the remaining graph with 5 colors. Put the vertex back in.





# Graph Coloring

- **Theorem** (Five Color Theorem) The chromatic number of a planar graph is no greater than five.

**Proof** (by induction on the number of vertices)  
w.l.o.g., assume that the graph is connected.

If the vertex has degree less than 5, or if it has degree 5 and only  $\leq 4$  colors are used for vertices connected to it, we can pick an available color for it.




# Graph Coloring

- **Theorem** (Five Color Theorem) The chromatic number of a planar graph is no greater than five.

**Proof** (by induction on the number of vertices)

If the vertex has degree 5, and all 5 colors are connected to it, we label the vertices adjacent to the “special” vertex (degree 5) 1 to 5 (in order).



# Graph Coloring

- **Theorem** (Five Color Theorem) The chromatic number of a planar graph is no greater than five.

**Proof** (by induction on the number of vertices)

We make a subgraph out of all the vertices colored 1 or 3. If the adjacent vertex colored 1 and the adjacent vertex colored 3 are not connected by a path in the subgraph.




# Graph Coloring

- **Theorem** (Five Color Theorem) The chromatic number of a planar graph is no greater than five.

**Proof** (by induction on the number of vertices)

We make a subgraph out of all the vertices colored 1 or 3. If the adjacent vertex colored 1 and the adjacent vertex colored 3 are not connected by a path in the subgraph.







# Graph Coloring

- **Theorem** (Five Color Theorem) The chromatic number of a planar graph is no greater than five.

**Proof** (by induction on the number of vertices)

We make a subgraph out of all the vertices colored 1 or 3. If the adjacent vertex colored 1 and the adjacent vertex colored 3 are not connected by a path in the subgraph.



# Graph Coloring

- **Theorem** (Five Color Theorem) The chromatic number of a planar graph is no greater than five.

**Proof** (by induction on the number of vertices)

On the other hand, if the vertices colored 1 and 3 are connected via a path in the subgraph, we do the **the same** for the vertices colored 2 and 4. Note that this will be a disconnected pair of subgraphs, separated by a path connecting the vertices colored 1 and 3 (**Why?**)




# Graph Coloring

- **Theorem** (Five Color Theorem) The chromatic number of a planar graph is no greater than five.

**Proof** (by induction on the number of vertices)

On the other hand, if the vertices colored 1 and 3 are connected via a path in the subgraph, we do the **the same** for the vertices colored 2 and 4. Note that this will be a disconnected pair of subgraphs, separated by a path connecting the vertices colored 1 and 3 (**Why?**)





# Graph Coloring

- **Theorem** (Five Color Theorem) The chromatic number of a planar graph is no greater than five.

**Proof** (by induction on the number of vertices)

On the other hand, if the vertices colored 1 and 3 are connected via a path in the subgraph, we do the **the same** for the vertices colored 2 and 4. Note that this will be a disconnected pair of subgraphs, separated by a path connecting the vertices colored 1 and 3 (**Why?**)




# Examples

- What is the chromatic number of  $K_n$ ,  $K_{m,n}$ ,  $C_n$ ?





# Examples

- What is the chromatic number of  $K_n$ ,  $K_{m,n}$ ,  $C_n$ ?






# Examples

- What is the chromatic number of  $K_n$ ,  $K_{m,n}$ ,  $C_n$ ?



# Examples





- What is the chromatic number of  $K_n$ ,  $K_{m,n}$ ,  $C_n$ ?





# Examples


- What is the chromatic number of  $K_n$ ,  $K_{m,n}$ ,  $C_n$ ?



# Applications of Graph Coloring

## ■ Scheduling Final Exams

Vertices represent courses, and there is an edge between two vertices if there is a common student in the courses.



Time Period

I

II

III

IV

Courses

1, 6

2

3, 5

4, 7



# Applications of Graph Coloring

## ■ Channel Assignments

Television channels 2 through 13 are assigned to stations in North America so that no two stations within 150 miles can operate on the same channel . How can the assignment of channels be modeled by graph coloring?



# Applications of Graph Coloring

## ■ Channel Assignments

Television channels 2 through 13 are assigned to stations in North America so that no two stations within 150 miles can operate on the same channel . How can the assignment of channels be modeled by graph coloring?

Graph Coloring  $\in$  NPC



# Next Lecture

- tree ...

