33. $\frac{1}{r} = \sin\theta + c\cos\theta$

34. $\frac{1}{y} = x^2 - 2 + ce^{-x^2/2}$

35. $y^2 = x^2 + cx - 1$

36. $\sin y = (1+x)(e^x + c)$

37. $(y - x + c)(xy + c) = 0$

38. $(2y - x^2 - c)(y - ce^x)(y + x - 1 - ce^{-x}) = 0$

39. $(x^2 - y^2 - 2c)(xy - c) = 0$

40. $(y^2 - x^2 - 2c)(3y^2 - 4x^3 - 6c) = 0$

41. $(y - c)(y + x^2 - c)(xy + cy + 1) = 0$

42. $(2y - x^2 - c)(2x - y^2 - c) = 0$

43. $4(x^2 + y)^3 = (3xy + 2x^3 + c)^2$

44. $y = \log(p^3 + p),\ x = 2\tan^{-1} p - p^{-1} + c$

45. $y = p^2x + p,\ (p-1)^2 x = \log p - p + c$

46. $4c^2x^2 - 8cy - 1 = 0$

47. General solution : $y = cx + c - c^2$
Singular solution : $4y = (x+1)^2$

48. General solution : $y = cx + \sin^{-1} c$

Singular solution : $y = \sqrt{x^2 - 1} + \sin^{-1}\left(\frac{\sqrt{x^2-1}}{x}\right)$

49. General solution : $y = cx + \frac{a}{c^2}$

Singular solution : $2ax^2 = (2ac + x)^3$

50. General solution : $y = cx + c - c^2$

Singular solution : $4y = (x+1)^2$

51. General solution : $y = cx + \cos^{-1} c$

Singular solution : $y = \sqrt{x^2 - 1} + \cos^{-1}\left(\frac{1}{x}\sqrt{x^2 - 1}\right)$

52. $x = (\log p - p + c)(p-1)^{-2}$ and the given equation.

53. $y = c - \frac{1}{1+p^2},\ x = \frac{p}{1+p^2} + \tan^{-1} p$

54. $x = c - 2\{p + \log(p-1)\},\ y = c - 2\left\{\frac{1}{2}p^2 + p + \log(p-1)\right\}$

55. $y^2 = cx^2 + \frac{c}{c-1}$

56. General solution : $cxy - a^3c^2x = 1$
Singular solution : $x(xy^2 - 4a^3) = 0$

CHAPTER 2

Linear Differential Equations of Second Order

2.1 INTRODUCTION

A linear differential equation of the nth order has the form

$$\frac{d^n y}{dx^n} + P_1\frac{d^{n-1}y}{dx^{n-1}} + P_2\frac{d^{n-2}y}{dx^{n-2}} + \ldots + P_{n-1}\frac{dy}{dx} + P_n y = X \qquad \ldots(2.1)$$

where $P_1, P_2, \ldots, P_n, X$ are functions of x only or constants. The general solution of (2.1) contains n arbitrary constants.

Non-homogeneous

Differential equation (2.1) is said to be non-homogeneous if the right hand side of (2.1), $X \neq 0$. Otherwise

Homogeneous

when $X = 0$.

Linear Differential Equations of First Order with Constant Coefficients

A linear differential equation of first order with constant coefficients is of the form

$$\frac{dy}{dx} + Py = Q \qquad \ldots(2.2)$$

Using the symbol D for the differential operator $\frac{d}{dx}$, (2.2) becomes

$$(D + P)y = Q, \quad \text{or} \quad f(D)y = Q, \text{ where } f(D) = D + P \qquad \ldots(2.3)$$

Here P is constant or a function of x only and Q is constant or a function of x only.

$$\therefore \qquad \text{I.F.} = e^{\int P dx}$$

Multiplying both sides of (2.2) by $e^{\int P dx}$, we get

$$e^{\int P dx}\frac{dy}{dx} + Pye^{\int P dx} = Qe^{\int P dx}, \quad \text{or} \quad \frac{d}{dx}(ye^{\int P dx}) = Qe^{\int P dx}$$

or $$d(ye^{\int Pdx}) = (Qe^{\int Pdx})dx$$

Integrating, $$\int d(ye^{\int Pdx}) = \int (Qe^{\int Pdx})dx$$

$\therefore$ $$ye^{\int Pdx} = c + \int (Qe^{\int Pdx})dx,$$ where c is an arbitrary constant.

$\therefore$ $$y = cu + v, \text{ where } u = e^{-\int Pdx},\ v = e^{-\int Pdx}\int (Qe^{\int Pdx})dx \qquad ...(2.4)$$

Now, $$\frac{du}{dx} = -Pe^{-\int Pdx} = -Pu, \quad \text{or} \quad \frac{du}{dx} + Pu = 0$$

or $$\frac{d}{dx}(cu) + P(cu) = 0$$

Therefore cu is the general solution (since it contains one arbitrary constant c) of the corresponding homogeneous equation of (2.2).

Also, $$\frac{dv}{dx} = -Pe^{-\int Pdx}\int (Qe^{\int Pdx})dx + e^{-\int Pdx}Qe^{\int Pdx}$$

or $$\frac{dv}{dx} = -Pv + Q, \quad \text{or} \quad \frac{dv}{dx} + Pv = Q$$

Therefore v is a solution of (2.2).

Hence the general solution of (2.2) is (2.4) consisting of two parts, *i.e.*, cu and v, where cu is the general solution of the corresponding homogeneous equation known as *complementary function* and v is known as *particular integral* (since it is free from any arbitrary constant).

$\therefore$ *General solution (or complete solution)*

= *Complementary Function + Particular Integral*

or $$y = \text{C.F.} + \text{P.I.}$$

Linear Differential Equations of Second Order with Constant Coefficients

The general form of a linear differential equation of second order with constant coefficients is

$$\frac{d^2y}{dx^2} + P_1\frac{dy}{dx} + P_2y = X \qquad ...(2.5)$$

where P_1, P_2 are constants and X is a function of x only or constant.

Using the symbol D for the differential operator $\frac{d}{dx}$, (2.5) becomes

$$(D^2 + P_1D + P_2)y = X, \quad \text{or} \quad f(D)y = X, \text{ where } f(D) = D^2 + P_1D + P_2 \qquad ...(2.6)$$

When $X = 0$, then $f(D)y = 0$, or $$\frac{d^2y}{dx^2} + P_1\frac{dy}{dx} + P_2y = 0 \qquad ...(2.7)$$

is called the homogeneous equation of (2.5) or (2.6).

Definition: Two solutions $y_1(x)$, $y_2(x)$ of (2.7) are said to be *linearly independent* if $c_1y_1 + c_2y_2 = 0$ for arbitrary constants c_1, c_2 implies $c_1 = c_2 = 0$.

Theorem 1: Two solutions $y_1(x)$, $y_2(x)$ of (2.7) are linearly independent if and only if their Wronskian $W(y_1, y_2) = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} \neq 0$.

Proof: Let us suppose that y_1, y_2 are linearly dependent.

Then by definition there exist constants c_1, c_2 not all zero, such that

$$c_1y_1 + c_2y_2 = 0, \text{ therefore, } c_1y_1' + c_2y_2' = 0.$$

Since this system of two equations in c_1, c_2 has a nontrivial solution (*i.e.*, c_1, c_2 are not all zero), we must have $\begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = 0$, *i.e.*, $W(y_1, y_2) = 0$.

Conversely, let $W(y_1, y_2) = 0$, then there exists a nontrivial solution (*i.e.*, c_1, c_2 are not all zero) to the following homogeneous equations

$$c_1y_1 + c_2y_2 = 0$$
$$c_1y_1' + c_2y_2' = 0$$

This implies there exist constants c_1, c_2 not all zero, such that $c_1y_1 + c_2y_2 = 0$ and hence y_1, y_2 are dependent.

Theorem 2: If $y_1(x)$ and $y_2(x)$ are any two linearly independent solutions of the homogeneous differential equation (2.7), then the general solution of (2.7) is given by $y(x) = c_1y_1(x) + c_2y_2(x)$, where c_1 and c_2 are two arbitrary constants.

Proof: Since $y_1(x)$, $y_2(x)$ are solutions of (2.7), therefore

$$\frac{d^2y_1}{dx^2} + P_1\frac{dy_1}{dx} + P_2y_1 = 0 \qquad ...(2.8)$$

and $$\frac{d^2y_2}{dx^2} + P_1\frac{dy_2}{dx} + P_2y_2 = 0 \qquad ...(2.9)$$

If c_1, c_2 are two arbitrary constants, then

$$\frac{d^2}{dx^2}(c_1y_1 + c_2y_2) + P_1\frac{d}{dx}(c_1y_1 + c_2y_2) + P_2(c_1y_1 + c_2y_2)$$

$$= c_1\left(\frac{d^2y_1}{dx^2} + P_1\frac{dy_1}{dx} + P_2y_1\right) + c_2\left(\frac{d^2y_2}{dx^2} + P_1\frac{dy_2}{dx} + P_2y_2\right)$$

$$= c_1 \cdot 0 + c_2 \cdot 0 = 0 \qquad \text{[by (2.8) and (2.9)]}$$

Hence $y(x) = c_1y_1(x) + c_2y_2(x)$ is the general solution of (2.7), since it contains two arbitrary constants c_1 and c_2.

Note: This superposition principle is not applicable to non-homogeneous and non-linear equations.

Theorem 3: If $u(x)$ be the general solution of the homogeneous differential equation (2.7) and $v(x)$ be a particular solution (*i.e.*, free from any arbitrary constant) of (2.5), then the general solution of the differential equation (2.5) is $y = u(x) + v(x)$.

Proof: Since $u(x)$ is the general solution of (2.7),

therefore, $$\frac{d^2u}{dx^2}+P_1\frac{du}{dx}+P_2u=0 \qquad ...(2.10)$$

Since $v(x)$ is a solution of (2.5),

therefore, $$\frac{d^2v}{dx^2}+P_1\frac{dv}{dx}+P_2v=X \qquad ...(2.11)$$

Adding (2.10) and (2.11), we get

$$\frac{d^2}{dx^2}(u+v)+P_1\frac{d}{dx}(u+v)+P_2(u+v)=X, \text{ or } \frac{d^2y}{dx^2}+P_1\frac{dy}{dx}+P_2y=X$$

This shows that $y=u(x)+v(x)$ is the general solution of (2.5).

Note: To obtain the general (or complete) solution of (2.5), we have first to obtain two independent solutions of (2.7) and any solution (free from any arbitrary constant) of (2.5). The sum of the general solution of (2.7) and a particular solution of (2.5) will be the general solution of (2.5).

The expression which is the general (or complete) solution of (2.7) is known as the *Complementary Function* of (2.5) and any particular solution of (2.5) is known as the *Particular Integral* of (2.5).

Thus the process of solving

$$\frac{d^2y}{dx^2}+P_1\frac{dy}{dx}+P_2y=X$$

involves following steps:

(*i*) Replace X by 0 and then find the complete solution $u=c_1u_1+c_2u_2$ of the reduced equation.

(*ii*) Find any particular solution of the original equation.

(*iii*) Sum of the complementary function found in step (*i*) and the particular integral found in step (*ii*) gives the general solution of the given differential equation.

These conclusions are also valid for general linear differential equations with constant coefficients.

$\therefore$ *General Solution (or Complete Solution)*

= *Complementary Function + Particular Integral*

or $y=$ C.F. + P.I.

2.2 METHOD FOR FINDING THE COMPLEMENTARY FUNCTION (C.F.)

1. In finding the complementary function, right hand side of the given differential equation is replaced by zero.
2. The C.F. of a linear second order differential equation with constant coefficients can be found from

$$\frac{d^2y}{dx^2}+P_1\frac{dy}{dx}+P_2y=0, \quad \text{or} \quad (D^2+P_1D+P_2)y=0 \qquad ...(2.12)$$

where $D\equiv\dfrac{d}{dx}$ and P_1, P_2 are constants.

Let e^{mx}, where m is a constant, be a solution of (2.12).

Substituting $y=e^{mx}$ in (2.12), we get

$$(m^2+P_1m+P_2)e^{mx}=0, \quad \text{or} \quad m^2+P_1m+P_2=0 \qquad (\because e^{mx}\neq 0) \qquad ...(2.13)$$

The algebraic equation (2.13) is known as the *characteristic* or *auxiliary equation* of (2.12). Also $y=e^{mx}$ is known as the *trial solution* of (2.12).

Note: If we write (2.12) in the form $f(D)y=0$, then the auxiliary equation is $f(m)=0$.

3. Solve the auxiliary equation.

Case I: Roots are real and unequal

If m_1, m_2 are two real and distinct roots of (2.13), then C.F. $=c_1e^{m_1x}+c_2e^{m_2x}$, where c_1, c_2 are two arbitrary constants.

Case II: Roots are real and equal

Let the roots of the auxiliary equation (2.13) be real and equal to α. Then (2.12) can be written as

$$(D-\alpha)^2y=0, \quad \text{or} \quad (D^2-2\alpha D+\alpha^2)y=0, \quad \text{or} \quad \frac{d^2y}{dx^2}-2\alpha\frac{dy}{dx}+\alpha^2y=0 \qquad ...(2.14)$$

Let its solution be $y=ve^{\alpha x}$, where v is a function of x.

Substituting $y=ve^{\alpha x}$, in (2.14), we get

$$\alpha^2ve^{\alpha x}+2\alpha e^{\alpha x}\frac{dv}{dx}+e^{\alpha x}\frac{d^2v}{dx^2}-2\alpha\left(\alpha ve^{\alpha x}+e^{\alpha x}\frac{dv}{dx}\right)+\alpha^2ve^{\alpha x}=0$$

or $$\frac{d^2v}{dx^2}=0 \qquad (\because e^{\alpha x}\neq 0)$$

The solution of this equation is $v=c_1x+c_2$.

Hence C.F. $=(c_1x+c_2)e^{\alpha x}$, where c_1, c_2 are arbitrary constants.

Case III: Roots are complex

Let the auxiliary equation (2.13) has complex roots $\alpha\pm i\beta$ (α, β real), then

$$\begin{aligned}\text{C.F.}&=c_1e^{(\alpha+i\beta)x}+c_2e^{(\alpha-i\beta x)}=e^{\alpha x}(c_1e^{i\beta x}+c_2e^{-i\beta x})\\&=e^{\alpha x}[c_1(\cos\beta x+i\sin\beta x)+c_2(\cos\beta x-i\sin\beta x)]\\&=e^{\alpha x}[(c_1+c_2)\cos\beta x+i(c_1-c_2)\sin\beta x]\\&=e^{\alpha x}(A\cos\beta x+B\sin\beta x).\end{aligned}$$

ILLUSTRATIVE EXAMPLES

Example 1: Solve: $\dfrac{d^2y}{dx^2}+(a+b)\dfrac{dy}{dx}+aby=0$, where a, b are unequal.

Solution: The given equation can be written as

$$\{D^2+(a+b)D+ab\}y=0, \quad \text{where} \quad D\equiv\frac{d}{dx}.$$

Let $y = e^{mx}$ be a trial solution. Then the auxiliary equation is $m^2 + (a+b)m + ab = 0$, or $(m+a)(m+b) = 0$

$\therefore \quad m = -a, -b.$

Hence the general solution is

$$y = c_1 e^{-ax} + c_2 e^{-bx}, \; c_1, c_2 \text{ are arbitrary constants.}$$

Example 2: Solve the equation $\frac{d^2y}{dx^2} - 4\frac{dy}{dx} + 4y = 0$ and find the particular solution if $y = 1$, $\frac{dy}{dx} = -2$ when $x = 0$.

Solution: The given equation can be written as

$$(D^2 - 4D + 4)y = 0, \text{ where } D \equiv \frac{d}{dx}.$$

Let $y = e^{mx}$ be a trial solution. Then the auxiliary equation is $m^2 - 4m + 4 = 0$, or $(m-2)^2 = 0$. Therefore, $m = 2, 2$.

Hence the general solution is

$$y = (c_1 x + c_2)e^{2x}, \text{ where } c_1, c_2 \text{ are two arbitrary constants.}$$

$$\therefore \quad \frac{dy}{dx} = c_1 e^{2x} + 2(c_1 x + c_2)e^{2x} = (2c_1 x + c_1 + 2c_2)e^{2x}.$$

By the given condition $y = 1$, $\frac{dy}{dx} = -2$ when $x = 0$.

$\therefore \quad c_2 = 1, \qquad c_1 + 2c_2 = -2, \quad \text{or} \quad c_1 = -4 \qquad (\because c_2 = 1).$

Therefore the particular solution is

$$y = (1 - 4x)e^{2x}.$$

Example 3: Solve: $\frac{d^2y}{dx^2} + 8\frac{dy}{dx} + 25y = 0$.

Solution: The given equation can be written as

$$(D^2 + 8D + 25)y = 0, \quad \text{where} \quad D \equiv \frac{d}{dx}.$$

Let $y = e^{mx}$ be a trial solution. Then the auxiliary equation is $m^2 + 8m + 25 = 0$.

Thus $m = -4 \pm 3i$ and the general solution is $y = e^{-4x}(c_1 \cos 3x + c_2 \sin 3x)$, where c_1, c_2 are two arbitrary constants.

Example 4: Solve: $\frac{d^2s}{dt^2} + n^2 s = 0$; $s = a$, $\frac{ds}{dt} = 0$ when $t = 0$.

Solution: The given equation can be written as

$$(D^2 + n^2)s = 0, \quad \text{where} \quad D \equiv \frac{d}{dt}.$$

Let $s = e^{mt}$ be a trial solution. Then the auxiliary equation is $m^2 + n^2 = 0$, or $m = \pm in$.

Hence the general solution is

$$s = c_1 \cos nt + c_2 \sin nt, \text{ where } c_1, c_2 \text{ are two arbitrary constants.}$$

$$\therefore \quad \frac{ds}{dt} = -c_1 n \sin nt + c_2 n \cos nt.$$

By the given condition $s = a$, $\frac{ds}{dt} = 0$ when $t = 0$.

$\therefore \quad c_1 = a, \quad c_2 n = 0, \quad \text{or} \quad c_2 = 0 \qquad (\because n \neq 0).$

Therefore, the required solution is $s = a \cos nt$.

2.3 THE OPERATOR D

The operators D, D^2, D^3, ... stand respectively for $\frac{d}{dx}, \frac{d^2}{dx^2}, \frac{d^3}{dx^3}, \ldots$. Also $\frac{1}{D}$ or D^{-1}, $\frac{1}{D^2}$ or D^{-2}, ... are used to denote the inverse operators, *i.e.*, the operators which integrate respectively once, twice, ...

This symbolic operator D largely satisfies the ordinary laws of algebra. We have $D(u+v) = Du + Dv$, so D satisfies distributive law. Also $(D+a)u = Du + au = au + Du = (a+D)u$ and $Dau = aDu$, so D is commutative with a constant a. It is also noted that $D^m D^n u = D^{m+n}u$ where m, n are integers. If α, β are constants, then

$$(D+\alpha)(D+\beta)u = [D^2 + (\alpha+\beta)D + \alpha\beta]u$$
$$= (D+\beta)(D+\alpha)u.$$

Thus D obeys the fundamental laws of algebra except that it is not commutative with variables.

2.4 RULES FOR FINDING PARTICULAR INTEGRAL (P.I.)

Inverse operator $\frac{1}{f(D)}$

The expression $\frac{1}{f(D)}X$ denotes a function of x, not containing any arbitrary constant, which gives X when operated by $f(D)$, *i.e.*, $f(D)\left\{\frac{1}{f(D)}X\right\} = X$. Also $\frac{1}{f(D)}\{f(D)X\} = X$. Therefore, the operator $\frac{1}{f(D)}$ is the inverse of the operator $f(D)$ and vice-versa.

Results (*i*) $\frac{1}{D}X = \int X\,dx \qquad$ (*ii*) $\frac{1}{D-a}X = e^{ax}\int Xe^{-ax}dx$

Proof: (*i*) Let $y = \frac{1}{D}X$, therefore, $Dy = D\left(\frac{1}{D}X\right)$

or $$\frac{dy}{dx} = X \qquad \text{(by definition)}$$

$\therefore$ $$dy = X\,dx, \quad \text{by integration,} \quad y = \int X\,dx.$$

(*ii*) Let $y = \frac{1}{D-a}X$, therefore, $(D-a)y = (D-a)\left(\frac{1}{D-a}X\right)$

or $$Dy - ay = X \qquad \text{(by definition)}$$

$\therefore$ $$\frac{dy}{dx} - ay = X(x), \text{ which is a linear equation in } y \qquad ...(2.15)$$

$\therefore$ $$\text{I.F.} = e^{-\int a dx} = e^{-ax}$$

Multiplying both sides of (2.15) by e^{-ax}, we get

$$e^{-ax}\frac{dy}{dx} - aye^{-ax} = Xe^{-ax}, \quad \text{or} \quad \frac{d}{dx}(ye^{-ax}) = Xe^{-ax}$$

or $$d(ye^{-ax}) = Xe^{-ax}dx$$

Integrating, $$\int d(ye^{-ax}) = \int Xe^{-ax}dx.$$

or $$ye^{-ax} = \int Xe^{-ax}dx.$$

$\therefore$ $$y = \frac{1}{D-a}X = e^{ax}\int Xe^{-ax}dx.$$

General method of finding particular integral (P.I.)

Let us consider the following linear differential equation of second order with constant coefficients

$$\frac{d^2y}{dx^2} + P_1\frac{dy}{dx} + P_2 y = X$$

or $$f(D)y = X, \text{ where } f(D) = D^2 + P_1 D + P_2,\ D \equiv \frac{d}{dx}.$$

Here P_1, P_2 are constants and X is a function of x only or constant.

Let m_1, m_2 be the roots of the auxiliary equation of the corresponding homogeneous equation, then

$$f(D) = (D - m_1)(D - m_2).$$

Let $\frac{1}{f(D)}$ can be resolved into partial fractions, say,

$$\frac{1}{f(D)} = \frac{A_1}{D-m_1} + \frac{A_2}{D-m_2}, \text{ where } A_1, A_2 \text{ are constants.}$$

Then $$\text{P.I.} = \frac{1}{f(D)}X = \frac{A_1}{D-m_1}X + \frac{A_2}{D-m_2}X$$

$$= A_1 e^{m_1 x}\int Xe^{-m_1 x}dx + A_2 e^{m_2 x}\int Xe^{-m_2 x}dx \quad \text{[by Result (ii)]} \qquad ...(2.16)$$

Note: Since (2.16) is a particular solution, it must not contain any arbitrary constant.

ILLUSTRATIVE EXAMPLES

Example 1: Solve: $\frac{d^2y}{dx^2} - 5\frac{dy}{dx} + 6y = e^{4x}$. (Arch–201/13)

Solution: The given equation can be written as

$$(D^2 - 5D + 6)y = e^{4x}, \quad \text{or} \quad (D-3)(D-2)y = e^{4x}, \text{ where } D \equiv \frac{d}{dx}.$$

First step: Determination of complementary function (C.F.)

The complementary function is found from

$$(D-3)(D-2) = 0.$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $(m-3)(m-2) = 0$, the roots of which are 2, 3.

$\therefore$ $$\text{C.F.} = c_1 e^{2x} + c_2 e^{3x}.$$

Second step: Determination of particular integral (P.I.)

$$\text{P.I.} = \frac{1}{(D-3)(D-2)}e^{4x} = \left(\frac{1}{D-3} - \frac{1}{D-2}\right)e^{4x}$$

$$= \frac{1}{D-3}e^{4x} - \frac{1}{D-2}e^{4x} = e^{3x}\int e^{4x}\cdot e^{-3x}dx - e^{2x}\int e^{4x}\cdot e^{-2x}dx$$

$$= e^{3x}\int e^{x}dx - e^{2x}\int e^{2x}dx = e^{3x}\cdot e^{x} - e^{2x}\cdot\frac{e^{2x}}{2}$$

$$= \frac{e^{4x}}{2}.$$

Third step: The general solution is therefore $y = \text{C.F.} + \text{P.I.} = c_1 e^{2x} + c_2 e^{3x} + \frac{e^{4x}}{2}$, where c_1, c_2 are arbitrary constants.

Example 2: Solve: $\frac{d^2y}{dx^2} - y = 4xe^{x}$.

Solution: The given equation can be written as

$$(D^2 - 1)y = 4xe^{x}, \quad \text{or} \quad (D-1)(D+1)y = 4xe^{x}, \text{ where } D \equiv \frac{d}{dx}.$$

First step: Determination of complementary function (C.F.)

The complementary function is found from

$$(D-1)(D+1)y = 0.$$

Indian Institute of Engineering Science Ramanujan Central Library

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $(m-1)(m+1) = 0$, the roots of which are $-1, 1$.

$$\therefore \qquad \text{C.F.} = c_1 e^{-x} + c_2 e^{x}.$$

Second step: Determination of particular integral (P.I.)

$$\text{P.I.} = \frac{1}{(D-1)(D+1)} 4xe^x = \frac{1}{2}\left(\frac{1}{D-1} - \frac{1}{D+1}\right) 4xe^x$$

$$= 2 \cdot \frac{1}{D-1} xe^x - 2 \cdot \frac{1}{D+1} xe^x = 2e^x \int xe^x \cdot e^{-x} dx - 2e^{-x} \int xe^x \cdot e^x dx$$

$$= 2e^x \int x\,dx - 2e^{-x} \int xe^{2x} dx$$

$$= 2e^x \cdot \frac{x^2}{2} - 2e^{-x}\left[x\int e^{2x} dx - \int\left\{\left(\frac{dx}{dx}\right)\int e^{2x} dx\right\} dx\right]$$

$$= x^2 e^x - 2e^{-x}\left[\frac{xe^{2x}}{2} - \frac{e^{2x}}{4}\right] = e^x\left(x^2 - x + \frac{1}{2}\right).$$

Third step: The general solution is therefore

$$y = \text{C.F.} + \text{P.I.} = c_1 e^{-x} + c_2 e^x + e^x\left(x^2 - x + \frac{1}{2}\right),$$

where c_1, c_2 are two arbitrary constants.

Example 3: Solve the following differential equation

$$\frac{d^2y}{dx^2} + a^2 y = \sec ax$$

with the symbolic operator D.

Solution: The given differential equation can be written as $(D^2 + a^2)y = \sec ax$,

or $$(D + ia)(D - ia)y = \sec ax, \text{ where } D \equiv \frac{d}{dx}.$$

First step: Determination of complementary function (C.F.)

The complementary function is found from

$$(D + ia)(D - ia)y = 0.$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $(m + ia)(m - ia) = 0$, the roots of which are $-ia, ia$.

$$\therefore \qquad \text{C.F.} = c_1 \cos ax + c_2 \sin ax.$$

Second step: Determination of particular integral (P.I.)

$$\text{P.I.} = \frac{1}{(D+ia)(D-ia)} \sec ax = \frac{1}{2ia}\left(\frac{1}{D-ia} - \frac{1}{D+ia}\right)\sec ax \qquad \text{...(1)}$$

Now, $$\frac{1}{D-ia}\sec ax = e^{iax}\int e^{-iax} \sec ax\, dx$$

$$= e^{iax}\int \frac{(\cos ax - i\sin ax)}{\cos ax} dx \qquad [\because e^{i\theta} = \cos\theta + i\sin\theta]$$

$$= e^{iax}\left(x + \frac{i}{a}\log\cos ax\right)$$

$$= (\cos ax + i\sin ax)\left(x + \frac{i}{a}\log\cos ax\right)$$

$$= \left(x\cos ax - \frac{1}{a}\sin ax \log\cos ax\right)$$

$$+ i\left(x\sin ax + \frac{1}{a}\cos ax\log\cos ax\right)$$

Similarly (replacing i by $-i$), we have

$$\frac{1}{D+ia}\sec ax = \left(x\cos ax - \frac{1}{a}\sin ax\log\cos ax\right)$$

$$-i\left(x\sin ax + \frac{1}{a}\cos ax\log\cos ax\right)$$

Therefore, from (1) we get,

$$\text{P.I.} = \frac{1}{2ia}\left(\frac{1}{D-ia}\sec ax - \frac{1}{D+ia}\sec ax\right)$$

$$= \frac{x\sin ax}{a} + \frac{\cos ax\log\cos ax}{a^2}.$$

Third step: The general solution is therefore $y = \text{C.F.} + \text{P.I.} = c_1 \cos ax + c_2 \sin ax + \dfrac{x\sin ax}{a} + \dfrac{\cos ax\log\cos ax}{a^2}$, where c_1, c_2 are two arbitrary constants.

2.5 SHORT METHODS FOR FINDING PARTICULAR INTEGRALS (P.I.) IN SOME SPECIAL CASES

For the differential equation

$$\frac{d^2y}{dx^2} + P_1\frac{dy}{dx} + P_2 y = X, \text{ or } f(D)y = X, \text{ where } f(D) = D^2 + P_1 D + P_2,$$

$$D \equiv \frac{d}{dx}, \quad \text{P.I.} = \frac{1}{f(D)} X.$$

Let us consider the following special types:

Case I: $X = x^m$, m is a positive integer

$$\text{P.I.} = \frac{1}{f(D)} x^m = \{f(D)\}^{-1} x^m.$$

Expand $\{f(D)\}^{-1}$ in ascending powers of D and operate on x^m. The terms of this expansion beyond mth power of D need not be considered, because the result of their operation on x^m becomes zero.

Note: This result is also valid for general nth $(n > 2)$ order linear differential equations with constant coefficients.

Example: Solve: $\dfrac{d^2y}{dx^2} - 4\dfrac{dy}{dx} + 4y = x^2 + x.$

Solution: The given differential equation can be written as $(D^2 - 4D + 4)y = x^2 + x$, or $(D-2)^2 y = x^2 + x$, where $D \equiv \dfrac{d}{dx}$.

First step: Determination of complementary function (C.F.). The complementary function is found from

$$(D-2)^2 y = 0.$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $(m-2)^2 = 0$, the roots of which are 2, 2.

$$\therefore \qquad \text{C.F.} = (c_1 x + c_2)e^{2x}.$$

Second step: Determination of particular integral (P.I.)

$$\text{P.I.} = \frac{1}{(D-2)^2}(x^2+x) = \frac{1}{4\left(1-\frac{D}{2}\right)^2}(x^2+x) = \frac{1}{4}\left(1-\frac{D}{2}\right)^{-2}(x^2+x)$$

$$= \frac{1}{4}\left\{1 + D + \frac{3}{4}D^2 + \ldots\right\}(x^2+x)$$

$$= \frac{1}{4}\left\{x^2 + x + 2x + 1 + \frac{3}{2}\right\} = \frac{1}{4}\left(x^2 + 3x + \frac{5}{2}\right).$$

Third step: The general solution is therefore $y = \text{C.F.} + \text{P.I.} = (c_1 x + c_2)e^{2x} + \dfrac{1}{4}\left(x^2 + 3x + \dfrac{5}{2}\right)$, where c_1, c_2 are two arbitrary constants.

Note: Remember

$$(1+x)^{-1} = 1 - x + x^2 - x^3 + \ldots$$

$$(1-x)^{-1} = 1 + x + x^2 + x^3 + \ldots$$

$$(1+x)^{-2} = 1 - 2x + 3x^2 - 4x^3 + \ldots$$

$$(1-x)^{-2} = 1 + 2x + 3x^2 + 4x^3 + \ldots$$

Case II: $X = e^{ax}$, a is a constant. Then

$$(i)\ \text{P.I.} = \frac{1}{f(D)}e^{ax} = \frac{e^{ax}}{f(a)}, \text{ if } f(a) \neq 0 \qquad \ldots(2.17)$$

$$(ii)\ \text{P.I.} = \frac{1}{f(D)}e^{ax} = x\frac{e^{ax}}{f'(a)}, \text{ if } f(a) = 0 \text{ but } f'(a) \neq 0 \qquad \ldots(2.18)$$

$$(iii)\ \text{P.I.} = \frac{1}{f(D)}e^{ax} = x^2\frac{e^{ax}}{f''(a)}, \text{ if } f(a) = f'(a) = 0 \qquad \ldots(2.19)$$

Proof: (*i*) Here $De^{ax} = ae^{ax}$, $D^2e^{ax} = a^2e^{ax}$.

$$\therefore \qquad (D^2 + P_1D + P_2)e^{ax} = (a^2 + P_1a + P_2)e^{ax}, \text{ i.e., } f(D)e^{ax} = f(a)e^{ax}.$$

Operating on both sides by $\dfrac{1}{f(D)}$, we have

$$\frac{1}{f(D)}\{f(D)e^{ax}\} = \frac{1}{f(D)}\{f(a)e^{ax}\}, \quad \text{or} \quad e^{ax} = f(a)\frac{1}{f(D)}e^{ax}$$

$$\therefore \qquad \frac{1}{f(D)}e^{ax} = \frac{e^{ax}}{f(a)}, \text{ provided } f(a) \neq 0.$$

(*ii*) If $f(a) = 0$, *i.e.*, a is a root of the auxiliary equation of the corresponding homogeneous differential equation, the above rule fails and we proceed as follows.

Here $D - a$ is a factor of $f(D)$. Suppose

$$f(D) = (D-a)\,\varphi(D), \text{ where } f'(a) = \varphi(a) \neq 0. \text{ Then}$$

$$\frac{1}{f(D)}e^{ax} = \frac{1}{D-a}\cdot\frac{1}{\varphi(D)}e^{ax} = \frac{1}{D-a}\cdot\frac{1}{\varphi(a)}e^{ax} \qquad \text{[by (2.17)]}$$

$$= \frac{1}{f'(a)}\cdot\frac{1}{D-a}e^{ax} = \frac{1}{f'(a)}e^{ax}\int e^{ax}\cdot e^{-ax}dx$$

$$= \frac{1}{f'(a)}e^{ax}\int dx = \frac{xe^{ax}}{f'(a)}, \text{ provided } f(a) = 0,\ f'(a) \neq 0.$$

(*iii*) If $f(a) = f'(a) = 0$, then $f(D) = (D-a)^2$, for the differential equation under consideration.

$$\therefore \qquad \frac{1}{f(D)}e^{ax} = \frac{1}{(D-a)^2}e^{ax} = \frac{1}{D-a}\cdot\frac{1}{D-a}e^{ax} = \frac{1}{D-a}e^{ax}\int e^{ax}\cdot e^{-ax}dx$$

$$= \frac{1}{D-a}xe^{ax} = e^{ax}\int xe^{ax}\cdot e^{-ax}dx = e^{ax}\int x\,dx$$

$$= \frac{x^2}{2}e^{ax} = x^2\frac{e^{ax}}{f''(a)} \qquad [\because f''(D) = 2], \text{ provided } f(a) = f'(a) = 0.$$

Note: These results are also valid for general nth $(n > 2)$ order linear differential equations with constant coefficients provided for (*iii*) $f''(a) \neq 0$ and so on.

Example: Solve: $\dfrac{d^2y}{dx^2} - 3\dfrac{dy}{dx} + 2y = e^x$, if $y = 3$ and $\dfrac{dy}{dx} = 3$, when $x = 0$.

Solution: The given differential equation can be written as $(D^2 - 3D + 2)y = e^x$, or $(D-1)(D-2)y = e^x$, where $D \equiv \dfrac{d}{dx}$.

First step: Determination of complementary function (C.F.)

The complementary function is found from

$$(D-1)(D-2)y = 0.$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $(m-1)(m-2) = 0$, the roots of which are 1, 2.

$$\therefore \qquad \text{C.F.} = c_1 e^x + c_2 e^{2x}.$$

Second step: Determination of particular integral (P.I.)

$$\text{P.I.} = \frac{1}{(D-1)(D-2)} e^x = \frac{1}{(D-1)(1-2)} e^x = -\frac{1}{D-1} e^x$$

$$= -e^x \int e^x \cdot e^{-x} dx = -e^x \int dx = -xe^x.$$

Third step: The general solution is therefore $y = \text{C.F.} + \text{P.I.} = c_1 e^x + c_2 e^{2x} - xe^x$, where c_1, c_2 are two arbitrary constants.

$$\therefore \qquad \frac{dy}{dx} = c_1 e^x + 2c_2 e^{2x} - e^x - xe^x = (c_1 - 1 - x)e^x + 2c_2 e^{2x}.$$

By question, $y = \dfrac{dy}{dx} = 3$, when $x = 0$.

$$\therefore \qquad \left.\begin{aligned} c_1 + c_2 &= 3 \\ c_1 - 1 + 2c_2 &= 3 \end{aligned}\right\} \quad \text{solving, } c_1 = 2,\ c_2 = 1.$$

Hence the required solution is $y = (1-x)e^x + 2e^{2x}$.

Case III: $X = e^{ax}V$, where V is any function of x.

Then
$$\text{P.I.} = \frac{1}{f(D)} e^{ax} V = e^{ax} \cdot \frac{1}{f(D+a)} V. \qquad \text{...(2.20)}$$

Proof: If u is a function of x, then

$$D(e^{ax}u) = e^{ax}Du + ae^{ax}u = e^{ax}(D+a)u$$

$$D^2(e^{ax}u) = e^{ax}D^2u + 2ae^{ax}Du + a^2e^{ax}u$$
$$= e^{ax}(D+a)^2 u.$$

$$\therefore \qquad (D^2 + P_1 D + P_2)e^{ax}u = e^{ax}\{(D+a)^2 + P_1(D+a) + P_2\}u$$

$$\therefore \qquad f(D)e^{ax}u = e^{ax} f(D+a)u$$

Operating on both sides by $\dfrac{1}{f(D)}$, we have

$$\frac{1}{f(D)}\{f(D)e^{ax}u\} = \frac{1}{f(D)}\{e^{ax} f(D+a)u\}$$

or
$$e^{ax}u = \frac{1}{f(D)}\{e^{ax} f(D+a)u\}$$

Let us put $f(D+a)u = V$, *i.e.*, $u = \dfrac{1}{f(D+a)} V$, therefore

$$e^{ax} \frac{1}{f(D+a)} V = \frac{1}{f(D)}(e^{ax}V), \text{ i.e., } \frac{1}{f(D)}(e^{ax}V) = e^{ax} \frac{1}{f(D+a)} V.$$

Note: This result is also valid for general nth $(n > 2)$ order linear differential equations with constant coefficients.

Example 1: Solve: $\dfrac{d^2y}{dx^2} - 2\dfrac{dy}{dx} + y = xe^x$. [M–201/13]

Solution: The given differential equation can be written as $(D-1)^2 y = xe^x$, where $D \equiv \dfrac{d}{dx}$.

First step: Determination of C.F.

The complementary function is found from $(D-1)^2 y = 0$.

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $(m-1)^2 = 0$, the roots of which are 1, 1.

$$\therefore \qquad \text{C.F.} = (C_1 + C_2 x)e^x$$

Second step: Determination of P.I.

$$\text{P.I.} = \frac{1}{(D-1)^2} xe^x = e^x \frac{1}{(D+1-1)^2} x = e^x \frac{1}{D^2} x$$

$$= e^x \frac{1}{D} \frac{x^2}{2} = e^x \frac{x^3}{6}.$$

Third step: The general solution is given by $y = (C_1 + C_2 x)e^x + \dfrac{1}{6}x^3 e^x$, where C_1, C_2 are two arbitrary constants.

Example 2: Solve: $\dfrac{d^2y}{dx^2} - 4y = x \sinh x$.

Solution: The given differential equation can be written as $(D^2 - 4)y = x \sinh x$, or $(D-2)(D+2)y = x \sinh x$, where $D \equiv \dfrac{d}{dx}$.

First step: Determination of complementary function (C.F.)

The complementary function is found from

$$(D-2)(D+2)y = 0.$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $(m-2)(m+2) = 0$, the roots of which are 2, –2.

$$\therefore \qquad \text{C.F.} = c_1 e^{2x} + c_2 e^{-2x}$$

Second step: Determination of particular integral (P.I.)

$$\text{P.I.} = \frac{1}{D^2 - 4} x \sinh x = \frac{1}{D^2 - 4} x \frac{1}{2}(e^x - e^{-x})$$

$$= \frac{1}{2}\left\{\frac{1}{D^2-4}e^x x - \frac{1}{D^2-4}e^{-x}x\right\} = \frac{1}{2}\left\{e^x \frac{1}{(D+1)^2-4}x - e^{-x}\frac{1}{(D-1)^2-4}x\right\}$$

$$= \frac{1}{2}\left\{e^x \frac{1}{D^2+2D-3}x - e^{-x}\frac{1}{D^2-2D-3}x\right\}$$

$$= \frac{1}{2}\left[\frac{e^x}{-3}\left\{1-\left(\frac{2D}{3}+\frac{D^2}{3}\right)\right\}^{-1}x - \frac{e^{-x}}{-3}\left\{1+\left(\frac{2D}{3}-\frac{D^2}{3}\right)\right\}^{-1}x\right]$$

$$= -\frac{1}{6}\left[e^x\left(1+\frac{2D}{3}+\frac{D^2}{3}+\ldots\right)x - e^{-x}\left(1-\frac{2D}{3}+\frac{D^2}{3}+\ldots\right)x\right]$$

$$= -\frac{1}{6}\left[e^x\left(x+\frac{2}{3}\right)-e^{-x}\left(x-\frac{2}{3}\right)\right] = -\frac{x}{3}\left(\frac{e^x-e^{-x}}{2}\right)-\frac{2}{9}\left(\frac{e^x+e^{-x}}{2}\right)$$

$$= -\frac{x}{3}\sinh x - \frac{2}{9}\cosh x.$$

Third step: The general solution is therefore $y = \text{C.F.} + \text{P.I.} = c_1e^{2x} + c_2e^{-2x} - \frac{x}{3}\sinh x - \frac{2}{9}\cosh x$, where c_1, c_2 are two arbitrary constants.

Note: (*i*) $\sinh x = \frac{1}{2}(e^x - e^{-x})$, $\cosh x = \frac{1}{2}(e^x + e^{-x})$,

$$\tanh x = \frac{\sinh x}{\cosh x},\quad \coth x = \frac{1}{\tanh x},\quad \text{cosech } x = \frac{1}{\sinh x},$$

$$\text{sech } x = \frac{1}{\cosh x},\quad \cosh^2 x - \sinh^2 x = 1$$

(*ii*) $\frac{d}{dx}\sinh x = \cosh x$, $\frac{d}{dx}\cosh x = \sinh x$.

Case IV: $X = \sin(ax+b)$ or $\cos(ax+b)$. Then

(*i*) $$\text{P.I.} = \frac{1}{f(D)}\sin(ax+b) = \frac{1}{\varphi(D^2)}\sin(ax+b)$$

$$= \frac{1}{\varphi(-a^2)}\sin(ax+b), \text{ provided } \varphi(-a^2)\neq 0, \qquad \ldots(2.21)$$

(*ii*) $$\text{P.I.} = \frac{1}{f(D)}\cos(ax+b) = \frac{1}{\varphi(D^2)}\cos(ax+b)$$

$$= \frac{1}{\varphi(-a^2)}\cos(ax+b), \text{ provided } \varphi(-a^2)\neq 0, \qquad \ldots(2.22)$$

if $f(D) = \varphi(D^2)$ and it is possible to express $f(D)$ in terms of $\varphi(D^2)$.

Proof: (*i*) Now

$$D\sin(ax+b) = a\cos(ax+b)$$

$$D^2\sin(ax+b) = -a^2\sin(ax+b)$$

$$D^3\sin(ax+b) = -a^3\cos(ax+b)$$

$$D^4\sin(ax+b) = a^4\sin(ax+b)$$

$$\therefore \quad D^2\sin(ax+b) = (-a^2)\sin(ax+b),$$

$$(D^2)^2\sin(ax+b) = (-a^2)^2\sin(ax+b).$$

In general $(D^2)^r\sin(ax+b) = (-a^2)^r\sin(ax+b)$, $r = 1, 2, 3, \ldots$

$$\therefore \quad \varphi(D^2)\sin(ax+b) = \varphi(-a^2)\sin(ax+b).$$

Operating on both sides by $\frac{1}{\varphi(D^2)}$, we have

$$\frac{1}{\varphi(D^2)}\{\varphi(D^2)\sin(ax+b)\} = \frac{1}{\varphi(D^2)}\{\varphi(-a^2)\sin(ax+b)\}$$

or $$\sin(ax+b) = \varphi(-a^2)\frac{1}{\varphi(D^2)}\sin(ax+b)$$

$$\therefore \quad \frac{1}{\varphi(D^2)}\sin(ax+b) = \frac{1}{\varphi(-a^2)}\sin(ax+b), \text{ provided } \varphi(-a^2)\neq 0.$$

(*ii*) Proceed as in (*i*).

Note: If $\varphi(-a^2) = 0$, proceed with the method illustrated in Example 2 below, or use the rule:

$$\frac{1}{f(D)}\{\sin(ax+b) \text{ or } \cos(ax+b)\} = x\frac{1}{f'(D)}\{\sin(ax+b) \text{ or } \cos(ax+b)\}$$

Example 1: Solve $\frac{d^2y}{dx^2} + y = 2\cos^2 x$.

Solution: The given differential equation can be written as $(D^2+1)y = 2\cos^2 x$, where $D \equiv \frac{d}{dx}$.

First step: Determination of complementary function (C.F.).

The complementary function is found from

$$(D^2+1)\,y = 0.$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $m^2 + 1 = 0$, the roots of which are $i, -i$.

$$\therefore \quad \text{C.F.} = c_1\cos x + c_2\sin x.$$

Second step: Determination of particular integral (P.I.).

$$\text{P.I.} = \frac{1}{D^2+1}2\cos^2 x = \frac{1}{D^2+1}(1+\cos 2x) = (1+D^2)^{-1}1 + \frac{1}{D^2+1}\cos 2x$$

$$= (1-D^2+\ldots)1 + \frac{1}{-(2)^2+1}\cos 2x \qquad \text{[replacing } D^2 \text{ by } -(2)^2 \text{ in the second term]}$$

$$= 1 - \frac{1}{3}\cos 2x.$$

Third step: The general solution is therefore

$$y = \text{C.F.} + \text{P.I.} = c_1 \cos x + c_2 \sin x + 1 - \frac{1}{3}\cos 2x,$$

where c_1, c_2 are two arbitrary constants.

Example 2: Solve $\dfrac{d^2y}{dx^2} - 2\dfrac{dy}{dx} + 2y = x + e^x \cos x$.

Solution: Let us write the given differential equation as

$$(D^2 - 2D + 2)\,y = x + e^x \cos x, \text{ where } D \equiv \frac{d}{dx}.$$

First step: Determination of complementary function (C.F.).
The complementary function is found from

$$(D^2 - 2D + 2)y = 0$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $m^2 - 2m + 2 = 0$, the roots of which are $1 \pm i$.

$$\therefore \qquad \text{C.F.} = e^x(c_1 \cos x + c_2 \sin x).$$

Second step: Determination of particular integral (P.I.).

$$\text{P.I.} = \frac{1}{D^2 - 2D + 2}(x + e^x \cos x) = \frac{1}{D^2 - 2D + 2}x + \frac{1}{D^2 - 2D + 2}(e^x \cos x)$$

$$= \frac{1}{2}\left\{1 - \left(D - \frac{D^2}{2}\right)\right\}^{-1} x + e^x \frac{1}{(D+1)^2 - 2(D+1) + 2}\cos x$$

$$= \frac{1}{2}\left(1 + D - \frac{D^2}{2} + \ldots\right)x + e^x \frac{1}{D^2 + 1}\cos x \qquad \ldots(1)$$

In $\dfrac{1}{D^2+1}\cos x$, if we replace D^2 by -1^2, the method fails.

$$\text{Now, } \frac{1}{D^2+1}(\cos x + i \sin x) = \frac{1}{D^2+1}e^{ix} = e^{ix}\frac{1}{(D+i)^2+1}1$$

$$= e^{ix}\frac{1}{D^2 + 2iD}1 = e^{ix}\frac{1}{2iD}\left(1 + \frac{D}{2i}\right)^{-1}1$$

$$= e^{ix}\frac{1}{2iD}\left(1 - \frac{D}{2i} + \ldots\right)1 = e^{ix}\cdot\frac{1}{2i}\cdot\frac{1}{D}1 = \frac{e^{ix}}{2i}x$$

$$= \frac{(\cos x + i \sin x)x}{2i} = \frac{1}{2}x \sin x - \frac{i}{2}x \cos x$$

$$\therefore \qquad \frac{1}{D^2+1}(\cos x + i \sin x) = \frac{1}{2}x \sin x - \frac{i}{2}x \cos x$$

Equating real part from both sides, we get

$$\frac{1}{D^2+1}\cos x = \frac{1}{2}x \sin x$$

Hence, from (1), we get

$$\text{P.I.} = \frac{1}{2}(x+1) + \frac{1}{2}xe^x \sin x.$$

Third step: The general solution is therefore

$$y = \text{C.F.} + \text{P.I.} = e^x(c_1 \cos x + c_2 \sin x) + \frac{1}{2}(x+1) + \frac{1}{2}xe^x \sin x,$$

where c_1, c_2 are two arbitrary constants.

Note: $\dfrac{1}{D^2+1}\cos x = x\dfrac{1}{2D}\cos x = \dfrac{x}{2}\displaystyle\int \cos x\, dx = \dfrac{x}{2}\sin x$

Example 3: Solve $(D^2 - 4D + 3)y = \sin 3x \cos 2x$, where $D \equiv \dfrac{d}{dx}$.

Solution:
First step: Determination of complementary function (C.F.).
The complementary function is found from

$$(D^2 - 4D + 3)y = 0.$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $m^2 - 4m + 3 = 0$, or $m^2 - 3m - m + 3 = 0$, or $(m-3)(m-1) = 0$, the roots are $m = 3, 1$.

$$\therefore \qquad \text{C.F.} = c_1 e^{3x} + c_2 e^x.$$

Second step: Determination of particular integral (P.I.).

$$\text{P.I.} = \frac{1}{D^2 - 4D + 3}(\sin 3x \cos 2x) = \frac{1}{D^2 - 4D + 3}\left\{\frac{1}{2}(\sin 5x + \sin x)\right\}$$

$$= \frac{1}{2}\frac{D^2 + 3 + 4D}{(D^2+3)^2 - 16D^2}\sin 5x + \frac{1}{2}\frac{D^2 + 3 + 4D}{(D^2+3)^2 - 16D^2}\sin x$$

$$= \frac{1}{2}\frac{D^2 + 4D + 3}{(-5^2+3)^2 - 16(-5^2)}\sin 5x + \frac{1}{2}\frac{D^2 + 4D + 3}{(-1^2+3)^2 - 16(-1^2)}\sin x$$

$$= \frac{1}{2}\cdot\frac{1}{884}(-25 \sin 5x + 20 \cos 5x + 3 \sin 5x)$$

$$+ \frac{1}{2}\cdot\frac{1}{20}(-\sin x + 4 \cos x + 3 \sin x)$$

$$= \frac{1}{884}(10 \cos 5x - 11 \sin 5x) + \frac{1}{20}(\sin x + 2 \cos x)$$

Third step: The general solution is therefore

$$y = \text{C.F.} + \text{P.I.} = c_1 e^{3x} + c_2 e^x + \frac{1}{884}(10 \cos 5x - 11 \sin 5x) + \frac{1}{20}(\sin x + 2 \cos x),$$

where c_1, c_2 are arbitrary constants.

Case V: $X = xV$, where V is any function of x.

Then
$$\text{P.I.} = \frac{1}{f(D)}xV = \left\{x - \frac{1}{f(D)}f'(D)\right\}\frac{1}{f(D)}V \qquad \ldots(2.23)$$

Proof: Let $$V_1 = \frac{1}{f(D)}V.$$

Now, $$D(xV_1) = xDV_1 + V_1$$

$$D^2(xV_1) = DV_1 + xD^2V_1 + DV_1 = xD^2V_1 + 2DV_1$$

$$\therefore \quad (D^2 + P_1D + P_2)xV_1 = xD^2V_1 + 2DV_1 + P_1xDV_1 + P_1V_1 + P_2xV_1$$

$$= x(D^2 + P_1D + P_2)V_1 + (2D + P_1)V_1$$

Thus $$f(D)xV_1 = xf(D)V_1 + f'(D)V_1$$

or $$f(D)x\left\{\frac{1}{f(D)}V\right\} = xf(D)\left\{\frac{1}{f(D)}V\right\} + f'(D)\left\{\frac{1}{f(D)}V\right\} \qquad \left[\because V_1 = \frac{1}{f(D)}V\right]$$

or $$f(D)x\left\{\frac{1}{f(D)}V\right\} = xV + f'(D)\left\{\frac{1}{f(D)}V\right\}$$

or $$x\left\{\frac{1}{f(D)}V\right\} = \frac{1}{f(D)}xV + \frac{1}{f(D)}f'(D)\left\{\frac{1}{f(D)}V\right\}$$

$$\therefore \quad \frac{1}{f(D)}xV = \left\{x - \frac{1}{f(D)}f'(D)\right\}\frac{1}{f(D)}V.$$

Note: (*i*) By successive applications of this method the P.I. of $X = x^mV$, m is a positive integer, may be found.

(*ii*) The result (2.23) is also valid for general nth ($n > 2$) order linear differential equations with constant coefficients.

Example 1: Solve $\dfrac{d^2y}{dx^2} + 4y = x\cos x.$

Solution: Let us write the given differential equation as $(D^2 + 4)y = x\cos x$, where $D \equiv \dfrac{d}{dx}$.

First step: Determination of complementary function (C.F.).

The complementary function is found from

$$(D^2 + 4)y = 0.$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $m^2 + 4 = 0$, the roots of which are $\pm 2i$.

$$\therefore \quad \text{C.F.} = c_1\cos 2x + c_2\sin 2x.$$

Second step: Determination of particular integral (P.I.).

$$\text{P.I.} = \frac{1}{D^2+4}x\cos x = \left(x - \frac{2D}{D^2+4}\right)\frac{1}{D^2+4}\cos x$$

$$= \left(x - \frac{2D}{D^2+4}\right)\frac{\cos x}{-1^2+4} \qquad \text{(replacing } D^2 \text{ by } -1^2\text{)}$$

$$= \frac{x}{3}\cos x + \frac{2}{3}\cdot\frac{1}{D^2+4}\sin x = \frac{x}{3}\cos x + \frac{2}{3}\cdot\frac{\sin x}{-1^2+4}$$

$$= \frac{x}{3}\cos x + \frac{2}{9}\sin x.$$

Third step: The general solution is therefore

$$y = \text{C.F.} + \text{P.I.} = c_1\cos 2x + c_2\sin 2x + \frac{x}{3}\cos x + \frac{2}{9}\sin x,$$

where c_1, c_2 are two arbitrary constants.

Example 2: Solve $\dfrac{d^2y}{dx^2} - y = x^2\sin x.$

Solution: Let us write the given differential equation as $(D^2 - 1)y = x^2\sin x$, where $D \equiv \dfrac{d}{dx}$.

First step: Determination of complementary function (C.F.).

The complementary function is found from

$$(D^2 - 1)y = 0.$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $m^2 - 1 = 0$, the roots of which are ± 1.

$$\therefore \quad \text{C.F.} = c_1e^x + c_2e^{-x}.$$

Second step: Determination of particular integral (P.I.).

$$\text{P.I.} = \frac{1}{D^2-1}x^2\sin x = \frac{1}{D^2-1}\{x(x\sin x)\}$$

$$= \left(x - \frac{2D}{D^2-1}\right)\frac{1}{D^2-1}(x\sin x) = \left(x - \frac{2D}{D^2-1}\right)\left(x - \frac{2D}{D^2-1}\right)\frac{1}{D^2-1}\sin x$$

$$= \left(x - \frac{2D}{D^2-1}\right)\left(x - \frac{2D}{D^2-1}\right)\frac{\sin x}{-1^2-1} \qquad \text{(replacing } D^2 \text{ by } -1^2\text{)}$$

$$= -\frac{1}{2}\left(x - \frac{2D}{D^2-1}\right)\left(x\sin x - \frac{1}{D^2-1}2\cos x\right) = -\frac{1}{2}\left(x - \frac{2D}{D^2-1}\right)\left(x\sin x - \frac{2\cos x}{-1^2-1}\right)$$

$$= -\frac{1}{2}\left\{x^2\sin x + x\cos x - \frac{1}{D^2-1}2(\sin x + x\cos x - \sin x)\right\}$$

$$= -\frac{1}{2}(x^2\sin x + x\cos x) + \frac{1}{D^2-1}x\cos x$$

$$= -\frac{1}{2}(x^2\sin x + x\cos x) + \left(x - \frac{2D}{D^2-1}\right)\frac{1}{D^2-1}\cos x$$

$$= -\frac{1}{2}(x^2\sin x + x\cos x) + \left(x - \frac{2D}{D^2-1}\right)\frac{\cos x}{-1^2-1}$$

$$= -\frac{1}{2}(x^2\sin x + x\cos x) - \frac{1}{2}\left(x\cos x + \frac{1}{D^2-1}2\sin x\right)$$

$$= -\frac{1}{2}(x^2\sin x + x\cos x) - \frac{1}{2}\left(x\cos x + \frac{2\sin x}{-1^2-1}\right)$$

$$= -\frac{1}{2}x^2\sin x - x\cos x + \frac{1}{2}\sin x$$

Third step: The general solution is therefore $y = \text{C.F.} + \text{P.I.} = c_1e^x + c_2e^{-x} - \frac{1}{2}x^2\sin x - x\cos x + \frac{1}{2}\sin x$, where c_1, c_2 are two arbitrary constants.

MISCELLANEOUS EXAMPLES

Example 1: Solve the differential equation

$$\frac{d^2y}{dx^2} - 5\frac{dy}{dx} + 6y = e^x\cos x \qquad \textit{(W.B.U.T. 2002, 2011)}$$

Solution: Let us write the given differential equation as

$$(D^2 - 5D + 6)y = e^x\cos x, \text{ where } D \equiv \frac{d}{dx}.$$

First step: Determination of complementary function (C.F.).

The complementary function is found from

$$(D^2 - 5D + 6)y = 0.$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $m^2 - 5m + 6 = 0$, or $(m-3)(m-2) = 0$, the roots of which are 3, 2.

$$\therefore \qquad \text{C.F.} = c_1e^{3x} + c_2e^{2x}.$$

Second step: Determination of particular integral (P.I.).

$$\text{P.I.} = \frac{1}{D^2 - 5D + 6}e^x\cos x = e^x\frac{1}{(D+1)^2 - 5(D+1) + 6}\cos x$$

$$= e^x\frac{1}{D^2 - 3D + 2}\cos x = e^x\frac{D^2 + 2 + 3D}{(D^2+2)^2 - 9D^2}\cos x$$

$$= e^x\frac{D^2 + 3D + 2}{(-1^2+2)^2 - 9(-1^2)}\cos x$$

$$= \frac{e^x}{10}(D^2 + 3D + 2)\cos x = \frac{e^x}{10}(-\cos x - 3\sin x + 2\cos x)$$

$$= \frac{e^x}{10}(\cos x - 3\sin x).$$

Third step: The general solution is therefore

$$y = \text{C.F.} + \text{P.I.} = c_1e^{3x} + c_2e^{2x} + \frac{e^x}{10}(\cos x - 3\sin x), \text{ where } c_1, c_2 \text{ are}$$

two arbitrary constants.

Example 2: Solve $(D^2 - 2D)y = e^x\sin x$, where $D \equiv \frac{d}{dx}$. (W.B.U.T. 2

Solution:

First step: Determination of complementary function (C.F.).

The complementary function is found from

$$(D^2 - 2D)y = 0.$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $m^2 - 2m = 0$, or $m(m-2) = 0$, the roots of which are 0, 2.

$$\therefore \qquad \text{C.F.} = c_1e^{0x} + c_2e^{2x} = c_1 + c_2e^{2x}.$$

Second step: Determination of particular integral (P.I.).

$$\text{P.I.} = \frac{1}{D^2 - 2D}e^x\sin x = e^x\frac{1}{(D+1)^2 - 2(D+1)}\sin x = e^x\frac{1}{D^2 - 1}\sin x$$

$$= e^x\frac{\sin x}{-1^2 - 1} = -\frac{1}{2}e^x\sin x.$$

Third step: The general solution is therefore

$$y = \text{C.F.} + \text{P.I.} = c_1 + c_2e^{2x} - \frac{1}{2}e^x\sin x, \text{ where, } c_1, c_2 \text{ are two arbitrary}$$

constants.

Example 3: Solve: $(D^2 + 4)y = x\sin^2 x$, where $D \equiv \frac{d}{dx}$. *(W.B.U.T. 2008, 2010)*

Solution:

First step: Determination of complementary function (C.F.).

The complementary function is found from

$$(D^2 + 4)y = 0.$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $m^2 + 4 = 0$, the roots are $m = \pm 2i$.

$$\therefore \qquad \text{C.F.} = c_1\cos 2x + c_2\sin 2x.$$

Second step:

$$\text{P.I.} = \frac{1}{D^2 + 4}x\sin^2 x = \frac{1}{2}\cdot\frac{1}{D^2 + 4}x(1 - \cos 2x) = \frac{1}{2}\cdot\frac{1}{D^2 + 4}x - \frac{1}{2}\frac{1}{D^2 + 4}x\cos 2x \qquad ...(1)$$

Now,

$$\frac{1}{D^2 + 4}x = \frac{1}{4}\left(1 + \frac{D^2}{4}\right)^{-1}x = \frac{1}{4}\left(1 - \frac{D^2}{4} + \ldots\right)x = \frac{x}{4} \qquad ...(2)$$

$$\frac{1}{D^2 + 4}x\cos 2x = \left(x - \frac{2D}{D^2 + 4}\right)\frac{1}{D^2 + 4}\cos 2x \qquad ...(3)$$

In $\frac{1}{D^2 + 4}\cos 2x$, if we replace D^2 by -2^2, the method fails.

$$\frac{1}{D^2+4}(\cos 2x + i\sin 2x) = \frac{1}{D^2+4}e^{2ix} = e^{2ix}\frac{1}{(D+2i)^2+4}1$$

$$= e^{2ix}\frac{1}{D^2+4iD}1 = e^{2ix}\frac{1}{4iD}\left(1+\frac{D}{4i}\right)^{-1}1 = e^{2ix}\frac{1}{4iD}\left(1-\frac{D}{4i}+\ldots\right)1$$

$$= e^{2ix}\cdot\frac{1}{4i}\cdot\frac{1}{D}1 = \frac{e^{2ix}}{4i}x = \frac{(\cos 2x + i\sin 2x)x}{4i} = \frac{1}{4}x\sin 2x - \frac{i}{4}x\cos 2x$$

$$\therefore \quad \frac{1}{D^2+4}(\cos 2x + i\sin 2x) = \frac{1}{4}x\sin 2x - \frac{i}{4}x\cos 2x.$$

Equating real and imaginary parts from both sides, we get

$$\frac{1}{D^2+4}\cos 2x = \frac{1}{4}x\sin 2x, \quad \frac{1}{D^2+4}\sin 2x = -\frac{1}{4}x\cos 2x \qquad \ldots(4)$$

Therefore, from (3), we get

$$\frac{1}{D^2+4}x\cos 2x = \left(x - \frac{2D}{D^2+4}\right)\frac{1}{4}x\sin 2x = \frac{1}{4}x^2\sin 2x - \frac{1}{2}\frac{1}{D^2+4}D(x\sin 2x)$$

$$= \frac{1}{4}x^2\sin 2x - \frac{1}{2}\cdot\frac{1}{D^2+4}(\sin 2x + 2x\cos 2x)$$

$$= \frac{1}{4}x^2\sin 2x - \frac{1}{2}\cdot\frac{1}{D^2+4}\sin 2x - \frac{1}{D^2+4}x\cos 2x$$

$$\therefore \quad 2\cdot\frac{1}{D^2+4}x\cos 2x = \frac{1}{4}x^2\sin 2x - \frac{1}{2}\cdot\frac{1}{D^2+4}\sin 2x$$

$$= \frac{1}{4}x^2\sin 2x + \frac{1}{8}x\cos 2x \qquad \text{[by (4)]}$$

$$\therefore \quad \frac{1}{D^2+4}x\cos 2x = \frac{1}{8}x^2\sin 2x + \frac{1}{16}x\cos 2x \qquad \ldots(5)$$

From (1), (2), (3) and (5), we get

$$\text{P.I.} = \frac{x}{8} - \frac{x^2\sin 2x}{16} - \frac{x\cos 2x}{32}.$$

Third step: The general solution is therefore

$$y = \text{C.F.} + \text{P.I.} = c_1\cos 2x + c_2\sin 2x + \frac{x}{8} - \frac{x^2\sin 2x}{16} - \frac{x\cos 2x}{32},$$

where c_1, c_2 are two arbitrary constants.

Example 4: Solve the differential equation

$$\frac{d^2y}{dx^2} - 5\frac{dy}{dx} + 6y = x^2e^{3x} \qquad \textit{(W.B.U.T. 2009, 2010)}$$

Solution: Let us write the given differential equation as

$$(D^2 - 5D + 6)y = x^2e^{3x}, \text{ where } D \equiv \frac{d}{dx}.$$

First step: Determination of complementary function (C.F.).

The complementary function is found from

$$(D^2 - 5D + 6)y = 0.$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $m^2 - 5m + 6 = 0$, or $(m-3)(m-2) = 0$, the roots are $m = 2, 3$.

$$\therefore \quad \text{C.F.} = c_1e^{2x} + c_2e^{3x}.$$

Second step: Determination of particular integral (P.I.).

$$\text{P.I.} = \frac{1}{D^2 - 5D + 6}(x^2e^{3x}) = e^{3x}\cdot\frac{1}{(D+3)^2 - 5(D+3) + 6}x^2$$

$$= e^{3x}\cdot\frac{1}{D^2+D}x^2 = e^{3x}\cdot\frac{1}{D}(1+D)^{-1}x^2$$

$$= e^{3x}\cdot\frac{1}{D}(1 - D + D^2 - D^3 + \ldots)x^2 = e^{3x}\frac{1}{D}(x^2 - 2x + 2)$$

$$= e^{3x}\left(\frac{1}{3}x^3 - x^2 + 2x\right).$$

Third step: The general solution is therefore

$$y = \text{C.F.} + \text{P.I.} = c_1e^{2x} + c_2e^{3x} + e^{3x}\left(\frac{1}{3}x^3 - x^2 + 2x\right),$$

where c_1, c_2 are two arbitrary constants.

Example 5: Solve $\dfrac{d^2y}{dx^2} + 4\dfrac{dy}{dx} + 4y = 3\sin x + 4\cos x$, $y(0) = 1$ and $y'(0) = 0$.

Solution: Let us write the given differential equation as

$$(D^2 + 4D + 4)y = 3\sin x + 4\cos x, \text{ or } (D+2)^2y = 3\sin x + 4\cos x,$$

where $D \equiv \dfrac{d}{dx}$.

First step: Determination of complementary function (C.F.).

The complementary function is found from

$$(D+2)^2y = 0.$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $(m+2)^2 = 0$, the roots are $m = -2, -2$.

$$\therefore \quad \text{C.F.} = (c_1 + c_2x)e^{-2x}.$$

Second step: Determination of particular integral (P.I.).

$$\text{P.I.} = \frac{1}{(D+2)^2}(3\sin x + 4\cos x) = \frac{(D-2)^2}{(D^2-4)^2}(3\sin x + 4\cos x)$$

$$= \frac{D^2-4D+4}{(-1^2-4)^2}(3\sin x+4\cos x)$$

$$= \frac{1}{25}\{-3\sin x-4\cos x-4(3\cos x-4\sin x)+12\sin x+16\cos x\}$$

$$= \frac{1}{25}(25\sin x) = \sin x.$$

Third step: The general solution is therefore

$$y = \text{C.F.} + \text{P.I.} = (c_1+c_2x)e^{-2x}+\sin x, \text{ where } c_1, c_2 \text{ are two arbitrary constants.}$$

$$\therefore \quad y'(x) = \frac{dy}{dx} = c_2e^{-2x}-2(c_1+c_2x)e^{-2x}+\cos x.$$

By question, $y(0) = 1$ and $y'(0) = 0$.

$\therefore \quad c_1 = 1, \quad c_2-2c_1+1 = 0, \quad$ or $\quad c_2 = 1 \qquad (\because c_1 = 1)$

Hence the required solution is

$$y = (1+x)e^{-2x}+\sin x.$$

Example 6: Solve: $\dfrac{d^2y}{dx^2}-3\dfrac{dy}{dx}+2y = xe^{3x}+\sin 2x.$

Solution: Let us write the given differential equation as

$$(D^2-3D+2)y = xe^{3x}+\sin 2x, \text{ where } D \equiv \frac{d}{dx}.$$

First step: Determination of complementary function (C.F.).

The complementary function is found from

$$(D^2-3D+2)y = 0.$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $m^2-3m+2 = 0$, or $(m-2)(m-1) = 0$, the roots are $m = 1, 2$.

$$\therefore \quad \text{C.F.} = c_1e^x+c_2e^{2x}.$$

Second step: Determination of particular integral (P.I.).

$$\text{P.I.} = \frac{1}{D^2-3D+2}(xe^{3x}+\sin 2x)$$

$$= \frac{1}{D^2-3D+2}(e^{3x}x)+\frac{1}{D^2-3D+2}\sin 2x$$

$$= e^{3x}\frac{1}{(D+3)^2-3(D+3)+2}x+\frac{D^2+2+3D}{(D^2+2)^2-9D^2}\sin 2x$$

$$= e^{3x}\frac{1}{D^2+3D+2}x+\frac{D^2+3D+2}{(-2^2+2)^2-9(-2^2)}\sin 2x$$

$$= \frac{e^{3x}}{2}\left\{1+\left(\frac{3D}{2}+\frac{D^2}{2}\right)\right\}^{-1}x+\frac{1}{40}(-4\sin 2x+6\cos 2x+2\sin 2x)$$

$$= \frac{e^{3x}}{2}\left\{1-\frac{3D}{2}-\frac{D^2}{2}+\ldots\right\}x+\frac{1}{20}(3\cos 2x-\sin 2x)$$

$$= \frac{e^{3x}}{2}\left(x-\frac{3}{2}\right)+\frac{1}{20}(3\cos 2x-\sin 2x).$$

Third step: The general solution is therefore

$$y = \text{C.F.} + \text{P.I.} = c_1e^x+c_2e^{2x}+\frac{e^{3x}}{2}\left(x-\frac{3}{2}\right)+\frac{1}{20}(3\cos 2x-\sin 2x), \text{ where } c_1, c_2 \text{ are}$$

two arbitrary constants.

Example 7: Solve $\dfrac{d^2y}{dx^2}-2\dfrac{dy}{dx}+y = xe^x\sin x.$

Solution: Let us write the given differential equation as

$$(D^2-2D+1)y = xe^x\sin x, \text{ or } (D-1)^2y = xe^x\sin x, \text{ where } D \equiv \frac{d}{dx}.$$

First step: Determination of complementary function (C.F.).

The complementary function is found from

$$(D-1)^2y = 0.$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $(m-1)^2 = 0$, the roots are $m = 1, 1$.

$$\therefore \quad \text{C.F.} = (c_1+c_2x)e^x.$$

Second step: Determination of particular integral (P.I.).

$$\text{P.I.} = \frac{1}{(D-1)^2}e^xx\sin x = e^x\frac{1}{(D+1-1)^2}x\sin x = e^x\frac{1}{D^2}x\sin x$$

$$= e^x\frac{1}{D}\int x\sin x\,dx = e^x\frac{1}{D}\{x(-\cos x)-\int 1\cdot(-\cos x)\,dx\}$$

$$= e^x\int(-x\cos x+\sin x)dx$$

$$= e^x[-\{x\sin x-\int 1\cdot\sin x\,dx\}-\cos x] = e^x(-x\sin x-\cos x-\cos x)$$

$$= -e^x(x\sin x+2\cos x).$$

Third step: The general solution is therefore

$$y = \text{C.F.} + \text{P.I.} = (c_1+c_2x)e^x-e^x(x\sin x+2\cos x), \text{ where } c_1, c_2 \text{ are two arbitrary}$$

constants.

Example 8: Solve $\dfrac{d^2y}{dx^2}-4\dfrac{dy}{dx}+4y = 8x^2e^{2x}\sin 2x.$

Solution: Let us write the given differential equation as

$(D^2-4D+4)y = 8x^2e^{2x}\sin 2x$, or $(D-2)^2y = 8x^2e^{2x}\sin 2x$,

where $D \equiv \dfrac{d}{dx}$.

First step: Determination of complementary function (C.F.).

The complementary function is found from

$$(D-2)^2y = 0.$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $(m-2)^2 = 0$, the roots are $m = 2, 2$.

$$\therefore \qquad \text{C.F.} = (c_1+c_2x)e^{2x}.$$

Second step: Determination of particular integral (P.I.)

$$\text{P.I.} = \frac{1}{(D-2)^2}(8x^2e^{2x}\sin 2x) = 8e^{2x}\frac{1}{(D+2-2)^2}(x^2\sin 2x)$$

$$= 8e^{2x}\frac{1}{D^2}(x^2\sin 2x) = 8e^{2x}\frac{1}{D}\int x^2\sin 2x\,dx$$

$$= 8e^{2x}\frac{1}{D}\left\{x^2\left(-\frac{\cos 2x}{2}\right)-\int 2x\left(-\frac{\cos 2x}{2}\right)dx\right\}$$

$$= 8e^{2x}\frac{1}{D}\left\{-\frac{x^2}{2}\cos 2x + x\frac{\sin 2x}{2}-\int 1\cdot\frac{\sin 2x}{2}dx\right\}$$

$$= 8e^{2x}\int\left(-\frac{x^2}{2}\cos 2x+\frac{x}{2}\sin 2x+\frac{\cos 2x}{4}\right)dx$$

$$= 8e^{2x}\left[\left\{-\frac{x^2}{2}\frac{\sin 2x}{2}-\int(-x)\frac{\sin 2x}{2}dx\right\}+\left\{\int\frac{x}{2}\sin 2x\,dx\right\}+\frac{\sin 2x}{8}\right]$$

$$= 8e^{2x}\left\{\left(-\frac{x^2}{4}+\frac{1}{8}\right)\sin 2x+\int x\sin 2x\,dx\right\}$$

$$= 8e^{2x}\left\{\left(\frac{1}{8}-\frac{x^2}{4}\right)\sin 2x + x\left(-\frac{\cos 2x}{2}\right)-\int 1\cdot\left(-\frac{\cos 2x}{2}\right)dx\right\}$$

$$= 8e^{2x}\left\{\left(\frac{1}{8}-\frac{x^2}{4}\right)\sin 2x-\frac{x}{2}\cos 2x+\frac{\sin 2x}{4}\right\}$$

$$= e^{2x}\{(1-2x^2)\sin 2x-4x\cos 2x+2\sin 2x\}$$

$$= e^{2x}\{(3-2x^2)\sin 2x-4x\cos 2x\}.$$

Ramanujan Central Library

Third step: The general solution is therefore

$$y = \text{C.F.} + \text{P.I.} = e^{2x}\{c_1+c_2x+(3-2x^2)\sin 2x-4x\cos 2x\},$$ where c_1, c_2 are two arbitrary constants.

Example 9: Solve $\dfrac{d^2y}{dx^2}-6\dfrac{dy}{dx}+13y = 8e^{3x}\sin 4x+2^x.$

Solution: Let us write the given differential equation as

$$(D^2-6D+13)y = 8e^{3x}\sin 4x+2^x, \quad \text{where } D \equiv \frac{d}{dx}.$$

First step: Determination of complementary function (C.F.).

The complementary function is found from

$$(D^2-6D+13)y = 0.$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $m^2-6m+13 = 0$, the roots are $m = 3 \pm 2i$.

$$\therefore \qquad \text{C.F.} = e^{3x}(c_1\cos 2x+c_2\sin 2x)$$

Second step: Determination of particular integral (P.I.)

$$\text{P.I.} = \frac{1}{D^2-6D+13}(8e^{3x}\sin 4x+2^x)$$

$$= \frac{1}{D^2-6D+13}(8e^{3x}\sin 4x)+\frac{1}{D^2-6D+13}e^{x\log 2}$$

$$= 8e^{3x}\frac{1}{(D+3)^2-6(D+3)+13}\sin 4x+\frac{e^{x\log 2}}{(\log 2)^2-6\log 2+13}$$

$$= 8e^{3x}\cdot\frac{1}{D^2+4}\sin 4x+\frac{2^x}{(\log 2)^2-6\log 2+13}$$

$$= 8e^{3x}\frac{1}{-4^2+4}\sin 4x+\frac{2^x}{(\log 2)^2-6\log 2+13}$$

$$= -\frac{2}{3}e^{3x}\sin 4x+\frac{2^x}{(\log 2)^2-6\log 2+13}.$$

Indian Institute of Engineering Science and Technology Shibpur
Ramanujan Central Library

Third step: The general solution is therefore

$$y = \text{C.F.} + \text{P.I.} = e^{3x}(c_1\cos 2x+c_2\sin 2x)-\frac{2}{3}e^{3x}\sin 4x+\frac{2^x}{(\log 2)^2-6\log 2+13},$$

where c_1, c_2 are two arbitrary constants.

Example 10: Solve $(D^2+1)y = 3\cos^2 x+2\sin^3 x.$

Solution:

First step: Determination of complementary function (C.F.).

The complementary function is found from

$$(D^2+1)y = 0.$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $m^2 + 1 = 0$, the roots are $m = \pm i$.

$$\therefore \qquad \text{C.F.} = c_1 \cos x + c_2 \sin x.$$

Second step: Determination of particular integral (P.I.)

Now, $\qquad 3\cos^2 x = \frac{3}{2}(1+\cos 2x).$

$$\sin 3x = 3\sin x - 4\sin^3 x. \quad \therefore\ 2\sin^3 x = \frac{1}{2}(3\sin x - \sin 3x).$$

$$\therefore \qquad \text{P.I.} = \frac{1}{D^2+1}(3\cos^2 x + 2\sin^3 x)$$

$$= \frac{3}{2}\frac{1}{D^2+1}(1+\cos 2x) + \frac{1}{2}\cdot\frac{1}{D^2+1}(3\sin x - \sin 3x)$$

$$= \frac{3}{2}(1+D^2)^{-1}1 + \frac{3}{2}\cdot\frac{1}{D^2+1}\cos 2x + \frac{3}{2}\frac{1}{D^2+1}\sin x - \frac{1}{2}\frac{1}{D^2+1}\sin 3x$$

$$= \frac{3}{2}(1-D^2+\ldots)1 + \frac{3}{2}\cdot\frac{1}{-2^2+1}\cos 2x + \frac{3}{2}\cdot\frac{1}{D^2+1}\sin x - \frac{1}{2}\cdot\frac{\sin 3x}{-3^2+1}$$

$$= \frac{3}{2} - \frac{1}{2}\cos 2x + \frac{3}{2}\frac{1}{D^2+1}\sin x + \frac{1}{16}\sin 3x \qquad \ldots(1)$$

In $\frac{1}{D^2+1}\sin x$, if we replace D^2 by -1^2, the method fails.

$$\frac{1}{D^2+1}(\cos x + i\sin x) = \frac{1}{D^2+1}e^{ix} = e^{ix}\frac{1}{(D+i)^2+1}1 = e^{ix}\frac{1}{D^2+2iD}1.$$

$$= e^{ix}\frac{1}{2iD}\left(1+\frac{D}{2i}\right)^{-1}1 = e^{ix}\frac{1}{2iD}\left(1-\frac{D}{2i}+\ldots\right)1 = \frac{e^{ix}}{2i}\frac{1}{D}1$$

$$= \frac{e^{ix}}{2i}x = \frac{x(\cos x + i\sin x)}{2i} = \frac{1}{2}x\sin x - i\frac{x}{2}\cos x$$

$$\therefore \qquad \frac{1}{D^2+1}(\cos x + i\sin x) = \frac{1}{2}x\sin x - i\frac{x}{2}\cos x.$$

Equating imaginary part from both sides, we get

$$\frac{1}{D^2+1}\sin x = -\frac{x}{2}\cos x.$$

Therefore, from (1), we have

$$\text{P.I.} = \frac{3}{2} - \frac{1}{2}\cos 2x - \frac{3}{4}x\cos x + \frac{1}{16}\sin 3x.$$

Third step: The general solution is therefore

$$y = \text{C.F.} + \text{P.I.} = c_1\cos x + c_2\sin x + \frac{3}{2} - \frac{1}{2}\cos 2x - \frac{3}{4}x\cos x + \frac{1}{16}\sin 3x,$$

where c_1, c_2 are two arbitrary constants.

Example 11: Solve $\frac{d^2y}{dx^2} + a^2 y = \tan ax \ (a \neq 0)$

with the symbolic operator D, where $D \equiv \frac{d}{dx}$.

Solution: The given equation can be written as $(D^2 + a^2)\, y = \tan ax$, or $(D+ia)(D-ia)y = \tan ax$, where $D \equiv \frac{d}{dx}$.

First step: Determination of complementary function (C.F.).

The complementary function is found from

$$(D+ia)(D-ia)y = 0.$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $(m+ia)(m-ia) = 0$, the roots are $m = \pm ia$.

$$\therefore \qquad \text{C.F.} = c_1\cos ax + c_2\sin ax.$$

Second step: Determination of particular integral (P.I.).

$$\text{P.I.} = \frac{1}{(D+ia)(D-ia)}\tan ax = \frac{1}{2ia}\left(\frac{1}{D-ia} - \frac{1}{D+ia}\right)\tan ax \qquad \ldots(1)$$

Now, $\qquad \frac{1}{D-ia}\tan ax = e^{iax}\int e^{-iax}\tan ax\, dx \qquad \left[\because \frac{1}{D-a}X = e^{ax}\int Xe^{-ax}dx\right]$

$$= e^{iax}\int(\cos ax - i\sin ax)\tan ax\, dx \qquad [\because e^{i\theta} = \cos\theta + i\sin\theta]$$

$$= e^{iax}\left[\int \sin ax\, dx - i\int\frac{\sin^2 ax}{\cos ax}dx\right]$$

$$= e^{iax}\left[-\frac{1}{a}\cos ax - i\int\frac{(1-\cos^2 ax)}{\cos ax}dx\right]$$

$$= e^{iax}\left[-\frac{1}{a}\cos ax - i\int\sec ax\, dx + i\int\cos ax\, dx\right]$$

$$= \frac{e^{iax}}{a}(-\cos ax - i\log|\sec ax + \tan ax| + i\sin ax)$$

$$= \frac{1}{a}(\cos ax + i\sin ax)\{-\cos ax - i(\log|\sec ax + \tan ax| - \sin ax)\}$$

$$= \frac{1}{a}(\sin ax\log|\sec ax + \tan ax| - 1) - \frac{i}{a}\cos ax\log|\sec ax + \tan ax|$$

then u is a function of four variables x_1, x_2, x_3 and x_4.

Similarly (replacing i by $-i$), we get

$$\frac{1}{D+ia}\tan ax = \frac{1}{a}(\sin ax \log|\sec ax + \tan ax| - 1) + \frac{i}{a}\cos ax \log|\sec ax + \tan ax|$$

Therefore, from (1):

$$\text{P.I.} = \frac{1}{2ia}\left(\frac{1}{D-ia}\tan ax - \frac{1}{D+ia}\tan ax\right)$$

$$= -\frac{1}{a^2}\cos ax \log|\sec ax + \tan ax|.$$

Third step: The general solution is therefore

$$y = \text{C.F.} + \text{P.I.} = c_1 \cos ax + c_2 \sin ax - \frac{1}{a^2}\cos ax \log|\sec ax + \tan ax|,$$

where c_1, c_2 are two arbitrary constants.

2.6 METHOD OF VARIATION OF PARAMETERS

It is a more powerful method of finding a particular solution of any linear non-homogeneous differential equation of second order even with variable coefficients also provided its complementary function is known. We state and prove the result in the following theorem.

Theorem

Consider the linear differential equation of second order of the form

$$\frac{d^2y}{dx^2} + P_1\frac{dy}{dx} + P_2 y = X \qquad ...(2.24)$$

where P_1, P_2 and X are functions of x or constants, then

$$\text{P.I.} = -y_1\int\frac{y_2 X}{W}dx + y_2\int\frac{y_1 X}{W}dx,$$

where y_1 and y_2 are two linearly independent solutions of

$$\frac{d^2y}{dx^2} + P_1\frac{dy}{dx} + P_2 y = 0 \qquad ...(2.25)$$

and $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$ is called the Wronskian of y_1, y_2.

Proof: Let the general solution of (2.25) is

$$y = c_1 y_1 + c_2 y_2$$

where c_1, c_2 are two arbitrary constants and y_1, y_2 are two linearly independent solutions of (2.25).

Let us assume that P.I. of (2.24) is

$$y_p = u_1 y_1 + u_2 y_2 \qquad ....(2.26)$$

where u_1, u_2 are unknown functions of x.

Differentiating both sides of (2.26) w.r.t. x, we get

$$y_p' = u_1 y_1' + u_2 y_2' + u_1' y_1 + u_2' y_2 \qquad ...(2.27)$$

Let us choose u_1, u_2 in such a manner that

$$u_1' y_1 + u_2' y_2 = 0 \qquad ...(2.28)$$

Then (2.27) becomes

$$y_p' = u_1 y_1' + u_2 y_2' \qquad ...(2.29)$$

$$\therefore \quad y_p'' = u_1 y_1'' + u_2 y_2'' + u_1' y_1' + u_2' y_2' \qquad ...(2.30)$$

Putting these values of y_p, y_p', y_p'' in (2.24) and rearranging, we get

$$u_1(y_1'' + P_1 y_1' + P_2 y_1) + u_2(y_2'' + P_1 y_2' + P_2 y_2) + u_1' y_1' + u_2' y_2' = X$$

or

$$u_1' y_1' + u_2' y_2' = X \quad [\because y_1, y_2 \text{ are solutions of (2.25)}] \qquad ...(2.31)$$

Solving (2.28) and (2.31), we get

$$u_1' = \frac{\begin{vmatrix} 0 & y_2 \\ X & y_2' \end{vmatrix}}{\begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}} = \frac{-y_2 X}{W} \quad \text{and} \quad u_2' = \frac{\begin{vmatrix} y_1 & 0 \\ y_1' & X \end{vmatrix}}{\begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}} = \frac{y_1 X}{W}$$

Here $W(y_1, y_2) = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} \neq 0$ because y_1, y_2 are linearly independent functions.

Integrating we get,

$$u_1 = -\int\frac{y_2 X}{W}dx \quad \text{and} \quad u_2 = \int\frac{y_1 X}{W}dx$$

Putting these values of u_1, u_2 in (2.26), we get

$$\text{P.I.} = -y_1\int\frac{y_2 X}{W}dx + y_2\int\frac{y_1 X}{W}dx.$$

ILLUSTRATIVE EXAMPLES

Example 1: Apply the method of variation of parameters to solve

$$\frac{d^2y}{dx^2} + 4y = \tan 2x.$$

Solution: Let us write the given differential equation as $(D^2+4)y = \tan 2x$, where $D \equiv \frac{d}{dx}$.

First step: Determination of complementary function (C.F.).

The complementary function is found from

$$(D^2+4)y = 0.$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $m^2 + 4 = 0$, the roots are $m = \pm 2i$.

$$\therefore \qquad \text{C.F.} = c_1 \cos 2x + c_2 \sin 2x.$$

Second step: Determination of particular integral (P.I.).

Let $y_1 = \cos 2x$, $y_2 = \sin 2x$ and $X = \tan 2x$.

$$\therefore \qquad W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \begin{vmatrix} \cos 2x & \sin 2x \\ -2\sin 2x & 2\cos 2x \end{vmatrix} = 2 \neq 0.$$

Hence y_1, y_2 are two linearly independent solutions of the corresponding homogeneous equation of the given differential equation.

$$\therefore \qquad \text{P.I.} = -y_1 \int \frac{y_2 X}{W} dx + y_2 \int \frac{y_1 X}{W} dx$$

$$= -\cos 2x \int \frac{\sin 2x \tan 2x}{2} dx + \sin 2x \int \frac{\cos 2x \tan 2x}{2} dx$$

$$= -\frac{1}{2}\cos 2x \int \frac{1-\cos^2 2x}{\cos 2x} dx + \frac{1}{2}\sin 2x \int \sin 2x dx$$

$$= -\frac{1}{2}\cos 2x \left\{ \int \sec 2x\, dx - \int \cos 2x\, dx \right\} - \frac{1}{4}\sin 2x \cos 2x$$

$$= -\frac{1}{4}\cos 2x \{\log(\sec 2x + \tan 2x) - \sin 2x\} - \frac{1}{4}\sin 2x \cos 2x$$

$$= -\frac{1}{4}\cos 2x \log(\sec 2x + \tan 2x).$$

Third step: The general solution is therefore

$$y = \text{C.F.} + \text{P.I.} = c_1 \cos 2x + c_2 \sin 2x - \frac{1}{4}\cos 2x \log(\sec 2x + \tan 2x),$$

where c_1, c_2 are two arbitrary constants.

Example 2: Solve by the method of variation of parameters

$$\frac{d^2y}{dx^2} + 2\frac{dy}{dx} + y = \frac{e^{-x}}{x^2}.$$

Solution: Let us write the given differential equation as

$$(D^2 + 2D + 1)y = \frac{e^{-x}}{x^2}, \quad \text{or} \quad (D+1)^2 y = \frac{e^{-x}}{x^2}, \text{ where } D \equiv \frac{d}{dx}.$$

First step: Determination of complementary function (C.F.).

The complementary function is found from

$$(D+1)^2 y = 0.$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $(m+1)^2 = 0$, the roots are $m = -1, -1$.

$$\therefore \qquad \text{C.F.} = (c_1 + c_2 x)e^{-x} = c_1 e^{-x} + c_2 x e^{-x}.$$

Second step: Determination of particular integral (P.I.)

Let $y_1 = e^{-x}$, $y_2 = xe^{-x}$ and $X = \dfrac{e^{-x}}{x^2}$.

$$\therefore \qquad W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \begin{vmatrix} e^{-x} & xe^{-x} \\ -e^{-x} & (1-x)e^{-x} \end{vmatrix} = e^{-2x} \neq 0.$$

Hence y_1, y_2 are two linearly independent solutions of the corresponding homogeneous equation of the given differential equation.

$$\therefore \qquad \text{P.I.} = -y_1 \int \frac{y_2 X}{W} dx + y_2 \int \frac{y_1 X}{W} dx$$

$$= -e^{-x} \int \frac{xe^{-x}}{e^{-2x}} \cdot \frac{e^{-x}}{x^2} dx + xe^{-x} \int \frac{e^{-x}}{e^{-2x}} \cdot \frac{e^{-x}}{x^2} dx$$

$$= -e^{-x} \int \frac{dx}{x} + xe^{-x} \int x^{-2} dx$$

$$= -e^{-x} \log x + xe^{-x}\left(-\frac{1}{x}\right) = -(1 + \log x)e^{-x}.$$

Third step: The general solution is therefore

$$y = \text{C.F.} + \text{P.I.} = (c_1 + c_2 x)e^{-x} - (1 + \log x)e^{-x},$$

where c_1, c_2 are two arbitrary constants.

Example 3: Solve by the method of variation of parameters:

$$\frac{d^2y}{dx^2} + a^2 y = \sec ax \qquad \text{(W.B.U.T. 2010)}$$

Solution: Let us write the given differential equation as

$$(D^2 + a^2)y = \sec ax, \text{ where } D \equiv \frac{d}{dx}.$$

First step: Determination of complementary function (C.F.).

The complementary function is found from

$$(D^2 + a^2)y = 0.$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $m^2 + a^2 = 0$, the roots ar $m = \pm ia$.

$$\therefore \qquad \text{C.F.} = c_1 \cos ax + c_2 \sin ax.$$

Second step: Determination of particular integral (P.I.)

Let $y_1 = \cos ax$, $y_2 = \sin ax$ and $X = \sec ax$.

$$\therefore \qquad W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \begin{vmatrix} \cos ax & \sin ax \\ -a\sin ax & a\cos ax \end{vmatrix} = a \neq 0.$$

Hence y_1, y_2 are two linearly independent solutions of the corresponding homogeneous equation of the given differential equation.

$$\therefore \quad \text{P.I.} = -y_1\int \frac{y_2 X}{W}dx + y_2\int \frac{y_1 X}{W}dx$$

$$= -\cos ax\int \frac{\sin ax \sec ax}{a}dx + \sin ax\int \frac{\cos ax \sec ax}{a}dx$$

$$= -\frac{1}{a^2}\cos ax \log(\sec ax) + \frac{1}{a}x\sin ax.$$

Third step: The general solution is therefore

$$y = \text{C.F.} + \text{P.I.} = c_1\cos ax + c_2\sin ax - \frac{1}{a^2}\cos ax\log(\sec ax) + \frac{1}{a}x\sin ax \text{ (provided } a \neq 0),$$

where c_1, c_2 are two arbitrary constants.

Example 4: Apply the method of variation of parameters to solve the equation

$$\frac{d^2y}{dx^2} + y = \sec^3 x\tan x \qquad \textit{(W.B.U.T. 2007, 2012)}$$

Solution: Let us write the given differential equation as

$$(D^2+1)y = \sec^3 x\tan x, \text{ where } D \equiv \frac{d}{dx}.$$

First step: Determination of complementary function (C.F.).

The complementary function is found from

$$(D^2+1)y = 0.$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $m^2 + 1 = 0$, the roots are $m = \pm i$.

$$\therefore \quad \text{C.F.} = c_1\cos x + c_2\sin x.$$

Second step: Determination of particular integral (P.I.)

Let $y_1 = \cos x, \quad y_2 = \sin x$ and $X = \sec^3 x\tan x$.

$$\therefore \quad W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \begin{vmatrix} \cos x & \sin x \\ -\sin x & \cos x \end{vmatrix} = 1 \neq 0.$$

Hence y_1, y_2 are two linearly independent solutions of the corresponding homogeneous equation of the given differential equation.

$$\therefore \quad \text{P.I.} = -y_1\int \frac{y_2 X}{W}dx + y_2\int \frac{y_1 X}{W}dx$$

$$= -\cos x\int \sin x\sec^3 x\tan x\,dx + \sin x\int \cos x\sec^3 x\tan x\,dx$$

$$= -\cos x\int \sec^2 x\tan^2 x\,dx + \sin x\int \sec^2 x\tan x\,dx$$

$$= -\frac{1}{3}\cos x\tan^3 x + \frac{1}{2}\sin x\tan^2 x$$

$$= -\frac{1}{3}\sin x\tan^2 x + \frac{1}{2}\sin x\tan^2 x = \frac{1}{6}\sin x\tan^2 x.$$

Third step: The general solution is therefore

$$y = \text{C.F.} + \text{P.I.} = c_1\cos x + c_2\sin x + \frac{1}{6}\sin x\tan^2 x,$$

where c_1, c_2 are two arbitrary constants.

Example 5: Apply the method of variation of parameters to solve

$$\frac{d^2y}{dx^2} + 4y = 4\sec^2 2x \qquad \textit{(W.B.U.T. 2006)}$$

Solution: Let us write the given differential equation as

$$(D^2+4)y = 4\sec^2 2x, \text{ where } D \equiv \frac{d}{dx}.$$

First step: Determination of complementary function (C.F.).

The complementary function is found from

$$(D^2+4)y = 0.$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $m^2 + 4 = 0$, the roots are $m = \pm 2i$.

$$\therefore \quad \text{C.F.} = c_1\cos 2x + c_2\sin 2x.$$

Second step: Determination of particular integral (P.I.)

Let $y_1 = \cos 2x, \quad y_2 = \sin 2x$ and $X = 4\sec^2 2x$.

$$\therefore \quad W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \begin{vmatrix} \cos 2x & \sin 2x \\ -2\sin 2x & 2\cos 2x \end{vmatrix} = 2 \neq 0.$$

Hence y_1, y_2 are two linearly independent solutions of the corresponding homogeneous equation of the given differential equation.

$$\therefore \quad \text{P.I.} = -y_1\int \frac{y_2 X}{W}dx + y_2\int \frac{y_1 X}{W}dx$$

$$= -\cos 2x\int \frac{1}{2}(\sin 2x)4\sec^2 2x\,dx + \sin 2x\int \frac{1}{2}(\cos 2x)4\sec^2 2x\,dx.$$

$$= -\cos 2x\int 2\sec 2x\tan 2x\,dx + \sin 2x\int 2\sec 2x\,dx$$

$$= -\cos 2x\sec 2x + \sin 2x\log(\sec 2x + \tan 2x).$$

$$= -1 + \sin 2x\log(\sec 2x + \tan 2x).$$

Third step: The general solution is therefore

$$y = \text{C.F.} + \text{P.I.} = c_1\cos 2x + c_2\sin 2x - 1 + \sin 2x\log(\sec 2x + \tan 2x),$$

where c_1, c_2 are two arbitrary constants.

Example 6: Solve by the method of variation of parameters

$$\frac{d^2y}{dx^2} - 3\frac{dy}{dx} + 2y = 9e^x \qquad \text{(W.B.U.T. 2004)}$$

Solution: Let us write the given differential equation as

$$(D^2 - 3D + 2)y = 9e^x, \quad \text{where } D \equiv \frac{d}{dx}$$

First step: Determination of complementary function (C.F.).

The complementary function is found from

$$(D^2 - 3D + 2)y = 0.$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $m^2 - 3m + 2 = 0$, or $(m-2)(m-1) = 0$, the roots are $m = 1, 2$.

$$\therefore \qquad \text{C.F.} = c_1e^x + c_2e^{2x}.$$

Second step: Determination of particular integral (P.I.)

Let $y_1 = e^x$, $y_2 = e^{2x}$ and $X = 9e^x$.

$$\therefore \qquad W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \begin{vmatrix} e^x & e^{2x} \\ e^x & 2e^{2x} \end{vmatrix} = e^{3x} \neq 0.$$

Hence y_1, y_2 are two linearly independent solutions of the corresponding homogeneous equation of the given differential equation.

$$\therefore \qquad \text{P.I.} = -y_1\int \frac{y_2X}{W}dx + y_2\int \frac{y_1X}{W}dx$$

$$= -e^x\int \frac{e^{2x}9e^x}{e^{3x}}dx + e^{2x}\int \frac{e^x9e^x}{e^{3x}}dx$$

$$= -e^x\int 9dx + e^{2x}\int 9e^{-x}dx$$

$$= -9xe^x - 9e^{2x}\cdot e^{-x} = -9(x+1)e^x.$$

Third step: The general solution is therefore

$$y = \text{C.F.} + \text{P.I.} = c_1e^x + c_2e^{2x} - 9(x+1)e^x,$$

where c_1, c_2 are two arbitrary constants.

Example 7: Solve by the method of variation of parameters

$$(D^2 - 3D + 2)y = \frac{e^x}{1+e^x}, \quad \text{where } D \equiv \frac{d}{dx}.$$

Solution:

First step: Determination of complementary function (C.F.).

The complementary function is found from

$$(D^2 - 3D + 2)y = 0.$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $m^2 - 3m + 2 = 0$, or $(m-2)(m-1) = 0$, the roots are $m = 1, 2$.

$$\therefore \qquad \text{C.F.} = c_1e^x + c_2e^{2x}.$$

Second step: Determination of particular integral (P.I.)

Let $y_1 = e^x$, $y_2 = e^{2x}$ and $X = \dfrac{e^x}{1+e^x}$

$$\therefore \qquad W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \begin{vmatrix} e^x & e^{2x} \\ e^x & 2e^{2x} \end{vmatrix} = e^{3x} \neq 0.$$

Hence y_1, y_2 are two linearly independent solutions of the corresponding homogeneous equation of the given differential equation.

$$\therefore \qquad \text{P.I.} = -y_1\int \frac{y_2X}{W}dx + y_2\int \frac{y_1X}{W}dx$$

$$= -e^x\int \frac{e^{2x}}{e^{3x}}\cdot\frac{e^x}{(1+e^x)}dx + e^{2x}\int \frac{e^x}{e^{3x}}\cdot\frac{e^x}{(1+e^x)}dx$$

$$= -e^x\int \frac{dx}{1+e^x} + e^{2x}\int \frac{dx}{e^x(1+e^x)}$$

$$= -e^x\int \frac{dx}{1+e^x} + e^{2x}\int \frac{(1+e^x)-e^x}{e^x(1+e^x)}dx$$

$$= -(e^x + e^{2x})\int \frac{dx}{1+e^x} + e^{2x}\int e^{-x}dx$$

$$= -(e^x + e^{2x})\int \frac{e^{-x}}{e^{-x}+1}dx - e^{2x}\cdot e^{-x}$$

$$= (e^x + e^{2x})\log(e^{-x}+1) - e^x.$$

Third step: The general solution is therefore

$$y = \text{C.F.} + \text{P.I.} = c_1e^x + c_2e^{2x} + (e^x + e^{2x})\log(e^{-x}+1) - e^x,$$

where c_1, c_2 are two arbitrary constants.

Example 8: Solve by the method of variation of parameters

$$(D^2 - 2D + 1)y = e^x\log x, \quad \text{where } D \equiv \frac{d}{dx}.$$

Solution:

First step: Determination of complementary function (C.F.).

The complementary function is found from

$$(D^2 - 2D + 1)y = 0.$$

Let $y = e^{mx}$ be a trial solution, then the auxiliary equation becomes $m^2 - 2m + 1 = 0$, or $(m-1)^2 = 0$ the roots are $m = 1, 1$.

$$\therefore \qquad \text{C.F.} = (c_1 + c_2x)e^x = c_1e^x + c_2xe^x.$$

Second step: Determination of particular integral (P.I.)

Let $y_1 = e^x$, $y_2 = xe^x$ and $X = e^x\log x$.

$$\therefore \qquad W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \begin{vmatrix} e^x & xe^x \\ e^x & (1+x)e^x \end{vmatrix} = e^{2x} \neq 0.$$

Hence y_1, y_2 are two linearly independent solutions of the corresponding homogeneous equation of the given differential equation.

$$\therefore \qquad \text{P.I.} = -y_1 \int \frac{y_2 X}{W} dx + y_2 \int \frac{y_1 X}{W} dx$$

$$= -e^x \int \frac{xe^x}{e^{2x}} e^x \log x \, dx + xe^x \int \frac{e^x}{e^{2x}} e^x \log x \, dx$$

$$= -e^x \int x \log x \, dx + xe^x \int \log x \, dx$$

$$= -e^x \left(\frac{x^2}{2} \log x - \int \frac{1}{x} \cdot \frac{x^2}{2} dx \right) + xe^x \left(x \log x - \int \frac{1}{x} \cdot x \, dx \right)$$

$$= -e^x \left(\frac{x^2}{2} \log x - \frac{x^2}{4} \right) + xe^x (x \log x - x)$$

$$= \frac{1}{4} x^2 e^x (1 - 2\log x + 4 \log x - 4) = \frac{1}{4} x^2 e^x (2 \log x - 3).$$

Third step: The general solution is therefore

$$y = \text{C.F.} + \text{P.I.} = (c_1 + c_2 x)e^x + \frac{1}{4} x^2 e^x (2 \log x - 3),$$

where c_1, c_2 are two arbitrary constants.

Example 9: Find the general solution of

$$(1+x)\frac{d^2y}{dx^2} + x\frac{dy}{dx} - y = (1+x)^2$$

by the method of variation of parameters, it is given that $y = x$ and $y = e^{-x}$ are two linearly independent solutions of the corresponding homogeneous equation.

Solution: Let us write the given equation as

$$\frac{d^2y}{dx^2} + \frac{x}{1+x}\frac{dy}{dx} - \frac{1}{1+x} y = 1 + x.$$

It is a second order linear differential equation and it is given that x, e^{-x} are two linearly independent solutions of the corresponding homogeneous equation.

$$\therefore \qquad \text{C.F.} = c_1 x + c_2 e^{-x}$$

Let $y_1 = x, \quad y_2 = e^{-x}$ and $X = 1 + x$.

$$\therefore \qquad W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \begin{vmatrix} x & e^{-x} \\ 1 & -e^{-x} \end{vmatrix} = -xe^{-x} - e^{-x} = -(x+1)e^{-x} \neq 0.$$

since y_1, y_2 are linearly independent functions according to the question.

$$\therefore \qquad \text{P.I.} = -y_1 \int \frac{y_2 X}{W} dx + y_2 \int \frac{y_1 X}{W} dx$$

$$= -x \int \frac{e^{-x}(1+x)}{-(x+1)e^{-x}} dx + e^{-x} \int \frac{x(1+x)}{-(x+1)e^{-x}} dx$$

$$= x \int dx - e^{-x} \int xe^x dx = x^2 - e^{-x} \left\{ x \int e^x dx - \int \left(\frac{dx}{dx} \right) \left(\int e^x dx \right) dx \right\}$$

$$= x^2 - e^{-x}(xe^x - e^x) = x^2 - x + 1.$$

Therefore, the general solution is

$$y = \text{C.F.} + \text{P.I.} = c_1 x + c_2 e^{-x} + x^2 - x + 1,$$

where c_1, c_2 are two arbitrary constants.

Example 10: Solve by the method of variation of parameters

$$\frac{d^2y}{dx^2} + \frac{1}{x}\frac{dy}{dx} - \frac{1}{x^2} y = \log x \; (x > 0)$$

it is given that $y = x$ and $y = \frac{1}{x}$ are two linearly independent solutions of the corresponding homogeneous differential equation.

Solution: Here $x, \frac{1}{x}$ are two linearly independent solutions of the corresponding homogeneous equation of the given differential equation.

$$\therefore \qquad \text{C.F.} = c_1 x + \frac{c_2}{x}.$$

Let $y_1 = x, \quad y_2 = \frac{1}{x}$ and $X = \log x$.

$$\therefore \qquad W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \begin{vmatrix} x & \frac{1}{x} \\ 1 & -\frac{1}{x^2} \end{vmatrix} = -\frac{2}{x} \neq 0.$$

$$\therefore \qquad \text{P.I.} = -y_1 \int \frac{y_2 X}{W} dx + y_2 \int \frac{y_1 X}{W} dx$$

$$= -x \int \frac{1}{x} \left(-\frac{x}{2} \right) \log x \, dx + \frac{1}{x} \int x \left(-\frac{x}{2} \right) \log x \, dx$$

$$= \frac{x}{2} \int \log x \, dx - \frac{1}{2x} \int x^2 \log x \, dx$$

$$= \frac{x}{2} \int \left(x \log x - \int \frac{1}{x} \cdot x \, dx \right) - \frac{1}{2x} \left(\frac{x^3}{3} \log x - \int \frac{1}{x} \cdot \frac{x^3}{3} dx \right)$$

$$= \frac{x^2}{2} (\log x - 1) - \frac{1}{2x} \left(\frac{x^3}{3} \log x - \frac{x^3}{9} \right)$$

$$= \frac{x^2}{2}\left(\log x - 1 - \frac{1}{3}\log x + \frac{1}{9}\right) = \frac{x^2}{2}\left(\frac{2}{3}\log x - \frac{8}{9}\right)$$

$$= x^2\left(\frac{1}{3}\log x - \frac{4}{9}\right)$$

Therefore, the general solution is

$$y = \text{C.F.} + \text{P.I.} = c_1 x + \frac{c_2}{x} + x^2\left(\frac{1}{3}\log x - \frac{4}{9}\right)$$

where c_1, c_2 are two arbitrary constants.

2.7 CAUCHY-EULER EQUATION

An equation of the form

$$x^n \frac{d^n y}{dx^n} + P_1 x^{n-1}\frac{d^{n-1}y}{dx^{n-1}} + P_2 x^{n-2}\frac{d^{n-2}y}{dx^{n-2}} + \ldots + P_{n-1}x\frac{dy}{dx} + P_n y = X \qquad \ldots(2.32)$$

where $P_1, P_2, \ldots, P_n$ are constants and X is a function of x only, is known as Euler's homogeneous linear equation or, Cauchy-Euler equation. It can be transformed into linear differential equation with constant coefficients by the substitution $x = e^z$, or $z = \log x$.

Cauchy-Euler Equation of Second Order

The general form of Cauchy-Euler equation of second order is

$$x^2\frac{d^2y}{dx^2} + P_1 x\frac{dy}{dx} + P_2 y = X, \text{ or } (x^2D^2 + P_1 xD + P_2)y = X \qquad \ldots(2.33)$$

where $D \equiv \frac{d}{dx}$, P_1, P_2 are constants and X is a function of x only.

Let us substitute $x = e^z$, or $z = \log x$.

$$\therefore \quad \frac{dy}{dx} = \frac{dy}{dz}\cdot\frac{dz}{dx} = \frac{1}{x}\frac{dy}{dz}, \text{ or } x\frac{dy}{dx} = \frac{dy}{dz}.$$

$$\therefore \quad xDy = D'y, \quad \text{where } D \equiv \frac{d}{dx}, \quad D' \equiv \frac{d}{dz} \qquad \ldots(2.34)$$

Again differentiating both sides of $x\frac{dy}{dx} = \frac{dy}{dz}$ w.r.t. x, we get

$$\frac{dy}{dx} + x\frac{d^2y}{dx^2} = \frac{d}{dx}\left(\frac{dy}{dz}\right) = \frac{d}{dz}\left(\frac{dy}{dz}\right)\cdot\frac{dz}{dx} = \frac{1}{x}\frac{d^2y}{dz^2}.$$

$$\therefore \quad x\frac{dy}{dx} + x^2\frac{d^2y}{dx^2} = \frac{d^2y}{dz^2}$$

or

$$x^2\frac{d^2y}{dx^2} = \frac{d^2y}{dz^2} - x\frac{dy}{dx}, \quad \text{or} \quad x^2D^2y = D'^2y - D'y$$

$$\therefore \quad x^2D^2y = D'(D'-1)y, \text{ where } D \equiv \frac{d}{dx}, \ D' \equiv \frac{d}{dz} \qquad \ldots(2.35)$$

By (2.34), (2.35), the given equation (2.33) is converted into a linear equation with constant coefficients which can be solved by the previous methods.

Note: By applying the same procedures as in (2.34) and (2.35) it can be proved that

$$x^r\frac{d^ry}{dx^r} = D'(D'-1)(D'-2)\ldots(D'-r+1)y, \text{ where } D' \equiv \frac{d}{dz}, \ r = 1, 2, \ldots, n$$

for the differential equation (2.32).

ILLUSTRATIVE EXAMPLES

Example 1: Solve: (*i*) $x^2\frac{d^2y}{dx^2} - 2x\frac{dy}{dx} - 4y = x^4$. (*ii*) $x^2\frac{d^2y}{dx^2} - 3x\frac{dy}{dx} + 4y = 2x^2$ (*M–201/13*)

Solution: (*i*) Let us write the given equation as

$$(x^2D^2 - 2xD - 4)y = x^4, \text{ where } D \equiv \frac{d}{dx}. \qquad \ldots(1)$$

Putting $x = e^z$, *i.e.*, $z = \log x$, we have

$$xDy = D'y, x^2D^2y = D'(D'-1)y, \text{ where } D' \equiv \frac{d}{dz}$$

Therefore equation (1) reduces to

$$[D'(D'-1) - 2D' - 4]y = e^{4z}$$

or

$$(D'^2 - 3D' - 4)y = e^{4z}, \text{ where } D' \equiv \frac{d}{dz} \qquad \ldots(2)$$

First step: The complementary function is found from

$$(D'^2 - 3D' - 4)y = 0.$$

Let $y = e^{mz}$ be a trial solution, then the auxiliary equation becomes $m^2 - 3m - 4 = 0$, or $(m-4)(m+1) = 0$, the roots are $m = 4, -1$.

$$\therefore \quad \text{C.F.} = c_1e^{4z} + c_2e^{-z}$$

Second step: Determination of particular integral (P.I.)

$$\text{P.I.} = \frac{1}{D'^2 - 3D' - 4}e^{4z} = \frac{1}{(D'-4)(D'+1)}e^{4z} = \frac{1}{D'-4}\cdot\frac{1}{4+1}e^{4z}$$

$$= \frac{e^{4z}}{5}\cdot\frac{1}{(D'+4)-4}1 = \frac{e^{4z}}{5}\cdot\frac{1}{D'}1 = \frac{e^{4z}}{5}\cdot z$$

Third step: The general solution is therefore

$$y = \text{C.F.} + \text{P.I.} = c_1e^{4z} + c_2e^{-z} + \frac{ze^{4z}}{5}$$

$$= c_1x^4 + \frac{c_2}{x} + \frac{1}{5}x^4\log x \qquad [\because e^z = x]$$

where c_1, c_2 are two arbitrary constants.

(*ii*) Solve: $x^2\dfrac{d^2y}{dx^2}-3x\dfrac{dy}{dx}+4y=2x^2$. (*M–201/13*)

Let us write the given equation as

$$(x^2D^2-3xD+4)y=2x^2, \text{ where } D\equiv\frac{d}{dx} \qquad ...(1)$$

Putting $x=e^z$, *i.e.*, $z=\log x$, we have

$$xDy=D'y;\ x^2D^2y=D'(D'-1)y, \text{ where } D'\equiv\frac{d}{dz}.$$

Therefore equation (1) reduces to

$$\{D'(D'-1)-3D'+4\}y=2e^{2z}$$

$$\Rightarrow \qquad (D'-2)^2y=2e^{2z}, \text{ where } D'\equiv\frac{d}{dz}.$$

First step: The complementary function is found from: $(D'-2)^2y=0$.

Let $y=e^{mz}$ be a trial solution, then the auxiliary equation becomes $(m-2)^2=0 \Rightarrow m=2, 2$.

$$\therefore \qquad \text{C.F.}=(C_1+C_2z)e^{2z}=(C_1+C_2\log x)x^2$$

Second step: Determination of P.I.

$$\text{P.I.}=\frac{1}{(D'-2)^2}2e^{2z}=2e^{2z}\frac{1}{(D'+2-2)^2}$$

$$=2e^{2z}\cdot\frac{1}{D'^2}=e^{2z}z^2=x^2(\log x)^2$$

Third step: The general solution is given by

$$y=\text{C.F.}+\text{P.I.}=(C_1+C_2\log x)x^2+x^2(\log x)^2,$$

where C_1, C_2 are two arbitrary constants.

Example 2: Solve $x^2\dfrac{d^2y}{dx^2}+x\dfrac{dy}{dx}+y=\sin(\log x^2)$.

Solution: Let us write the given equation as

$$(x^2D^2+xD+1)y=\sin(\log x^2), \text{ where } D\equiv\frac{d}{dx}. \qquad ...(1)$$

Putting $x=e^z$, *i.e.*, $z=\log x$, we have

$$xDy=D'y,\ x^2D^2y=D'(D'-1)y, \text{ where } D'\equiv\frac{d}{dz}.$$

Therefore equation (1) reduces to

$$\{D'(D'-1)+D'+1\}y=\sin 2z,$$

or

$$(D'^2+1)y=\sin 2z,\ D'\equiv\frac{d}{dx}. \qquad ...(2)$$

First step: The complementary function is found from

$$(D'^2+1)y=0.$$

Let $y=e^{mz}$ be a trial solution, then the auxiliary equation becomes $m^2+1=0$, the roots are $m=\pm i$.

$$\therefore \qquad \text{C.F.}=c_1\cos z+c_2\sin z$$

Second step: Determination of particular integral (P.I.)

$$\text{P.I.}=\frac{1}{D'^2+1}\sin 2z=\frac{1}{-2^2+1}\sin 2z=-\frac{1}{3}\sin 2z.$$

Third step: The general solution is therefore

$$y=\text{C.F.}+\text{P.I.}=c_1\cos z+c_2\sin z-\frac{1}{3}\sin 2z$$

$$=c_1\cos(\log x)+c_2\sin(\log x)-\frac{1}{3}\sin(\log x^2) \qquad (\because z=\log x),$$

where c_1, c_2 are two arbitrary constants.

Example 3: Solve: $x^2\dfrac{d^2y}{dx^2}-x\dfrac{dy}{dx}-3y=x^2\log x$. (W.B.U.T. 2008

Solution: Let us write the given equation as

$$(x^2D^2-xD-3)y=x^2\log x, \text{ where } D\equiv\frac{d}{dx} \qquad ...($$

Putting $x=e^z$, *i.e.*, $z=\log x$, we have

$$xDy=D'y,\ x^2D^2y=D'(D'-1)y, \text{ where } D'\equiv\frac{d}{dz}.$$

Therefore equation (1) reduces to

$$\{D'(D'-1)-D'-3\}y=ze^{2z},$$

or

$$(D'^2-2D'-3)y=ze^{2z}$$

First step: The complementary function is found from

$$(D'^2-2D'-3)y=0.$$

Let $y=e^{mz}$ be a trial solution, then the auxiliary equation becomes $m^2-2m-3=0$, or $(m$ $(m+1)=0$, the roots are $m=-1, 3$.

$$\therefore \qquad \text{C.F.}=c_1e^{-z}+c_2e^{3z}.$$

Second step: Determination of particular integral (P.I.)

$$\text{P.I.}=\frac{1}{D'^2-2D'-3}ze^{2z}=e^{2z}\frac{1}{(D'+2)^2-2(D'+2)-3}z$$

$$=e^{2z}\frac{1}{D'^2+2D'-3}z=-\frac{e^{2z}}{3}\left\{1-\left(\frac{2D'}{3}+\frac{D'^2}{3}\right)\right\}^{-1}z$$

$$=-\frac{e^{2z}}{3}\left(1+\frac{2D'}{3}+\frac{1}{3}D'^2+\ldots\right)z=-\frac{1}{3}e^{2z}\left(z+\frac{2}{3}\right)$$

$$=-\frac{1}{9}e^{2z}(3z+2).$$

Third step: The general solution is therefore

$$y=\text{C.F.}+\text{P.I.}=c_1e^{-z}+c_2e^{3z}-\frac{1}{9}e^{2z}(3z+2)$$

$$=\frac{c_1}{x}+c_2x^3-\frac{1}{9}x^2(3\log x+2) \qquad [\because$$

where c_1, c_2 are two arbitrary constants.

Example 4: Solve $x^2\dfrac{d^2y}{dx^2}-x\dfrac{dy}{dx}+4y = x\sin(\log x)$. *(W.B.U.T. 2005, 2011)*

Solution: Let us write the given equation as

$$(x^2D^2-xD+4)y = x\sin(\log x), \text{ where } D\equiv\frac{d}{dx} \quad ...(1)$$

Putting $x=e^z$, *i.e.*, $z=\log x$, we have

$$xDy = D'y,\ x^2D^2y = D'(D'-1)y, \text{ where } D'\equiv\frac{d}{dz}.$$

Therefore equation (1) reduces to

$$\{D'(D'-1)-D'+4\}y = e^z\sin z,$$

or

$$(D'^2-2D'+4)y = e^z\sin z. \quad ...(2)$$

First step: The complementary function is found from

$$(D'^2-2D'+4)y = 0.$$

Let $y=e^{mz}$ be a trial solution, then the auxiliary equation becomes $m^2-2m+4=0$, the roots are $m=1\pm i\sqrt{3}$.

$$\therefore \quad \text{C.F.} = e^z(c_1\cos\sqrt{3}z+c_2\sin\sqrt{3}z)$$

Second step: Determination of particular integral (P.I.)

$$\text{P.I.} = \frac{1}{D'^2-2D'+4}(e^z\sin z) = e^z\frac{1}{(D'+1)^2-2(D'+1)+4}\sin z$$

$$= e^z\frac{1}{D'^2+3}\sin z = e^z\frac{1}{-1^2+3}\sin z = \frac{1}{2}e^z\sin z.$$

Third step: The general solution is therefore

$$y = \text{C.F.}+\text{P.I.} = e^z(c_1\cos\sqrt{3}z+c_2\sin\sqrt{3}z)+\frac{1}{2}e^z\sin z$$

$$= x\{c_1\cos(\sqrt{3}\log x)+c_2\sin(\sqrt{3}\log x)\}+\frac{1}{2}x\sin(\log x)$$

$[\because x=e^z, \text{ or } z=\log x]$

Here c_1, c_2 are two arbitrary constants.

Example 5: Solve $x^2\dfrac{d^2y}{dx^2}+x\dfrac{dy}{dx}-y = \sin(\log x)+x\cos(\log x)$ *(W.B.U.T. 2004)*

Solution: Let us write the given equation as

$$(x^2D^2+xD-1)y = \sin(\log x)+x\cos(\log x), \text{ where } D\equiv\frac{d}{dx} \quad ...(1)$$

Putting $x=e^z$, *i.e.*, $z=\log x$, we have

$$xDy = D'y,\ x^2D^2y = D'(D'-1)y, \text{ where } D'\equiv\frac{d}{dz}.$$

Therefore equation (1) reduces to

$$\{D'(D'-1)+D'-1\}y = \sin z+e^z\cos z$$

or

$$(D'^2-1)y = \sin z+e^z\cos z.$$

First step: The complementary function is found from

$$(D'^2-1)y = 0.$$

Let $y=e^{mz}$ be a trial solution, then the auxiliary equation becomes $m^2-1=0$, the roots are $m=\pm1$.

$$\therefore \quad \text{C.F.} = c_1e^z+c_2e^{-z}.$$

Second step: Determination of particular integral (P.I.)

$$\text{P.I.} = \frac{1}{D'^2-1}(\sin z+e^z\cos z) = \frac{1}{D'^2-1}\sin z+\frac{1}{D'^2-1}e^z\cos z$$

$$= \frac{1}{-1^2-1}\sin z+e^z\frac{1}{(D'+1)^2-1}\cos z = -\frac{1}{2}\sin z+e^z\frac{1}{D'^2+2D'}\cos z$$

$$= -\frac{1}{2}\sin z+e^z\frac{D'^2-2D'}{D'^4-4D'^2}\cos z$$

$$= -\frac{1}{2}\sin z+e^z\frac{D'^2-2D'}{(-1^2)^2-4(-1^2)}\cos z$$

$$= -\frac{1}{2}\sin z+e^z\cdot\frac{1}{5}(-\cos z+2\sin z).$$

Third step: The general solution is therefore

$$y = \text{C.F.}+\text{P.I.} = c_1e^z+c_2e^{-z}-\frac{1}{2}\sin z+\frac{e^z}{5}(2\sin z-\cos z)$$

$$= c_1x+\frac{c_2}{x}-\frac{1}{2}\sin(\log x)+\frac{x}{5}\{2\sin(\log x)-\cos(\log x)\}$$

$[\because x=e^z, \text{ i.e., } z=\log x]$

Here c_1, c_2 are two arbitrary constants.

Example 6: Solve the differential equation

$$x^2\frac{d^2y}{dx^2}+x\frac{dy}{dx}+y = \log x\sin(\log x). \quad \textit{(W.B.U.T. 2002, 2012)}$$

Solution: Let us write the given equation as

$$(x^2D^2+xD+1)y = \log x\sin(\log x), \text{ where } D\equiv\frac{d}{dx} \quad ...(1)$$

Putting $x=e^z$, *i.e.*, $z=\log x$, we have

$$xDy = D'y,\ x^2D^2y = D'(D'-1)y, \text{ where } D'\equiv\frac{d}{dz}.$$

Therefore equation (1) reduces to

$$\{D'(D'-1)+D'+1\}y = z\sin z,$$

or

$$(D'^2+1)y = z\sin z.$$

First step: The complementary function is found from

$$(D'^2+1)y = 0.$$

Let $y = e^{mz}$ be a trial solution, then the auxiliary equation becomes $m^2 + 1 = 0$, the roots are $m = \pm i$.

$\therefore$ $$\text{C.F.} = c_1 \cos z + c_2 \sin z.$$

Second step: Determination of particular integral (P.I.)

$$\text{P.I.} = \frac{1}{D'^2+1}(z\sin z) = \left(z - \frac{2D'}{D'^2+1}\right)\frac{1}{D'^2+1}\sin z \qquad ...(2)$$

In $\frac{1}{D'^2+1}\sin z$, if we replace D'^2 by -1^2, the method fails.

Now, $$\frac{1}{D'^2+1}(\cos z + i\sin z) = \frac{1}{D'^2+1}e^{iz} = e^{iz}\frac{1}{(D'+i)^2+1}1$$

$$= e^{iz}\frac{1}{D'^2+2iD'}1 = e^{iz}\frac{1}{2iD'}\left(1+\frac{D'}{2i}\right)^{-1}1$$

$$= e^{iz}\frac{1}{2iD'}\left(1-\frac{D'}{2i}+\ldots\right)1 = \frac{e^{iz}}{2i}\frac{1}{D'}1 = \frac{e^{iz}}{2i}z$$

$$= \frac{z}{2i}(\cos z + i\sin z) = \frac{1}{2}z\sin z - i\frac{z}{2}\cos z.$$

$\therefore$ $$\frac{1}{D'^2+1}\cos z + i\frac{1}{D'^2+1}\sin z = \frac{1}{2}z\sin z - i\frac{z}{2}\cos z.$$

Equating real and imaginary parts from both sides, we get

$$\frac{1}{D'^2+1}\cos z = \frac{1}{2}z\sin z, \quad \frac{1}{D'^2+1}\sin z = -\frac{1}{2}z\cos z \qquad ...(3)$$

Therefore from (2), we get

$$\text{P.I.} = \left(z - \frac{2D'}{D'^2+1}\right)\left(-\frac{z}{2}\cos z\right) = -\frac{z^2}{2}\cos z + \frac{1}{D'^2+1}D'(z\cos z)$$

$$= -\frac{z^2}{2}\cos z + \frac{1}{D'^2+1}(\cos z - z\sin z)$$

$$= -\frac{z^2}{2}\cos z + \frac{1}{D'^2+1}\cos z - \frac{1}{D'^2+1}(z\sin z)$$

$$= -\frac{z^2}{2}\cos z + \frac{1}{2}z\sin z - \text{P.I.} \qquad \text{[by (2) and (3)]}$$

$\therefore$ $$2(\text{P.I.}) = -\frac{z^2}{2}\cos z + \frac{z}{2}\sin z.$$

or $$\text{P.I.} = -\frac{z^2}{4}\cos z + \frac{z}{4}\sin z.$$

Third step: The general solution is therefore

$$y = \text{C.F.} + \text{P.I.} = c_1\cos z + c_2\sin z - \frac{z^2}{4}\cos z + \frac{z}{4}\sin z$$

$$= c_1\cos(\log x) + c_2\sin(\log x) - \frac{1}{4}(\log x)^2\cos(\log x) + \frac{1}{4}(\log x)\sin(\log x)$$

$[\because x = e^z,\ i.e.,\ z = \log x]$

Here c_1, c_2 are two arbitrary constants.

Example 7: Solve $x^2\frac{d^2y}{dx^2} + 3x\frac{dy}{dx} + y = \frac{1}{(1-x)^2}$.

Solution: Let us write the given equation as

$$(x^2D^2 + 3xD + 1)y = \frac{1}{(1-x)^2}, \text{ where } D \equiv \frac{d}{dx} \qquad ...(1)$$

Putting $x = e^z$, *i.e.*, $z = \log x$, we have

$$xDy = D'y,\ x^2D^2y = D'(D'-1)y, \text{ where } D' \equiv \frac{d}{dz}.$$

Therefore, equation (1) reduces to

$$\{D'(D'-1) + 3D' + 1\}y = \frac{1}{(1-e^z)^2},$$

or $$(D'+1)^2y = \frac{1}{(1-e^z)^2}.$$

First step: The complementary function is found from

$$(D'+1)^2y = 0.$$

Let $y = e^{mz}$ be a trial solution, then the auxiliary equation becomes $(m+1)^2 = 0$, the roots are $m = -1, -1$.

$\therefore$ $$\text{C.F.} = (c_1 + c_2 z)e^{-z}.$$

Second step: Determination of particular integral (P.I.)

$$\text{P.I.} = \frac{1}{(D'+1)^2}\frac{1}{(1-e^z)^2} = \frac{1}{D'+1}\left\{\frac{1}{D'+1}\frac{1}{(1-e^z)^2}\right\} \qquad ...(2)$$

Now, $$\frac{1}{D'+1}\frac{1}{(1-e^z)^2} = e^{-z}\int\frac{e^z}{(1-e^z)^2}dz \qquad \left[\because \frac{1}{D-a}X = e^{ax}\int Xe^{-ax}dx\right]$$

$$= \frac{e^{-z}}{1-e^z}.$$

Therefore, from (2),

$$\text{P.I.} = \frac{1}{D'+1}\left(\frac{e^{-z}}{1-e^z}\right) = e^{-z}\int e^z\left(\frac{e^{-z}}{1-e^z}\right)dz = e^{-z}\int\frac{dz}{1-e^z}$$

$$= e^{-z}\int\frac{(1-e^z)+e^z}{1-e^z}dz = e^{-z}\int\left(1+\frac{e^z}{1-e^z}\right)dz$$

$$= e^{-z}\left\{\int dz + \int\frac{e^z}{1-e^z}dz\right\} = e^{-z}\{z - \log|1-e^z|\}$$

Third step: The general solution is therefore

$$y = \text{C.F.} + \text{P.I.} = (c_1 + c_2 z)e^{-z} + e^{-z}\{z - \log|1-e^z|\}$$

$$= \left\{c_1 + c_2\log x + \log\left(\frac{x}{|1-x|}\right)\right\}\frac{1}{x} \qquad [\because x = e^z, \text{ or } z = \log x]$$

where c_1, c_2 are two arbitrary constants.

Example 8: Solve $x^2\dfrac{d^2y}{dx^2}+4x\dfrac{dy}{dx}+2y=e^x$. (W.B.U.T. 2012)

Solution: Let us write the given equation as

$$(x^2D^2+4xD+2)y=e^x, \text{ where } D\equiv\frac{d}{dx} \qquad ...(1)$$

Putting $x=e^z$, *i.e.*, $z=\log x$, we have

$$xDy=D'y,\ x^2D^2y=D'(D'-1)y, \text{ where } D'\equiv\frac{d}{dz}.$$

Therefore equation (1) reduces to

$$\{D'(D'-1)+4D'+2\}y=e^{e^z},$$

or $$(D'^2+3D'+2)y=e^{e^z}.$$

First step: The complementary function is found from

$$(D'^2+3D'+2)y=0.$$

Let $y=e^{mz}$ be a trial solution, then the auxiliary equation becomes $m^2+3m+2=0$, or $(m+1)(m+2)=0$, the roots are $m=-1,-2$.

$$\therefore \quad \text{C.F.}=c_1e^{-z}+c_2e^{-2z}=\frac{c_1}{x}+\frac{c_2}{x^2} \qquad (\because e^z=x)$$

Second step: Determination of particular integral (P.I.)

$$\text{P.I.}=\frac{1}{D'^2+3D'+2}e^{e^z}=\frac{1}{(D'+1)(D'+2)}e^{e^z}=\left(\frac{1}{D'+1}-\frac{1}{D'+2}\right)e^{e^z}$$

$$=e^{-z}\int e^{e^z}e^z dz-e^{-2z}\int e^{e^z}e^{2z}dz \qquad \left[\because \frac{1}{D-a}X=e^{ax}\int Xe^{-ax}dx\right]$$

$$=\frac{1}{x}\int e^x dx-\frac{1}{x^2}\int xe^x dx \qquad [\because x=e^z,\ \therefore dx=e^z dz]$$

$$=\frac{1}{x}e^x-\frac{1}{x^2}\{xe^x-\int 1\cdot e^x dx\}=\frac{1}{x}e^x-\frac{e^x}{x^2}(x-1)$$

$$=\frac{e^x}{x^2}.$$

Third step: The general solution is therefore

$$y=\text{C.F.}+\text{P.I.}=\frac{c_1}{x}+\frac{c_2}{x^2}+\frac{e^x}{x^2},$$

where c_1, c_2 are two arbitrary constants.

Example 9: Solve $(x+a)^2\dfrac{d^2y}{dx^2}-4(x+a)\dfrac{dy}{dx}+6y=x$.

Solution: Put $x+a=u$, then

$$\frac{dy}{dx}=\frac{dy}{du}\frac{du}{dx}=\frac{dy}{du},\ \frac{d^2y}{dx^2}=\frac{d}{dx}\left(\frac{dy}{dx}\right)=\frac{d}{du}\left(\frac{dy}{du}\right)\frac{du}{dx}=\frac{d^2y}{du^2}$$

and hence the given equation reduces to

$$u^2\frac{d^2y}{du^2}-4u\frac{dy}{du}+6y=u-a$$

Let us write (1) as

$$(u^2D^2-4uD+6)y=u-a, \text{ where } D\equiv\frac{d}{du} \qquad ...(2)$$

Putting $u=e^z$, *i.e.*, $z=\log u$, we have

$$uDy=D'y,\ u^2D^2y=D'(D'-1)y, \text{ where } D'\equiv\frac{d}{dz}.$$

Therefore equation (2) changes to

$$\{D'(D'-1)-4D'+6\}y=e^z-a, \text{ or } (D'^2-5D'+6)y=e^z-a.$$

First step: The complementary function is found from

$$(D'^2-5D'+6)y=0.$$

Let $y=e^{mz}$ be a trial solution, then the auxiliary equation becomes $m^2-5m+6=0$, or $(m-3)(m-2)=0$, the roots are $m=3, 2$.

$$\therefore \quad \text{C.F.}=c_1e^{3z}+c_2e^{2z}.$$

Second step: Determination of particular integral (P.I.)

$$\text{P.I.}=\frac{1}{D'^2-5D'+6}(e^z-a)=\frac{1}{D'^2-5D'+6}e^z-\frac{1}{D'^2-5D'+6}a$$

$$=\frac{1}{1^2-5\cdot1+6}e^z-\frac{1}{6}\left\{1-\left(\frac{5}{6}D'-\frac{1}{6}D'^2\right)\right\}^{-1}a$$

$$=\frac{1}{2}e^z-\frac{1}{6}\left(1+\frac{5}{6}D'-\frac{1}{6}D'^2+\ldots\right)a$$

$$=\frac{1}{2}e^z-\frac{1}{6}a.$$

Third step: The general solution is therefore

$$y=\text{C.F.}+\text{P.I.}=c_1e^{3z}+c_2e^{2z}+\frac{1}{2}e^z-\frac{1}{6}a$$

$$=c_1(x+a)^3+c_2(x+a)^2+\frac{1}{2}(x+a)-\frac{1}{6}a \qquad [\because u=e^z,\ x+a=u],$$

where c_1, c_2 are two arbitrary constants.

2.8 SIMULTANEOUS LINEAR DIFFERENTIAL EQUATIONS

Here we shall consider linear differential equations with more than one dependent variable depending on only one independent variable. Such equations are known as simultaneous linear differential equations. Here we shall discuss simultaneous linear equations with constant coefficients only. To solve such equations completely, the number of equations must be equal to the number of dependent variables. There are two methods for solution of such a system.

Method-I

Let us consider the following simultaneous equations

$$f_1(D)x+f_2(D)y=f(t) \qquad ...(2.36)$$

$$g_1(D)x + g_2(D)y = g(t) \qquad ...(2.37)$$

where $D \equiv \frac{d}{dt}$, x, y are functions of t, f_1, f_2, g_1, g_2 are rational functions of D with constant coeffi-cients.

To eliminate y, we operate on both sides of (2.36) by $g_2(D)$ and on both sides of (2.37) by $f_2(D)$ and then subtracting, we get

$$\{g_2(D)f_1(D) - f_2(D)g_1(D)\}x = g_2(D)f(t) - f_2(D)g(t)$$

which is solved by the methods discussed earlier. Putting this value of x in either (2.36) or (2.37), we get the value of y.

Method-II

In this method we eliminate one dependent variable by differentiation and solve for the other variable (or variables) and then the value (or values) of the remaining variable (or variables) can be found. See example 3.

ILLUSTRATIVE EXAMPLES

Example 1: Solve $\frac{dx}{dt} - 7x + y = 0$ and $\frac{dy}{dt} - 2x - 5y = 0$. *(W.B.U.T. 2007)*

Solution: Let us write the given equation as

$$(D-7)x + y = 0 \qquad ...(1)$$

$$-2x + (D-5)y = 0 \qquad ...(2)$$

where $D \equiv \frac{d}{dt}$.

Operating both sides of (1) by $(D-5)$ and then subtracting from (2), we get

$$-\{(D-5)(D-7)+2\}x = 0, \quad \text{or} \quad (D^2 - 12D + 37)x = 0 \qquad ...(3)$$

Let $x = e^{mt}$ be a trial solution of (3), then the auxiliary equation becomes $m^2 - 12m + 37 = 0$, the roots are $m = 6 \pm i$.

$$\therefore \qquad x = e^{6t}(c_1 \cos t + c_2 \sin t)$$

$$\therefore \qquad \frac{dx}{dt} = 6e^{6t}(c_1 \cos t + c_2 \sin t) + e^{6t}(-c_1 \sin t + c_2 \cos t)$$

Therefore, from (1), we get

$$y = 7x - \frac{dx}{dt} = 7e^{6t}(c_1 \cos t + c_2 \sin t) - 6e^{6t}(c_1 \cos t + c_2 \sin t) - e^{6t}(-c_1 \sin t + c_2 \cos t)$$

$$= e^{6t}\{(c_1 - c_2)\cos t + (c_1 + c_2)\sin t\}$$

Therefore, the required solution is

$$x = e^{6t}(c_1 \cos t + c_2 \sin t)$$

$$y = e^{6t}\{(c_1 - c_2)\cos t + (c_1 + c_2)\sin t\},$$

where c_1, c_2 are arbitrary constants.

Example 2: Solve $\frac{dx}{dt} + 3x + y = e^t$, $\frac{dy}{dt} - x + y = e^{2t}$. *(W.B.U.T. 2004, 2012)*

Solution: Let us write the given equations as

$$(D+3)x + y = e^t \qquad ...(1)$$

$$-x + (D+1)y = e^{2t} \qquad ...(2)$$

where $D \equiv \frac{d}{dt}$.

Operating both sides of (1) by $(D+1)$ and then subtracting from (2), we get

$$-\{(D+1)(D+3)+1\}x = e^{2t} - (D+1)e^t, \text{ or } (D+2)^2 x = 2e^t - e^{2t} \qquad ...(3)$$

The complementary function of (3) is found from

$$(D+2)^2 x = 0.$$

Let $x = e^{mt}$ be a trial solution, then the auxiliary equation becomes $(m+2)^2 = 0$, the roots are $m = -2, -2$.

$$\therefore \qquad \text{C.F.} = (c_1 + c_2 t)e^{-2t}$$

$$\text{P.I.} = \frac{1}{(D+2)^2}(2e^t - e^{2t}) = \frac{1}{(D+2)^2}(2e^t) - \frac{1}{(D+2)^2}e^{2t}$$

$$= 2 \cdot \frac{1}{(1+2)^2}e^t - \frac{1}{(2+2)^2}e^{2t} = \frac{2}{9}e^t - \frac{1}{16}e^{2t}.$$

$$\therefore \qquad x = (c_1 + c_2 t)e^{-2t} + \frac{2}{9}e^t - \frac{1}{16}e^{2t}.$$

From (1), we have

$$y = e^t - (D+3)x = e^t - (D+3)\left\{(c_1 + c_2 t)e^{-2t} + \frac{2}{9}e^t - \frac{1}{16}e^{2t}\right\}$$

$$= e^t - \left\{c_2 e^{-2t} - 2(c_1 + c_2 t)e^{-2t} + \frac{2}{9}e^t - \frac{1}{8}e^{2t} + 3(c_1 + c_2 t)e^{-2t} + \frac{6}{9}e^t - \frac{3}{16}e^{2t}\right\}$$

$$= -c_2 e^{-2t} - (c_1 + c_2 t)e^{-2t} + \frac{1}{9}e^t + \frac{5}{16}e^{2t}.$$

Therefore, the required solution is

$$x = (c_1 + c_2 t)e^{-2t} + \frac{2}{9}e^t - \frac{1}{16}e^{2t}$$

$$y = -c_2 e^{-2t} - (c_1 + c_2 t)e^{-2t} + \frac{1}{9}e^t + \frac{5}{16}e^{2t},$$

where c_1, c_2 are arbitrary constants.

Indian Institute of Engineering Science and Technology, Shibpur
Ramanujan Central Library

Example 3: Solve $\frac{dx}{dt}+y=e^t$, $\frac{dy}{dt}-x=e^{-t}$. (W.B.U.T. 2003)

Solution: Here
$$\frac{dx}{dt}+y=e^t \quad ...(1)$$
$$\frac{dy}{dt}-x=e^{-t} \quad ...(2)$$

From (1), we have $y=e^t-\frac{dx}{dt}$, substituting this value of y in (2), we get

$$\frac{d}{dt}\left(e^t-\frac{dx}{dt}\right)-x=e^{-t}, \text{ or } e^t-\frac{d^2x}{dt^2}-x=e^{-t}$$

or
$$\frac{d^2x}{dt^2}+x=e^t-e^{-t}, \text{ or } (D^2+1)x=e^t-e^{-t}, \text{ where } D\equiv\frac{d}{dt}.$$

The complementary function is found from
$$(D^2+1)x=0.$$

Let $x=e^{mt}$ be a trial solution, then auxiliary equation becomes $m^2+1=0$, the roots are $m=\pm i$.

$$\therefore \quad \text{C.F.}=c_1\cos t+c_2\sin t.$$

$$\text{P.I.}=\frac{1}{D^2+1}(e^t-e^{-t})=\frac{1}{D^2+1}e^t-\frac{1}{D^2+1}e^{-t}$$
$$=\frac{1}{1^2+1}e^t-\frac{1}{(-1)^2+1}e^{-t}=\frac{1}{2}(e^t-e^{-t}).$$

$$\therefore \quad x=c_1\cos t+c_2\sin t+\frac{1}{2}(e^t-e^{-t}).$$

$$\therefore \quad \frac{dx}{dt}=-c_1\sin t+c_2\cos t+\frac{1}{2}(e^t+e^{-t}).$$

Now from (1),
$$y=e^t-\frac{dx}{dt}=e^t+c_1\sin t-c_2\cos t-\frac{1}{2}(e^t+e^{-t})$$
$$=c_1\sin t-c_2\cos t+\frac{1}{2}(e^t-e^{-t}).$$

Therefore, the required solution is
$$x=c_1\cos t+c_2\sin t+\frac{1}{2}(e^t-e^{-t})$$
$$y=c_1\sin t-c_2\cos t+\frac{1}{2}(e^t-e^{-t}),$$
where c_1, c_2 are arbitrary constants.

Example 4: Solve $\frac{dx}{dt}+y=1-\cos t$, $\frac{dy}{dt}+x=\sin t$.

Solution: Writing D for $\frac{d}{dt}$, the given equations become
$$Dx+y=1-\cos t \quad ...(1)$$
$$Dy+x=\sin t \quad ...(2)$$

Operating both sides of (1) with D, we get
$$D^2x+Dy=\sin t \quad ...(3)$$

Subtracting (2) from (3), we get
$$D^2x-x=0, \quad \text{or} \quad (D^2-1)x=0.$$

Let $x=e^{mt}$ be a trial solution, then the auxiliary equation becomes $m^2-1=0$, the roots are $m=\pm 1$.

$$\therefore \quad x=c_1e^t+c_2e^{-t}$$
$$\therefore \quad \frac{dx}{dt}=c_1e^t-c_2e^{-t}$$

Using this in (1), we get
$$y=1-\cos t-\frac{dx}{dt}=1-\cos t-c_1e^t+c_2e^{-t}$$

Therefore, the required solution is
$$x=c_1e^t+c_2e^{-t},\ y=1-\cos t-c_1e^t+c_2e^{-t},$$
where c_1, c_2 are arbitrary constants.

Example 5: Solve the following simultaneous equations:
$$\frac{dx}{dt}+5x-2y=t,\ \frac{dy}{dt}+2x+y=0,$$
it is given that $x=y=0$ when $t=0$.

Solution: Writing D for $\frac{d}{dt}$, the given equations become
$$(D+5)x-2y=t \quad ...(1)$$
$$2x+(D+1)y=0 \quad ...(2)$$

Operating both sides of (1) by $(D+1)$, multiplying both sides of (2) by 2 and then adding, we get
$$\{(D+1)(D+5)+4\}x=(D+1)t, \quad \text{or} \quad (D+3)^2x=1+t \quad ...(3)$$

The complementary function of (3) is found from
$$(D+3)^2x=0.$$

Let $x=e^{mt}$ be a trial solution, then the auxiliary equation becomes $(m+3)^2=0$, the roots are $m=-3,-3$.

$$\therefore \quad \text{C.F.}=(c_1+c_2t)e^{-3t}.$$

$$\text{P.I.} = \frac{1}{(D+3)^2}(1+t) = \frac{1}{9}\left(1+\frac{D}{3}\right)^{-2}(1+t) = \frac{1}{9}\left(1-2\frac{D}{3}+3\cdot\frac{D^2}{9}-\ldots\right)(1+t)$$

$$= \frac{1}{9}\left(1+t-\frac{2}{3}\right) = \frac{t}{9}+\frac{1}{27}.$$

$$\therefore \quad x = (c_1+c_2t)e^{-3t}+\frac{t}{9}+\frac{1}{27}. \qquad \ldots(4)$$

Using (1), we get

$$y = \frac{1}{2}(D+5)x-\frac{t}{2} = \frac{1}{2}(D+5)\left\{(c_1+c_2t)e^{-3t}+\frac{t}{9}+\frac{1}{27}\right\}-\frac{t}{2}$$

$$= \frac{1}{2}\left\{c_2e^{-3t}-3(c_1+c_2t)e^{-3t}+\frac{1}{9}+5(c_1+c_2t)e^{-3t}+\frac{5t}{9}+\frac{5}{27}\right\}-\frac{t}{2}$$

$$= \left(c_1+\frac{c_2}{2}+c_2t\right)e^{-3t}-\frac{2t}{9}+\frac{4}{27}. \qquad \ldots(5)$$

By question $x = y = 0$ when $t = 0$, therefore from (4) and (5), we get

$$c_1+\frac{1}{27} = 0,\ c_1+\frac{c_2}{2}+\frac{4}{27} = 0, \quad \therefore \quad c_1 = -\frac{1}{27},\ c_2 = -\frac{6}{27}.$$

Hence the required solution is

$$x = -\frac{1}{27}(1+6t)e^{-3t}+\frac{1}{27}(1+3t),$$

$$y = -\frac{2}{27}(2+3t)e^{-3t}+\frac{2}{27}(2-3t).$$

Example 6: Solve: $\dfrac{dx}{dt}+2\dfrac{dy}{dt}-2x+2y = 3e^t$

and $\quad 3\dfrac{dx}{dt}+\dfrac{dy}{dt}+2x+y = 4e^{2t}.$

Solution: Writing D for $\dfrac{d}{dt}$, the given equations become

$$(D-2)x+2(D+1)y = 3e^t \qquad \ldots(1)$$

$$(3D+2)x+(D+1)y = 4e^{2t} \qquad \ldots(2)$$

Multiplying both sides of (2) by 2 and then subtracting from (1), we get

$$(D-2-6D-4)x = 3e^t-8e^{2t}, \text{ or } (5D+6)x = 8e^{2t}-3e^t$$

$$\therefore \quad \frac{dx}{dt}+\frac{6}{5}x = \frac{8}{5}e^{2t}-\frac{3}{5}e^t, \text{ which is linear in } x, \qquad \ldots(3)$$

$$\therefore \quad \text{I.F.} = e^{\int\frac{6}{5}dt} = e^{\frac{6}{5}t}.$$

Multiplying both sides of (3) by $e^{\frac{6}{5}t}$, we get

$$e^{\frac{6}{5}t}\frac{dx}{dt}+\frac{6}{5}xe^{\frac{6}{5}t} = \left(\frac{8}{5}e^{2t}-\frac{3}{5}e^t\right)e^{\frac{6}{5}t}$$

or $$\frac{d}{dt}(xe^{\frac{6}{5}t}) = \frac{8}{5}e^{\frac{16}{5}t}-\frac{3}{5}e^{\frac{11}{5}t}$$

or $$d(xe^{\frac{6}{5}t}) = \left(\frac{8}{5}e^{\frac{16}{5}t}-\frac{3}{5}e^{\frac{11}{5}t}\right)dt.$$

Integrating, $$\int d(xe^{\frac{6}{5}t}) = \int\left(\frac{8}{5}e^{\frac{16}{5}t}-\frac{3}{5}e^{\frac{11}{5}t}\right)dt$$

$$\therefore \quad xe^{\frac{6}{5}t} = \frac{1}{2}e^{\frac{16}{5}t}-\frac{3}{11}e^{\frac{11}{5}t}+c_1$$

$$\therefore \quad x = \frac{1}{2}e^{2t}-\frac{3}{11}e^t+c_1e^{-\frac{6}{5}t}$$

$$\therefore \quad \frac{dx}{dt} = e^{2t}-\frac{3}{11}e^t-\frac{6}{5}c_1e^{-\frac{6}{5}t}$$

Therefore, from (1) we get

$$e^{2t}-\frac{3}{11}e^t-\frac{6}{5}c_1e^{-\frac{6}{5}t}-e^{2t}+\frac{6}{11}e^t-2c_1e^{-\frac{6}{5}t}+2\frac{dy}{dt}+2y = 3e^t$$

or $$2\frac{dy}{dt}+2y = \frac{30}{11}e^t+\frac{16}{5}c_1e^{-\frac{6}{5}t}$$

$$\therefore \quad \frac{dy}{dt}+y = \frac{15}{11}e^t+\frac{8}{5}c_1e^{-\frac{6}{5}t}, \text{ which is linear in } y. \qquad \ldots(4)$$

$$\therefore \quad \text{I.F.} = e^{\int dt} = e^t.$$

Multiplying both sides of (4) by e^t, we get

$$e^t\frac{dy}{dt}+ye^t = \frac{15}{11}e^{2t}+\frac{8}{5}c_1e^{-\frac{t}{5}}$$

or $$\frac{d}{dt}(ye^t) = \frac{15}{11}e^{2t}+\frac{8}{5}c_1e^{-\frac{t}{5}}$$

$$\therefore \quad d(ye^t) = \left(\frac{15}{11}e^{2t}+\frac{8}{5}c_1e^{-\frac{t}{5}}\right)dt$$

Integrating, $$\int d(ye^t) = \int\left(\frac{15}{11}e^{2t}+\frac{8}{5}c_1e^{-\frac{t}{5}}\right)dt$$

$$\therefore \quad ye^t = \frac{15}{22}e^{2t}-8c_1e^{-\frac{t}{5}}+c_2$$

or $$y = \frac{15}{22}e^{t} - 8c_1e^{-\frac{6t}{5}} + c_2e^{-t}$$

Hence the required solution is

$$x = \frac{1}{2}e^{2t} - \frac{3}{11}e^{t} + c_1e^{-\frac{6}{5}t}$$

$$y = \frac{15}{22}e^{t} - 8c_1e^{-\frac{6}{5}t} + c_2e^{-t},$$

where c_1, c_2 are arbitrary constants.

Example 7: Solve $\frac{dx}{dt} + \frac{dy}{dt} - 2y = \cos t + \sin t$

and $$\frac{dx}{dt} - \frac{dy}{dt} + 2x = \cos t - \sin t.$$

Solution: Writing D for $\frac{d}{dt}$, the given equations become

$$Dx + (D-2)y = \cos t + \sin t \quad \text{...(1)}$$

$$(D+2)x + Dy = \cos t - \sin t \quad \text{...(2)}$$

Operating both sides of (1) by D and both sides of (2) by $(D-2)$ and then adding, we get

$$[D^2 + (D-2)(D+2)]x = D(\cos t + \sin t) + (D-2)(\cos t - \sin t)$$

or $$(2D^2 - 4)x = -2\cos t, \quad \text{or} \quad (D^2 - 2)x = -\cos t.$$

The complementary function is found from

$$(D^2 - 2)x = 0.$$

Let $x = e^{mt}$ be a trial solution, then the auxiliary equation becomes $m^2 - 2 = 0$, the roots are $m = \pm\sqrt{2}$.

$\therefore$ $$\text{C.F.} = c_1e^{\sqrt{2}t} + c_2e^{-\sqrt{2}t}.$$

$$\text{P.I.} = \frac{1}{D^2-2}(-\cos t) = -\frac{1}{-1^2-2}\cos t = \frac{1}{3}\cos t.$$

$\therefore$ $$x = c_1e^{\sqrt{2}t} + c_2e^{-\sqrt{2}t} + \frac{1}{3}\cos t$$

$\therefore$ $$\frac{dx}{dt} = c_1\sqrt{2}e^{\sqrt{2}t} - c_2\sqrt{2}e^{-\sqrt{2}t} - \frac{1}{3}\sin t \quad \text{...(3)}$$

Adding (1) and (2), we get

$$2\frac{dx}{dt} + 2x - 2y = 2\cos t$$

$\therefore$ $$y = \frac{dx}{dt} + x - \cos t = c_1\sqrt{2}e^{\sqrt{2}t} - c_2\sqrt{2}e^{-\sqrt{2}t} - \frac{1}{3}\sin t$$
$$+ c_1e^{\sqrt{2}t} + c_2e^{-\sqrt{2}t} + \frac{1}{3}\cos t - \cos t$$

Hence the required solution is

$$x = c_1e^{\sqrt{2}t} + c_2e^{-\sqrt{2}t} + \frac{1}{3}\cos t$$

$$y = (1+\sqrt{2})c_1e^{\sqrt{2}t} + (1-\sqrt{2})c_2e^{-\sqrt{2}t} - \frac{2}{3}\cos t - \frac{1}{3}\sin t,$$

where c_1, c_2 are two arbitrary constants.

Example 8: Solve $\frac{d^2x}{dt^2} - \frac{dy}{dt} = x + 2t$ and $6\frac{dx}{dt} + 4\frac{dy}{dt} = -2t$.

Solution: Writing D for $\frac{d}{dt}$, the given equations become

$$(D^2 - 1)x - Dy = 2t \quad \text{...(1)}$$

$$6Dx + 4Dy = -2t \quad \text{...(2)}$$

Multiplying both sides of (1) by 4 and adding with (2), we get

$$(4D^2 + 6D - 4)x = 6t$$

The complementary function is found from

$$(4D^2 + 6D - 4)x = 0.$$

Let $x = e^{mt}$ be a trial solution, then the auxiliary equation becomes $4m^2 + 6m - 4 = 0$, or $4m^2 + 8m - 2m - 4 = 0$, or $2(2m-1)(m+2) = 0$, the roots are $m = \frac{1}{2}, -2$.

$\therefore$ $$\text{C.F.} = c_1e^{\frac{t}{2}} + c_2e^{-2t}$$

$$\text{P.I.} = \frac{1}{4D^2 + 6D - 4}(6t) = \frac{6}{-4}\left\{1 - \left(\frac{3}{2}D + D^2\right)\right\}^{-1}t$$

$$= -\frac{3}{2}\left(1 + \frac{3}{2}D + D^2 + \ldots\right)t = -\frac{3}{2}\left(t + \frac{3}{2}\right) = -\frac{3}{4}(2t+3)$$

$\therefore$ $$x = c_1e^{\frac{t}{2}} + c_2e^{-2t} - \frac{3}{4}(2t+3)$$

$\therefore$ $$\frac{dx}{dt} = \frac{c_1}{2}e^{\frac{t}{2}} - 2c_2e^{-2t} - \frac{3}{2}$$

Putting this value of $\frac{dx}{dt}$ in (2), we get

$$3c_1e^{\frac{t}{2}} - 12c_2e^{-2t} - 9 + 4\frac{dy}{dt} = -2t$$

or $$\frac{dy}{dt} = \frac{1}{4}(12c_2e^{-2t} - 3c_1e^{\frac{t}{2}} - 2t + 9)$$

$$\therefore \quad \int dy = \frac{1}{4}\int (12c_2e^{-2t} - 3c_1e^{\frac{t}{2}} - 2t + 9)\,dt$$

$$\therefore \quad y = -\frac{3}{2}c_2e^{-2t} - \frac{3}{2}c_1e^{\frac{t}{2}} - \frac{1}{4}t^2 + \frac{9}{4}t + c_3.$$

Hence the required solution is

$$x = c_1e^{\frac{t}{2}} + c_2e^{-2t} - \frac{3}{4}(2t+3)$$

$$y = -\frac{3}{2}c_1e^{\frac{t}{2}} - \frac{3}{2}c_2e^{-2t} - \frac{1}{4}t^2 + \frac{9}{4}t + c_3,$$

where c_1, c_2, c_3 are arbitrary constants.

Example 9: Solve the following simultaneous differential equations:

$$x\frac{dy}{dx} + z = 0, \quad x\frac{dz}{dx} + y = 0.$$

Solution: Here $\quad x\dfrac{dy}{dx} + z = 0 \qquad$ ...(1)

$$x\frac{dz}{dx} + y = 0 \qquad \text{...(2)}$$

Differentiating both sides of (1) w.r.t. x, we get

$$x\frac{d^2y}{dx^2} + \frac{dy}{dx} + \frac{dz}{dx} = 0, \quad \text{or} \quad x^2\frac{d^2y}{dx^2} + x\frac{dy}{dx} + x\frac{dz}{dx} = 0$$

$$\therefore \quad x^2\frac{d^2y}{dx^2} + x\frac{dy}{dx} - y = 0 \qquad \left[\text{Since from (2), } x\frac{dz}{dx} = -y\right].$$

or $$(x^2D^2 + xD - 1)y = 0, \quad \text{where } D \equiv \frac{d}{dx}. \qquad \text{...(3)}$$

Putting $x = e^t$, *i.e.*, $t = \log x$, we have

$$xDy = D'y, \; x^2D^2y = D'(D'-1)y, \quad \text{where } D' \equiv \frac{d}{dt}.$$

Therefore equation (3) reduces to

$$\{D'(D'-1) + D' - 1\}y = 0, \quad \text{or} \quad (D'^2 - 1)y = 0.$$

Let $y = e^{mt}$ be a trial solution, then the auxiliary equation becomes $m^2 - 1 = 0$, the roots are $m = \pm 1$.

$$\therefore \quad y = c_1e^t + c_2e^{-t}, \text{ i.e., } y = c_1x + \frac{c_2}{x} \qquad (\because x = e^t)$$

$$\therefore \quad \frac{dy}{dx} = c_1 - \frac{c_2}{x^2}$$

From (1), we have $z = -x\dfrac{dy}{dx} = -x\left(c_1 - \dfrac{c_2}{x^2}\right) = -c_1x + \dfrac{c_2}{x}$.

Hence the required solution is

$$y = c_1x + \frac{c_2}{x}, \quad z = -c_1x + \frac{c_2}{x},$$

where c_1, c_2 are arbitrary constants.

Example 10: Solve $\dfrac{dx}{dt} = y + z, \; \dfrac{dy}{dt} = z + x, \; \dfrac{dz}{dt} = x + y.$

Solution: Here, $\quad \dfrac{dx}{dt} = y + z \qquad$ ...(1)

$$\frac{dy}{dt} = z + x \qquad \text{...(2)}$$

$$\frac{dz}{dt} = x + y \qquad \text{...(3)}$$

From (1), we have $\quad \dfrac{d^2x}{dt^2} = \dfrac{dy}{dt} + \dfrac{dz}{dt} = (z+x) + (x+y) \qquad$ [by (2) and (3)]

$$= 2x + \frac{dx}{dt} \qquad \text{[by (1)]}$$

$$\therefore \quad \frac{d^2x}{dt^2} - \frac{dx}{dt} - 2x = 0. \qquad \text{...(4)}$$

Let $x = e^{mt}$ be a trial solution of (4), then the auxiliary equation becomes $m^2 - m - 2 = 0$, or $m^2 - 2m + m - 2 = 0$, or $(m-2)(m+1) = 0$, the roots are $m = 2, -1$.

$$\therefore \quad x = c_1e^{2t} + c_2e^{-t}$$

$$\therefore \quad \frac{dx}{dt} = 2c_1e^{2t} - c_2e^{-t}, \quad \text{or} \quad y + z = 2c_1e^{2t} - c_2e^{-t} \quad \text{[by (1)]} \qquad \text{...(5)}$$

From (2) and (3), we have $\dfrac{dy}{dt} - \dfrac{dz}{dt} = (z+x) - (x+y)$,

or $$\frac{d}{dt}(y-z) = -(y-z), \quad \text{or} \quad \frac{d(y-z)}{y-z} = -dt$$

Integrating, $$\int \frac{d(y-z)}{y-z} = -\int dt, \quad \text{or} \quad \log(y-z) = -t + \log c_3$$

or $$\log\left(\frac{y-z}{c_3}\right) = -t \quad \therefore y - z = c_3e^{-t} \qquad \text{...(6)}$$

Solving (5) and (6), we get

$$y = c_1e^{2t} - \frac{1}{2}(c_2 - c_3)e^{-t}, \quad z = c_1e^{2t} - \frac{1}{2}(c_2 + c_3)e^{-t}.$$

Therefore, the required solution is

$$x = c_1e^{2t} + c_2e^{-t}, \quad y = c_1e^{2t} - \frac{1}{2}(c_2 - c_3)e^{-t}$$

and $$z = c_1 e^{2t} - \frac{1}{2}(c_2 + c_3)e^{-t},$$

where c_1, c_2, c_3 are arbitrary constants.

MULTIPLE CHOICE QUESTIONS

1. The general solution of the ordinary differential equation $\frac{d^2y}{dx^2} + 4y = 0$ is

   (a) $Ae^{2x} + Be^{-2x}$ (b) $(A + Bx)e^{2x}$
   (c) $A\cos 2x + B\sin 2x$ (d) $(A + Bx)\cos 2x$.
   (where A and B are arbitrary constants). (W.B.U.T. 2006, Arch-201/13)

2. The general solution of $\frac{d^2y}{dx^2} = 0$ is

   (a) $y = a$ (b) $y = ax + b$
   (c) $y = ax^2 + bx + c$ (d) $y = a\cos x + b\sin x$.
   (where a, b, c are arbitrary constants.)

3. $y = c_1e^x + c_2e^{-x}$ (c_1, c_2 are constants) is the solution of the differential equation

   (a) $\frac{d^2y}{dx^2} - y = 0$ (b) $\frac{d^2y}{dx^2} + y = 0$
   (c) $\frac{d^2y}{dx^2} - 2y = 0$ (d) none of these.

4. $y = c_1\cos 2x + c_2\sin 2x$ (c_1, c_2 are constants) is the solution of

   (a) $\frac{d^2y}{dx^2} - 2y = 0$ (b) $\frac{d^2y}{dx^2} + 2y = 0$
   (c) $\frac{d^2y}{dx^2} - 4y = 0$ (d) $\frac{d^2y}{dx^2} + 4y = 0$.

5. The general solution of $(D-1)^2 y = 0$, where $D \equiv \frac{d}{dx}$ is

   (a) $y = c_1 + c_2x$ (b) $y = c_1e^x$
   (c) $y = (c_1 + c_2x)e^x$ (d) none of these.
   (where c_1, c_2 are arbitrary constants).

6. The general solution of $\frac{d^2y}{dx^2} - 5\frac{dy}{dx} + 6y = 0$ is

   (a) $y = c_1e^{-2x} + c_2e^{-3x}$ (b) $y = c_1e^{2x} + c_2e^{3x}$
   (c) $y = e^{2x} + e^{3x}$ (d) none of these.
   (where c_1, c_2 are arbitrary constants).

7. The general solution of the differential equation $D^2y + 9y = 0$ is

   (a) $Ae^{3x} + Be^{-3x}$ (b) $(A + Bx)e^{3x}$
   (c) $A\cos 3x + B\sin 3x$ (d) $(A + Bx)\sin 3x$. (W.B.U.T. 2008)

8. $\frac{1}{D-1}x^2$ is equal to

   (a) $x^2 + 2x + 2$ (b) $-(x^2 + 2x + 2)$
   (c) $2x - x^2$ (d) $-(2x - x^2)$. (W.B.U.T. 2008, 2010, 2013)

9. $\frac{1}{D-3}e^{3x}$ is equal to

   (a) xe^{3x} (b) $3e^{3x}$ (c) $\frac{1}{2}x^2e^{3x}$ (d) none of these.

10. $\frac{1}{(D-1)^2}e^x$ is equal to

    (a) e^x (b) xe^x (c) x^2e^x (d) $\frac{x^2}{2}e^x$.

11. $\frac{1}{D^2+1}\sin 2x =$

    (a) $\frac{1}{3}\sin 2x$ (b) $\frac{1}{3}\cos 2x$
    (c) $-\frac{1}{3}\sin 2x$ (d) none of these.

12. The particular integral of $\frac{d^2y}{dx^2} + y = x^2$ is

    (a) $x^2 - 2x + 2$ (b) $x^2 - 2$
    (c) $x^2 + 2$ (d) none of these.

13. $\frac{1}{D^2 + D + 1}\cos x =$

    (a) $\sin x$ (b) $-\sin x$ (c) $\cos x$ (d) none of these.

14. $\frac{1}{(D-2)(D-3)}e^{3x} =$

    (a) e^{3x} (b) $3e^{3x}$ (c) $\frac{1}{2}x^2e^{3x}$ (d) xe^{3x}.

15. $\frac{1}{D-a}X =$

    (a) $\int Xe^{-ax}dx$ (b) $e^{-ax}\int Xe^{ax}dx$ (c) $e^{ax}\int Xe^{-ax}dx$ (d) none of these.

16. The differential equation $x^2\frac{d^2y}{dx^2}-x\frac{dy}{dx}-3y=x$ is solved by the transformation

(a) $e^x=z$ (b) $\log x=z$ (c) $x=z$ (d) none of these.

17. Using the substitution $x=e^z$, the equation $x\frac{d^2y}{dx^2}+\frac{dy}{dx}=\frac{1}{x}$ reduces to

(a) $\frac{d^2y}{dz^2}=1$ (b) $\frac{d^2y}{dz^2}+\frac{dy}{dz}=1$

(c) $\frac{d^2y}{dz^2}-\frac{dy}{dz}=1$ (d) none of these.

18. The equation $x^2\frac{d^2y}{dx^2}+x\frac{dy}{dx}+y=\sin(\log x^2)$ is a Cauchy-Euler equation. The statement is

(a) True (b) False.

19. The differential equation involving x and t, obtained from the simultaneous equations $\frac{dx}{dt}=y$, $\frac{dy}{dt}=z$, $\frac{dz}{dt}=x$ is

(a) $\frac{d^3x}{dt^3}+x=0$ (b) $\frac{d^2x}{dt^2}-x=0$

(c) $\frac{d^3x}{dt^3}-x=0$ (d) none of these.

20. The solution of the system $\frac{dx}{dt}=y$, $\frac{dy}{dt}=-x$ is

(a) $x=c_1e^t+c_2e^{-t}$, $y=c_1e^t-c_2e^{-t}$

(b) $x=c_1\cos t+c_2\sin t$, $y=-c_1\sin t+c_2\cos t$

(c) $x=c_1\cos t+c_2\sin t$, $y=c_1\sin t-c_2\cos t$

(d) none of these.

21. The substitution $x=e^z$ transforms the differential equation $x^2\frac{d^2y}{dx^2}-5y=\log_e x$ to

(a) $\frac{d^2y}{dz^2}+\frac{dy}{dz}-5y=z$ (b) $\frac{d^2y}{dz^2}-\frac{dy}{dz}+5y=z$

(c) $\frac{d^2y}{dz^2}-\frac{dy}{dz}+3y=0$ (d) $\frac{d^2y}{dz^2}-\frac{dy}{dz}-5y=z$ (M–201/12)

22. The particular integral of $(D^2-4D+4)y=x^3e^{2x}$ is

(a) $\frac{e^{2x}x^4}{20}$ (b) $\frac{e^{2x}x^5}{20}$

(c) $\frac{e^{2x}x^4}{60}$ (d) $\frac{e^{x}x^5}{20}$ (M–201/12)

23. The particular integral of $\frac{d^2y}{dx^2}+y=\cos x$ is

(a) $\frac{1}{2}\sin x$ (b) $\frac{1}{2}\cos x$

(c) $\frac{1}{2}x\sin x$ (d) $\frac{1}{2}x\cos x$ [see page 98] (M–201/13)

24. The value of $\frac{1}{D}(x^2)$ is

(a) x (b) $2x$

(c) $\frac{x^3}{3}$ (d) x^2 (Arch–201/13)

25. The complementary function of the differential equation $\frac{d^2y}{dx^2}+3\frac{dy}{dx}+2y=0$ is

(a) $Ae^{-2x}+Be^{-x}$ (b) $Ae^{2x}+Be^{x}$

(c) $Ae^{2x}+Be^{-x}$ (d) $Ae^{-2x}+Be^{x}$ (Arch–201/13)

ANSWERS

1. (c)	2. (b)	3. (a)	4. (d)	5. (c)
6. (b)	7. (c)	8. (b)	9. (a)	10. (d)
11. (c)	12. (b)	13. (a)	14. (d)	15. (c)
16. (b)	17. (a)	18. (a)	19. (c)	20. (b)
21. (d)	22. (b)	23. (c)	24. (c)	25. (a).

PROBLEMS

Solve the following differential equations:

1. $\frac{d^2y}{dx^2}-4\frac{dy}{dx}+y=x^2-2x+2$.

2. $\frac{d^2y}{dx^2}-6\frac{dy}{dx}+9y=6e^{3x}+7e^{-2x}-\log 2$.

3. $\frac{d^2y}{dx^2}+3\frac{dy}{dx}+2y=4\cos^2 x$.

4. $\frac{d^2y}{dx^2}+2\frac{dy}{dx}+y=e^{2x}-\cos^2 x$.

5. $\frac{d^2y}{dx^2}-4y=\cosh(2x-1)+3^x$.

6. $\frac{d^2y}{dx^2}+4y=x^2+\cos 2x$.

7. $\frac{d^2y}{dx^2}-6\frac{dy}{dx}+25y=e^{2x}+\sin x+x$.

8. $(D^2+4)y=\sin 2x$, where $D\equiv\frac{d}{dx}$.

9. $\frac{d^2y}{dx^2}-3\frac{dy}{dx}+2y=\sin 2x$.

10. $\frac{d^2y}{dx^2}+6\frac{dy}{dx}+9y=24e^{-3x}$.

11. $(D^2-7D+6)y = (x-2)e^x$.

12. $\frac{d^2y}{dx^2}-2\frac{dy}{dx}+y = xe^x$. (W.B.U.T. 2013)

13. $\frac{d^2y}{dx^2}-3\frac{dy}{dx} = x+e^x\sin x$.

14. $(D^2-4D+4)y = x^3e^{2x}+xe^{2x}$.

15. $\frac{d^2y}{dx^2}+\frac{dy}{dx}-6y = 10e^{2x}-18e^{3x}-6x-11$.

16. $\frac{d^2y}{dx^2}+a^2y = \operatorname{cosec} ax \quad (a \neq 0)$.

17. $\frac{d^2y}{dx^2}+2y = x^2e^{3x}+e^x\cos 2x$.

18. $\frac{d^2y}{dx^2}-2\frac{dy}{dx}+4y = e^x\cos x$.

19. $\frac{d^2y}{dx^2}-y = x^2\cos x$.

20. $\frac{d^2x}{dt^2}+4\frac{dx}{dt}+4x = 3\sin 2t$, given that $x=0$, $\frac{dx}{dt}=0$ at $t=0$.

21. $\frac{d^2y}{dx^2}-7\frac{dy}{dx}+6y = 2e^{3x}$, given that $y=1$, $\frac{dy}{dx}=0$, when $x=0$.

Solve by the method of variation of parameters:

22. $\frac{d^2y}{dx^2}-6\frac{dy}{dx}+9y = \frac{e^{3x}}{x^2}$.

23. $\frac{d^2y}{dx^2}-y = \frac{2}{1+e^x}$.

24. $\frac{d^2y}{dx^2}+4y = 4\tan 2x$. (W.B.U.T. 2002)

25. $\frac{d^2y}{dx^2}+9y = \sec 3x$ (W.B.U.T. 2005, 2008, 2009)

26. $\frac{d^2y}{dx^2}+y = x\sin x$.

27. $\frac{d^2y}{dx^2}-2\frac{dy}{dx}+y = \frac{e^x}{x}$.

28. $\frac{d^2y}{dx^2}-2\frac{dy}{dx}+2y = e^x\tan x$.

29. $\frac{d^2y}{dx^2}-2\frac{dy}{dx} = e^x\sin x$.

30. $\frac{d^2y}{dx^2}+y = \frac{1}{1+\sin x}$.

31. $\frac{d^2y}{dx^2}+a^2y = \tan ax. \quad (a \neq 0)$

32. $\frac{d^2y}{dx^2}+y = \cos x$.

33. $(2x+1)(x+1)\frac{d^2y}{dx^2}+2x\frac{dy}{dx}-2y = (2x+1)^2$, it is given that $y=x$ and $y=(x+1)^{-1}$ are two linearly independent solutions of the corresponding homogeneous equation.

34. $x^2\frac{d^2y}{dx^2}-x(x+2)\frac{dy}{dx}+(x+2)y = x^3$, it is given that $y=x$, $y=xe^x$ are two linearly independent solutions of the corresponding homogeneous equation.

Solve the following differential equations:

35. $x^2\frac{d^2y}{dx^2}-x\frac{dy}{dx}+y = 0$.

36. $x^2\frac{d^2y}{dx^2}-3x\frac{dy}{dx}+4y = 2x^2$. (W.B.U.T. 2013)

37. $x^2\frac{d^2y}{dx^2}+5x\frac{dy}{dx}+4y = x\log x$.

38. $x^2\frac{d^2y}{dx^2}-2x\frac{dy}{dx}-4y = x^2+2\log x$.

39. $x^2\frac{d^2y}{dx^2}-x\frac{dy}{dx}+y = \log x$.

40. $(1+x)^2\frac{d^2y}{dx^2}+(1+x)\frac{dy}{dx}+y = 4\cos\log(1+x)$.

41. $(2+x)^2\frac{d^2y}{dx^2}+(2+x)\frac{dy}{dx}+4y = 2\sin\{2\log(2+x)\}$.

Solve the following system of simultaneous equations:

42. $\frac{dx}{dt}+\frac{dy}{dt}+2x+y = 0$, $\frac{dy}{dt}+5x+3y = 0$.

43. $\frac{dx}{dt}+4x+3y = t$, $\frac{dy}{dt}+2x+5y = e^t$.

44. $-\frac{dx}{dt}+3\frac{dy}{dt}+y = e^t$, $3\frac{dx}{dt}-\frac{dy}{dt}+x = 0$.

45. $\frac{dx}{dt}+2y+\sin t = 0$, $\frac{dy}{dt}-2x-\cos t = 0$, given $x=0$, $y=1$ at $t=0$.

46. $\frac{dx}{dt}+\frac{dy}{dt}-2y = 2\cos t-7\sin t$, $\frac{dx}{dt}-\frac{dy}{dt}+2x = 4\cos t-3\sin t$.

47. $\frac{dx}{dt}-5x-y = 0$, $\frac{dy}{dt}-y+4x = 0$.

48. $\frac{dx}{dt}+y = \sin t$, $\frac{dy}{dt}+x = \cos t$, given that $x=2$ and $y=0$ at $t=0$.

49. $\frac{dx}{dt}+2x+3y = 0$, $\frac{dy}{dt}+3x+2y = 2e^{2t}$.

50. $\frac{dx}{dt}+2y = e^t$, $\frac{dy}{dt}-2x = e^{-t}$.

51. $\frac{dx}{dt}+2x-3y = 5t$, $\frac{dy}{dt}-3x+2y = 2e^{2t}$.

52. $\frac{dx}{dt}+\frac{dy}{dt}+3x = \sin t$, $\frac{dx}{dt}+y-x = \cos t$.

ANSWERS

1. $y = c_1e^{(2+\sqrt{3})x}+c_2e^{(2-\sqrt{3})x}+x^2+6x+24$.

2. $y = (c_1+c_2x)e^{3x}+3x^2e^{3x}+\frac{7}{25}e^{-2x}-\frac{1}{9}\log 2$.

3. $y = c_1e^{-x}+c_2e^{-2x}+1+\frac{1}{10}(3\sin 2x-\cos 2x)$.

4. $y = (c_1+c_2x)e^{-x}+\frac{1}{2}+\frac{1}{5}(2\sin 2x+\cos 2x)$.

5. $y = c_1e^{2x} + c_2e^{-2x} + \frac{x}{4}\sinh(2x-1) + \frac{3^x}{(\log 3)^2 - 4}$

6. $y = c_1\cos 2x + c_2\sin 2x + \frac{1}{4}x^2 - \frac{1}{8} - \frac{1}{4}x\cos 2x.$

7. $y = e^{3x}(c_1\cos 4x + c_2\sin 4x) + \frac{1}{17}e^{2x} + \frac{1}{565}(23\sin x + 6\cos x) + \frac{x}{25} + \frac{6}{625}.$

8. $y = c_1\cos 2x + c_2\sin 2x - \frac{1}{4}x\cos 2x.$

9. $y = c_1e^x + c_2e^{2x} - \frac{1}{20}\sin 2x + \frac{3}{20}\cos 2x.$

10. $y = (c_1 + c_2x)e^{-3x} + 12x^2e^{-3x}.$

11. $y = c_1e^x + c_2e^{6x} + \left(\frac{9}{25}x - \frac{1}{10}x^2\right)e^x.$

12. $y = e^x(c_1 + c_2x) + \frac{1}{6}x^3e^x.$

13. $y = c_1 + c_2e^{3x} - \frac{1}{3}\left(\frac{1}{2}x^2 + \frac{1}{3}x\right) + \frac{e^x}{10}(\cos x - 3\sin x).$

14. $y = (c_1 + c_2x)e^{2x} + \frac{1}{20}x^5e^{2x} + \frac{1}{6}x^3e^{2x}.$

15. $y = c_1e^{2x} + c_2e^{-3x} + 2xe^{2x} - 3e^{3x} + x + 2.$

16. $y = c_1\cos ax + c_2\sin ax - \frac{x}{a}\cos ax + \frac{1}{a^2}\sin ax\log\sin ax.$

17. $y = c_1\cos\sqrt{2}x + c_2\sin\sqrt{2}x + \frac{e^{3x}}{121}\left(11x^2 - 12x + \frac{50}{11}\right) + \frac{e^x}{17}(4\sin 2x - \cos 2x).$

18. $y = e^x(c_1\cos\sqrt{3}x + c_2\sin\sqrt{3}x) + \frac{1}{2}e^x\cos x.$

19. $y = c_1e^x + c_2e^{-x} + \frac{1}{2}(1 - x^2)\cos x + x\sin x.$

20. $x = \frac{1}{8}(3 + 6t)e^{-2t} - \frac{3}{8}\cos 2t.$

21. $y = \frac{7}{5}e^x - \frac{1}{15}e^{6x} - \frac{1}{3}e^{3x}.$

22. $y = (c_1 + c_2x)e^{3x} - e^{3x}(\log x + 1).$

23. $y = c_1e^x + c_2e^{-x} - 1 + e^x\log(e^{-x} + 1) - e^{-x}\log(e^x + 1).$

24. $y = c_1\cos 2x + c_2\sin 2x - \cos 2x\log(\sec 2x + \tan 2x).$

25. $y = c_1\cos 3x + c_2\sin 3x + \frac{x}{3}\sin 3x + \frac{1}{9}\cos 3x\log\cos 3x.$

26. $y = c_1\cos x + c_2\sin x + \frac{x}{2}\sin x - \frac{x^2}{4}\cos x.$

27. $y = (c_1 + c_2x)e^x + xe^x\log x.$

28. $y = e^x(c_1\cos x + c_2\sin x) - e^x\cos x\log(\sec x + \tan x).$

29. $y = c_1 + c_2e^{2x} - \frac{1}{2}e^x\sin x.$

30. $y = c_1\cos x + c_2\sin x + \sin x\log(1 + \sin x) - x\cos x - 1.$

31. $y = c_1\cos ax + c_2\sin ax - \frac{1}{a^2}\cos ax\log|\sec ax + \tan ax|.$

32. $y = c_1\cos x + c_2\sin x + \frac{1}{2}x\sin x + \frac{1}{4}\cos x.$

33. $y = c_1x + \frac{c_2}{x+1} + \frac{x^2(4x+3)}{6(x+1)}.$

34. $y = x(c_1 + c_2e^x) - x(x+1).$

35. $y = (c_1 + c_2\log x)x.$

36. $y = (c_1 + c_2\log x)x^2 + x^2(\log x)^2.$

37. $y = x^{-2}(c_1 + c_2\log x) + \frac{x}{9}\left(\log x - \frac{2}{3}\right)$

38. $y = \frac{c_1}{x} + c_2x^4 - \frac{x^2}{6} - \frac{1}{2}\log x + \frac{3}{8}.$

39. $y = x(c_1 + c_2\log x) + \log x + 2.$

40. $y = c_1\cos\{\log(1+x)\} + c_2\sin\{\log(1+x)\} + 2\log(1+x)\sin\{\log(1+x)\}.$

41. $y = c_1\cos 2t + c_2\sin 2t - \frac{1}{2}t\cos^2 t$, where $t = \log(2+x)$.

42. $x = c_1\cos t + c_2\sin t, y = -\frac{1}{2}(c_1 + 3c_2)\sin t + \frac{1}{2}(c_2 - 3c_1)\cos t.$

43. $x = c_1e^{-2t} + c_2e^{-7t} - \frac{31}{196} + \frac{5}{14}t - \frac{1}{8}e^t,\ y = -\frac{2}{3}c_1e^{-2t} + c_2e^{-7t} + \frac{9}{98} - \frac{1}{7}t + \frac{5}{24}e^t.$

44. $x = c_1e^{-\frac{1}{2}t} + c_2e^{-\frac{1}{4}t} + \frac{1}{15}e^t,\ y = c_1e^{-\frac{1}{2}t} - c_2e^{-\frac{1}{4}t} + \frac{4}{15}e^t.$

45. $x = \cos 2t - \sin 2t - \cos t,\ y = \cos 2t + \sin 2t - \sin t.$

46. $x = c_1e^{\sqrt{2}t} + c_2e^{-\sqrt{2}t} + 3\cos t,\ y = (\sqrt{2}+1)c_1e^{\sqrt{2}t} - (\sqrt{2}-1)c_2e^{-\sqrt{2}t} + 2\sin t + c_3.$

47. $x = (c_1 + c_2x)e^{3x},\ y = (1 - 2x)(c_2 - 2c_1)e^{3x}.$

48. $x = e^t + e^{-t},\ y = e^{-t} - e^t + \sin t.$

49. $x = c_1e^t + c_2e^{-5t} + \frac{6}{7}e^{2t},\ y = c_2e^{-5t} - c_1e^t + \frac{8}{7}e^{2t}.$

50. $x = \frac{1}{5}e^t + \frac{2}{5}e^{-t} - c_1\sin 2t + c_2\cos 2t,\ y = \frac{2}{5}e^t + \frac{1}{5}e^{-t} + c_1\cos 2t + c_2\sin 2t.$

51. $x = c_1e^t + c_2e^{-5t} + \frac{6}{7}e^{2t} - 2t - \frac{13}{5},\ y = c_2e^t - c_1e^{-5t} - \frac{8}{7}e^{2t} - 3t - \frac{12}{5}.$

52. $x = c_1e^{-t} + c_2e^{3t} - \frac{1}{5}(\cos t - 2\sin t),\ y = 2c_1e^{-t} - 2c_2e^{3t} + \frac{1}{5}(\sin t + 2\cos t).$