## **DNA** fingerprinting

= a process for quantifying the similarity of the DNA of 2 or more individuals. Step 1. Chop up the DNA into pieces using an enzyme.

ACCGTGTGAACTTACGTGTGC TGGCACACTTGAATGCACACG

Step 1. Chop up the DNA into pieces using an enzyme.

ACCGTGTGAACTTACGTGTGC TGGCACACTTGAATGCACACG

The enzyme recognizes a particular sequence of bases:

GTGTG

Step 1. Chop up the DNA into pieces using an enzyme.

ACC<mark>GTGTGAACTTACGTGTGC</mark> TGGCACAC<mark>TTGAATGCACACG</mark>

The enzyme recognizes a particular sequence of bases:

GTGTG CACAC

Step 1. Chop up the DNA into pieces using an enzyme.

ACC<mark>GTGTGAACTTAC</mark>GTGTGC TGGCACAC<mark>TTGAATGCACAC</mark>G

The enzyme recognizes a particular sequence of bases:


GTGTG CACAC Step 1. Chop up the DNA into pieces using an enzyme.

ACC<mark>GTGTGAACTTAC</mark>GTGTGC TGGCACAC<mark>TTGAATGCACAC</mark>G

ACC

GTGTGAACTTAC

GTGTGC



ACCGTGTGAACTTACGCGTGC
TGGCACACTTGAATGCACACG
ACCGTGTGAACTTACGCGTGC
TGGCACACTTGAATGCGCGTGC
TGGTACACTTGAATGCGCGTGC

ACCGTGTGAACTTACGTGTGC
TGGCACACTTGAATGCACACG

ACCGTGTGAACTTACGCGTGC
TGGCACACTTGAATGCGCACG

ACCATGTGAACTTACGCGTGC
TGGTACACTTGAATGCGCACG

ACCGTGTGAACTTACGTGTGC
TGGCACACTTGAATGCACACG

ACCGTGTGAACTTACGCGTGC
TGGCACACTTGAATGCGCACG

mutation

ACCATGTGAACTTACGCGTGC
TGGTACACTTGAATGCGCACG

ACCGTGTGAACTTACGTGTGC
TGGCACACTTGAATGCACACG

ACCGTGTGAACTTACGCGTGC
TGGCACACTTGAATGCGCACG
mutation

ACCATGTGAACTTACGCGTGC
TGGTACACTTGAATGCGCACG
mutation

ACCGTGTGAACTTACGCGTGC
TGGCACACTTGAATGCACACG


ACCGTGTGAACTTACGCGTGC
TGGCACACTTGAATGCGCACG

mutation

ACCATGTGAACTTACGCGTGC
TGGTACACTTGAATGCGCACG

mutation


mutation




The more similar the DNA is between 2 individuals, the more similar will be the sizes of the fragments of DNA.


The more similar the DNA is between 2 individuals, the more similar will be the sizes of the fragments of DNA.

DNA.			8
1	2	3	4
3	3	3	3 -
18	15	12	18 —
20	20	20	20
20	20	22	20 —
36	36	35	36 ——
42	42	42	42 ———
44	44	49	44 ———
60	60	60	60 —
63	66	<b>70</b>	63







Step 2. Put the DNA fragments on a gel.





Step 3. Apply electric current to the gel.


- causes the fragments to move down the gel.
- small fragments move faster than large fragments.















1	2	3	4
3	3	3	3
18	15	12	18
20	20	20	20
20	20	22	20
36	36	35	36
42	42	42	42
44	44	49	44
60	60	60	60
63	66	<b>70</b>	63



## **Step 4. Compare the bands**

- The more bands two individuals have in common, the more similar their DNA is.
- If two columns show identical banding patterns, the DNA came from the same individual (or a twin).

