

SOLVING EQUATIONS

QUANTITATIVE ECONOMICS 2024

Piotr Żoch

October 30, 2024

INTRODUCTION

- Systems of linear equations.
- Nonlinear equations (to be added)

LINEAR SYSTEMS OF EQUATIONS

- One of the most common problems in scientific computation: solve

$$\mathbf{Ax} = \mathbf{b},$$

for \mathbf{x} , where \mathbf{A} is a square matrix and \mathbf{b} is a vector.

- Seems like an easy problem, but it will teach us many things.
- Multiple specialized libraries for numerical linear algebra: LAPACK, BLAS, IMKL...

DIRECT METHODS

- Elementary operations:
 - multiply a row by a scalar,
 - add a scalar multiple of a row to another row,
 - interchange two rows.
- Solve $\mathbf{Ax} = \mathbf{b}$ by using elementary row operations on the augmented matrix $[\mathbf{A} \mid \mathbf{b}]$.
- Transform \mathbf{A} into a reduced row echelon form.

DIRECT METHODS

- Two step procedure:
 - Forward elimination.

$$\left[\begin{array}{cccc|c} * & * & * & * & * \\ * & * & * & * & * \\ * & * & * & * & * \\ * & * & * & * & * \end{array} \right] \rightarrow \left[\begin{array}{cccc|c} * & * & * & * & * \\ & * & * & * & * \\ & * & * & * & * \\ & * & * & * & * \end{array} \right] \rightarrow \left[\begin{array}{cccc|c} * & * & * & * & * \\ & * & * & * & * \\ & & * & * & * \\ & & * & * & * \end{array} \right] \rightarrow \left[\begin{array}{cccc|c} * & * & * & * & * \\ & * & * & * & * \\ & & * & * & * \\ & & & * & * \end{array} \right]$$

- Backward elimination.

$$\left[\begin{array}{cccc|c} * & * & * & * & * \\ & * & * & * & * \\ & & * & * & * \\ & & & * & * \end{array} \right] \rightarrow \left[\begin{array}{cccc|c} * & * & * & * & * \\ & * & * & * & * \\ & & * & * & * \\ & & & * & * \end{array} \right] \rightarrow \left[\begin{array}{cccc|c} * & * & * & * & * \\ & * & * & * & * \\ & & * & * & * \\ & & & * & * \end{array} \right] \rightarrow \left[\begin{array}{cccc|c} * & * & * & * & * \\ & * & * & * & * \\ & & * & * & * \\ & & & * & * \end{array} \right]$$

FLOATING POINT NUMBERS

- Forward elimination: to deal with the first column we need n^2 operations, for the second $n^2 - 1$, for the third $n^2 - 2$ and so on.
- Backward elimination: to deal with the last column we need n operations, for the second to last $n - 1$, for the third to last $n - 2$ and so on.
- Forward elimination is $\mathcal{O}(n^3)$, backward elimination is $\mathcal{O}(n^2)$.

TRIANGULAR SYSTEMS

- Lower triangular system:

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ l_{21} & 1 & 0 & 0 \\ l_{31} & l_{32} & 1 & 0 \\ l_{41} & l_{42} & l_{43} & 1 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \end{bmatrix}$$

can be solved using forward elimination, starting from the top

$$y_i = b_i - \sum_{j=1}^{i-1} l_{ij} y_j.$$

UPPER TRIANGULAR SYSTEMS

- Upper triangular system:

$$\begin{bmatrix} u_{11} & u_{12} & u_{13} & u_{14} \\ 0 & u_{22} & u_{23} & u_{24} \\ 0 & 0 & u_{33} & u_{34} \\ 0 & 0 & 0 & u_{44} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \end{bmatrix}$$

can be solved using backward elimination, starting from the bottom

$$x_i = \frac{1}{u_{ii}} \left(y_i - \sum_{j=i+1}^n u_{ij} x_j \right).$$

- Suppose we can write

$$\mathbf{A} = \mathbf{LU}$$

where \mathbf{L} is a lower triangular matrix and \mathbf{U} is an upper triangular matrix.

- We have

$$\mathbf{L}(\mathbf{Ux}) = \mathbf{b} \rightarrow \mathbf{Ly} = \mathbf{b}$$

which we can solve for \mathbf{y} using forward elimination.

- We then solve

$$\mathbf{Ux} = \mathbf{y}$$

for \mathbf{x} using backward elimination.

- This is known as Gaussian elimination.
 1. Compute \mathbf{L} and \mathbf{U} .
 2. Solve $\mathbf{L}\mathbf{y} = \mathbf{b}$ for \mathbf{y} using forward elimination.
 3. Solve $\mathbf{U}\mathbf{x} = \mathbf{y}$ for \mathbf{x} using backward elimination.
- Step 1. is known as **LU decomposition**.
- Nice thing: we can keep \mathbf{L} and \mathbf{U} and recycle them for different \mathbf{b} .
- How to perform the decomposition?

MATRIX MULTIPLICATION BY OUTER PRODUCTS

- Write the columns of **A** as $\mathbf{a}_1, \dots, \mathbf{a}_n$.
- Write the rows of **B** as $\mathbf{b}_1^\top, \dots, \mathbf{b}_n^\top$.
- We have

$$\mathbf{AB} = \sum_{k=1}^n \mathbf{a}_k \mathbf{b}_k^\top.$$

- Useful: for triangular matrices **L**, **U** only the first outer product contributes to the first row and the first column of **LU**

$$\mathbf{e}_1^\top \sum_{k=1}^n \mathbf{l}_k \mathbf{u}_k^\top = l_{11} \mathbf{u}_1^\top, \quad \left(\sum_{k=1}^n \mathbf{l}_k \mathbf{u}_k^\top \right) \mathbf{e}_1 = u_{11} \mathbf{l}_1.$$

LU FACTORIZATION WITHOUT PIVOTING

Algorithm LU Factorization without Pivoting

Require: Matrix $\mathbf{A} \in \mathbb{R}^{n \times n}$

```
1: for  $j = 1$  to  $n$  do  
2:   for  $i = j + 1$  to  $n$  do  
3:      $a_{ij} = \frac{a_{ij}}{a_{jj}}$   
4:   for  $k = j + 1$  to  $n$  do  
5:      $a_{ik} = a_{ik} - a_{ij}a_{jk}$   
6:   end for  
7: end for  
8: end for
```

- This algorithm uses \mathbf{A} matrix to store \mathbf{L} and \mathbf{U} .
- Problem when $a_{jj} = 0$ at any step -

SOLVING THE SYSTEM

- We need

$$\sum_{j=1}^n \sum_{i=j+1}^n \left(1 + \sum_{k=j+1}^n 2 \right) = \frac{2}{3}n^3 - \frac{1}{2}n^2 - \frac{1}{6}n$$

operations to perform the factorization.

- We then need

$$\sum_{i=1}^n \left(2 + \sum_{j=1}^{i-1} 2 \right) = n^2 + n$$

operations for forward and backward substitution each.

- LU decomposition is the most costly step.

SOLVING THE SYSTEM

- We need

$$\sum_{j=1}^n \sum_{i=j+1}^n \left(1 + \sum_{k=j+1}^n 2 \right) = \frac{2}{3}n^3 - \frac{1}{2}n^2 - \frac{1}{6}n$$

operations to perform the factorization.

- We then need

$$\sum_{i=1}^n \left(2 + \sum_{j=1}^{i-1} 2 \right) = n^2 + n$$

operations for forward and backward substitution each.

- LU decomposition is the most costly step.

SOLVING THE SYSTEM

- In practice we use a similar method, but with **pivoting**
- PLU factorization:

$$\tilde{\mathbf{A}} = \mathbf{L}\mathbf{U},$$

where $\tilde{\mathbf{A}}$ is a matrix \mathbf{A} with its rows permuted.

- It works if and only if \mathbf{A} is non-singular.
- Asymptotically uses the same number of operations and LU without pivoting.

NORMS

- A **vector norm** is a function $\|\cdot\| : \mathbb{R}^n \rightarrow \mathbb{R}$ that satisfies:

1. $\|\mathbf{x}\| \geq 0$,
2. $\|\mathbf{x}\| = 0 \iff \mathbf{x} = \mathbf{0}$,
3. $\|a\mathbf{x}\| = |a| \|\mathbf{x}\|$,
4. $\|\mathbf{x} + \mathbf{y}\| \leq \|\mathbf{x}\| + \|\mathbf{y}\|$,

for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ and $a \in \mathbb{R}$.

- Common vector norms: $\ell_1, \ell_2, \ell_\infty$:

$$\|\mathbf{x}\|_1 = \sum_{i=1}^n |x_i|, \quad \|\mathbf{x}\|_2 = \sqrt{\sum_{i=1}^n x_i^2}, \quad \|\mathbf{x}\|_\infty = \max_{i=1, \dots, n} |x_i|.$$

NORMS

- For matrices we have **matrix norms**.
- A **Frobenius** norm is

$$\|\mathbf{A}\|_F = \sqrt{\sum_{i=1}^m \sum_{j=1}^n a_{ij}^2}.$$

- Imagine representing a matrix as a vector with columns stacked on top of each other.
- An **induced** matrix norm is

$$\|\mathbf{A}\|_p = \max_{\|\mathbf{x}\|_p=1} \|\mathbf{Ax}\|_p.$$

- In Julia: `norm(A)` is the Frobenius norm, `opnorm(A, p)` is the induced norm.

NORMS

- We have
 1. $\|\mathbf{Ax}\| \leq \|\mathbf{A}\| \|\mathbf{x}\|$,
 2. $\|\mathbf{AB}\| \leq \|\mathbf{A}\| \|\mathbf{B}\|$,
 3. for a square matrix, $\|\mathbf{A}^k\| \leq \|\mathbf{A}\|^k$ for any integer $k \geq 0$.
- Two common matrix norm are the 1-norm and the ∞ -norm:

$$\|\mathbf{A}\|_1 = \max_{j=1,\dots,n} \sum_{i=1}^m |a_{ij}|, \quad \|\mathbf{A}\|_\infty = \max_{i=1,\dots,m} \sum_{j=1}^n |a_{ij}|.$$

CONDITIONING OF LINEAR SYSTEMS

- Consider the perturbed system

$$\mathbf{A}(\mathbf{x} + \mathbf{h}) = \mathbf{b} + \mathbf{d}.$$

- The condition number is the relative change in the solution divided by the relative change in the data:

$$\kappa = \frac{\|\mathbf{h}\| / \|\mathbf{x}\|}{\|\mathbf{d}\| / \|\mathbf{b}\|} = \frac{\|\mathbf{h}\| \|\mathbf{b}\|}{\|\mathbf{d}\| \|\mathbf{x}\|}.$$

- Note that $\mathbf{h} = \mathbf{A}^{-1}\mathbf{d}$ so

$$\|\mathbf{h}\| \leq \|\mathbf{A}^{-1}\| \|\mathbf{d}\|.$$

CONDITIONING OF LINEAR SYSTEMS

- Use $\|\mathbf{h}\| \leq \|\mathbf{A}^{-1}\| \|\mathbf{d}\|$ to write

$$\frac{\|\mathbf{h}\| \|\mathbf{b}\|}{\|\mathbf{d}\| \|\mathbf{x}\|} \leq \frac{\|\mathbf{A}^{-1}\| \|\mathbf{d}\| \|\mathbf{A}\| \|\mathbf{x}\|}{\|\mathbf{d}\| \|\mathbf{x}\|} = \|\mathbf{A}^{-1}\| \|\mathbf{A}\| .$$

- We can prove that inequality is tight.
- The matrix **condition number** of an invertible square matrix \mathbf{A} is

$$\kappa(\mathbf{A}) = \|\mathbf{A}^{-1}\| \|\mathbf{A}\| .$$

CONDITIONING OF LINEAR SYSTEMS

- If $\mathbf{A}(\mathbf{x} + \Delta\mathbf{x}) = \mathbf{b} + \Delta\mathbf{b}$ then

$$\frac{\|\Delta\mathbf{x}\|}{\|\mathbf{x}\|} \leq \kappa(\mathbf{A}) \frac{\|\Delta\mathbf{b}\|}{\|\mathbf{b}\|}.$$

- The condition number is a measure of how sensitive the solution is to changes in the data.
- We can derive a similar result for perturbed \mathbf{A} .
- The condition number is at least equal to 1.
- A condition number of 10^k means that we lose k digits of precision.

CONDITIONING OF LINEAR SYSTEMS

- Suppose that we compute a "solution" $\tilde{\mathbf{x}}$ to the system $\mathbf{Ax} = \mathbf{b}$.
- We would like to compare $\tilde{\mathbf{x}}$ to the true solution \mathbf{x} - but we do not know \mathbf{x} .
- We can calculate the residual

$$\mathbf{r} = \mathbf{b} - \mathbf{A}\tilde{\mathbf{x}}$$

-
- We have

$$\frac{\|\mathbf{x} - \tilde{\mathbf{x}}\|}{\|\mathbf{x}\|} \leq \kappa(\mathbf{A}) \frac{\|\mathbf{r}\|}{\|\mathbf{b}\|}.$$

ITERATIVE METHODS

- We saw that Gaussian elimination is $\mathcal{O}(n^3)$.
- This is prohibitive for large n (unless a matrix has a special structure).
- But matrix-vector multiplication is $\mathcal{O}(n^2)$.
- In some cases we can apply repeated matrix-vector multiplication to solve the system.
- We call these **iterative methods**.

JACOBI METHOD

- Suppose we want to solve $\mathbf{Ax} = \mathbf{b}$.
- This can be written as

$$\sum_{j=1}^n a_{ij}x_j = b_i, \quad \text{or} \quad x_i = \frac{1}{a_{ii}} \left(b_i - \sum_{j \neq i} a_{ij}x_j \right).$$

- Start from some initial $\mathbf{x}^{(0)}$ and iterate:

$$x_i^{(k+1)} = \frac{1}{a_{ii}} \left(b_i - \sum_{j \neq i} a_{ij}x_j^{(k)} \right)$$

until $\mathbf{x}^{(k+1)}$ is close enough to $\mathbf{x}^{(k)}$.

GAUSS-SEIDEL METHOD

- We can also write

$$x_i = \frac{1}{a_{ii}} \left(b_i - \sum_{j=1}^{i-1} a_{ij}x_j - \sum_{j=i+1}^n a_{ij}x_j \right).$$

- The Gauss-Seidel method uses the newest values of x_j :

$$x_i^{(k+1)} = \frac{1}{a_{ii}} \left(b_i - \sum_{j=1}^{i-1} a_{ij}x_j^{(k+1)} - \sum_{j=i+1}^n a_{ij}x_j^{(k)} \right).$$

ITERATIVE METHODS

- Rewrite $\mathbf{A} = \mathbf{P} - \mathbf{N}$ so that the system is

$$\mathbf{P}\mathbf{x} = \mathbf{N}\mathbf{x} + \mathbf{b}.$$

- An iterative method is

$$\mathbf{P}\mathbf{x}^{(k+1)} = (\mathbf{N}\mathbf{x}^{(k)} + \mathbf{b}).$$

- \mathbf{P} is called a **preconditioner**
- Jacobi and Gauss-Seidel differ in the choice of \mathbf{P} .
- Since $\mathbf{x}^{k+1} = \mathbf{P}^{-1} (\mathbf{N}\mathbf{x}^{(k)} + \mathbf{b})$ a good preconditioner should be easy to invert.

ITERATIVE METHODS

- Why do we call \mathbf{P} a preconditioner?
- Suppose we have a system $\mathbf{Ax} = \mathbf{b}$.
- The condition number is $\kappa(\mathbf{A})$.
- Suppose we have a preconditioned system $\mathbf{P}^{-1}\mathbf{Ax} = \mathbf{P}^{-1}\mathbf{b}$.
- The condition number is $\kappa(\mathbf{P}^{-1}\mathbf{A})$.
- If $\mathbf{P} \approx \mathbf{A}^{-1}$, then $\kappa(\mathbf{P}^{-1}\mathbf{A}) \approx 1$.

ITERATIVE METHODS

- Iterative methods do not always converge.
- They are guaranteed to converge if **A** is **diagonally dominant**:

$$|a_{ii}| > \sum_{j \neq i} |a_{ij}|.$$

- There are better methods than Jacobi and Gauss-Seidel - the idea is to choose **P** so that the convergence is faster.