

# **Optimization applied 1:**

## **Maximum likelihood estimation**

---

Marcin Lewandowski

November 26, 2024

## MLE: Intuition (Bernoulli)

Let  $Y \sim \text{Bern}(p)$ .

- Suppose we observe 6 i.i.d coin flips (1 for heads, 0 for tails):

$$\mathbf{y} = \{1, 1, 1, 0, 1, 0\}$$

- Denote each flip as  $y_1, y_2, \dots, y_6$ , where  $y_1 = 1$  and  $y_6 = 1$ .

## MLE: Intuition (Bernoulli)

Let  $Y \sim \text{Bern}(p)$ .

- Suppose we observe 6 i.i.d coin flips (1 for heads, 0 for tails):

$$\mathbf{y} = \{1, 1, 1, 0, 1, 0\}$$

- Denote each flip as  $y_1, y_2, \dots, y_6$ , where  $y_1 = 1$  and  $y_6 = 1$ .

Recall: the probability mass function (PMF) of a Bernoulli r.v. is:

$$f(y; p) = p^y(1 - p)^{1-y}, \quad y \in \{0, 1\}$$

The probability of 1 is  $p$ , while the probability of 0 is  $(1 - p)$ .

## MLE: Intuition (Bernoulli)

Let  $Y \sim \text{Bern}(p)$ .

- Suppose we observe 6 i.i.d coin flips (1 for heads, 0 for tails):

$$\mathbf{y} = \{1, 1, 1, 0, 1, 0\}$$

- Denote each flip as  $y_1, y_2, \dots, y_6$ , where  $y_1 = 1$  and  $y_6 = 1$ .

Recall: the probability mass function (PMF) of a Bernoulli r.v. is:

$$f(y; p) = p^y(1 - p)^{1-y}, \quad y \in \{0, 1\}$$

The probability of 1 is  $p$ , while the probability of 0 is  $(1 - p)$ .

**Overarching Goal:** Estimate  $p$  given a vector of observed data.

## MLE: Intuition (Bernoulli)


- $y_i$  are independent
- The likelihood of observing  $\mathbf{y} = \{1, 1, 1, 0, 1, 0\}$  is:

$$L(p \mid \mathbf{y}) = \prod_{i=1}^6 p^{y_i} (1 - p)^{1-y_i}$$

# MLE: Intuition (Bernoulli)

- $y_i$  are independent
- The likelihood of observing  $\mathbf{y} = \{1, 1, 1, 0, 1, 0\}$  is:

$$L(p | \mathbf{y}) = \prod_{i=1}^6 p^{y_i} (1-p)^{1-y_i}$$



## MLE: Log likelihood function (Bernoulli)

$$L(p \mid \mathbf{y}) = \prod_{i=1}^n p^{y_i} \cdot (1 - p)^{1-y_i}$$

Necessary condition for maxima:

## MLE: Log likelihood function (Bernoulli)

$$L(p \mid \mathbf{y}) = \prod_{i=1}^n p^{y_i} \cdot (1 - p)^{1-y_i}$$

$$\log(L(p \mid \mathbf{y})) = \sum_{i=1}^n \log(p^{y_i}) + \sum_{i=1}^n \log(1 - p^{1-y_i})$$

Necessary condition for maxima:



## MLE: Log likelihood function (Bernoulli)

$$\begin{aligned} L(p \mid \mathbf{y}) &= \prod_{i=1}^n p^{y_i} \cdot (1-p)^{1-y_i} \\ \log(L(p \mid \mathbf{y})) &= \sum_{i=1}^n \log(p^{y_i}) + \sum_{i=1}^n \log(1-p^{1-y_i}) \\ &= \sum_{i=1}^n y_i \log(p) + \sum_{i=1}^n (1-y_i) \log(1-p) \end{aligned}$$

Necessary condition for maxima:

## MLE: Log likelihood function (Bernoulli)

$$\begin{aligned} L(p \mid \mathbf{y}) &= \prod_{i=1}^n p^{y_i} \cdot (1-p)^{1-y_i} \\ \log(L(p \mid \mathbf{y})) &= \sum_{i=1}^n \log(p^{y_i}) + \sum_{i=1}^n \log(1-p^{1-y_i}) \\ &= \sum_{i=1}^n y_i \log(p) + \sum_{i=1}^n (1-y_i) \log(1-p) \\ &= \log(p) \sum_{i=1}^n y_i + \log(1-p) \sum_{i=1}^n (1-y_i) \end{aligned}$$

Necessary condition for maxima:

## MLE: Log likelihood function (Bernoulli)

$$L(p | \mathbf{y}) = \prod_{i=1}^n p^{y_i} \cdot (1 - p)^{1-y_i}$$

$$\begin{aligned}\log(L(p | \mathbf{y})) &= \sum_{i=1}^n \log(p^{y_i}) + \sum_{i=1}^n \log(1 - p^{1-y_i}) \\ &= \sum_{i=1}^n y_i \log(p) + \sum_{i=1}^n (1 - y_i) \log(1 - p) \\ &= \log(p) \sum_{i=1}^n y_i + \log(1 - p) \sum_{i=1}^n (1 - y_i)\end{aligned}$$

Necessary condition for maxima:

$$\frac{\partial \log(L(p | \mathbf{y}))}{\partial p} = 0$$

## MLE: Log likelihood function (Bernoulli)

$$L(p | \mathbf{y}) = \prod_{i=1}^n p^{y_i} \cdot (1-p)^{1-y_i}$$

$$\begin{aligned}\log(L(p | \mathbf{y})) &= \sum_{i=1}^n \log(p^{y_i}) + \sum_{i=1}^n \log(1-p^{1-y_i}) \\ &= \sum_{i=1}^n y_i \log(p) + \sum_{i=1}^n (1-y_i) \log(1-p) \\ &= \log(p) \sum_{i=1}^n y_i + \log(1-p) \sum_{i=1}^n (1-y_i)\end{aligned}$$

Necessary condition for maxima:

$$\begin{aligned}\frac{\partial \log(L(p | \mathbf{y}))}{\partial p} &= 0 \\ \frac{\sum_{i=1}^n y_i}{p} - \frac{\sum_{i=1}^n (1-y_i)}{1-p} &= 0\end{aligned}$$

## MLE: Log likelihood function (Bernoulli) maximized

**Note:** In this simple example we can solve for  $p$ !

$$\frac{\partial \log (L(p \mid \mathbf{y}))}{\partial p} = 0$$

## MLE: Log likelihood function (Bernoulli) maximized

**Note:** In this simple example we can solve for  $p$ !

$$\frac{\partial \log(L(p | \mathbf{y}))}{\partial p} = 0$$
$$\frac{\sum_{i=1}^n y_i}{p} = \frac{\sum_{i=1}^n (1 - y_i)}{1 - p}$$

## MLE: Log likelihood function (Bernoulli) maximized

**Note:** In this simple example we can solve for  $p$ !

$$\frac{\partial \log(L(p | \mathbf{y}))}{\partial p} = 0$$

$$\frac{\sum_{i=1}^n y_i}{p} = \frac{\sum_{i=1}^n (1 - y_i)}{1 - p}$$

Using:  $y = \{1, 1, 1, 0, 1, 0\}$

## MLE: Log likelihood function (Bernoulli) maximized

**Note:** In this simple example we can solve for  $p$ !

$$\frac{\partial \log(L(p | \mathbf{y}))}{\partial p} = 0$$

$$\frac{\sum_{i=1}^n y_i}{p} = \frac{\sum_{i=1}^n (1 - y_i)}{1 - p}$$

Using:  $y = \{1, 1, 1, 0, 1, 0\}$

$$\frac{4}{p} = \frac{2}{1 - p}$$



## MLE: Log likelihood function (Bernoulli) maximized

**Note:** In this simple example we can solve for  $p$ !

$$\frac{\partial \log(L(p | \mathbf{y}))}{\partial p} = 0$$

$$\frac{\sum_{i=1}^n y_i}{p} = \frac{\sum_{i=1}^n (1 - y_i)}{1 - p}$$

Using:  $y = \{1, 1, 1, 0, 1, 0\}$

$$\frac{4}{p} = \frac{2}{1 - p}$$

$$4 - 4p = 2p$$

## MLE: Log likelihood function (Bernoulli) maximized

**Note:** In this simple example we can solve for  $p$ !

$$\frac{\partial \log(L(p | \mathbf{y}))}{\partial p} = 0$$

$$\frac{\sum_{i=1}^n y_i}{p} = \frac{\sum_{i=1}^n (1 - y_i)}{1 - p}$$

Using:  $y = \{1, 1, 1, 0, 1, 0\}$

$$\frac{4}{p} = \frac{2}{1 - p}$$

$$4 - 4p = 2p$$

$$-6p = -4$$

## MLE: Log likelihood function (Bernoulli) maximized

**Note:** In this simple example we can solve for  $p$ !

$$\frac{\partial \log(L(p | \mathbf{y}))}{\partial p} = 0$$

$$\frac{\sum_{i=1}^n y_i}{p} = \frac{\sum_{i=1}^n (1 - y_i)}{1 - p}$$

Using:  $y = \{1, 1, 1, 0, 1, 0\}$

$$\frac{4}{p} = \frac{2}{1 - p}$$

$$4 - 4p = 2p$$

$$-6p = -4$$

$$\hat{p} = \frac{4}{6}$$

## MLE: Log likelihood function (Bernoulli) maximized

**Note:** In this simple example we can solve for  $p$ !

$$\frac{\partial \log(L(p | \mathbf{y}))}{\partial p} = 0$$

$$\frac{\sum_{i=1}^n y_i}{p} = \frac{\sum_{i=1}^n (1 - y_i)}{1 - p}$$

Using:  $y = \{1, 1, 1, 0, 1, 0\}$

$$\frac{4}{p} = \frac{2}{1 - p}$$

$$4 - 4p = 2p$$


$$-6p = -4$$

$$\hat{p} = \frac{4}{6}$$

Let's see how to implement MLE in Julia!


# A task for you: the Poisson Distribution

$$f(y | \lambda) = \frac{e^{-\lambda} \lambda^y}{y!} ; y \in \{0, 1, 2, 3, \dots\}$$



# The Poisson Distribution

$$f(y | \lambda) = \frac{e^{-\lambda} \lambda^y}{y!} ; y \in \{0, 1, 2, 3, \dots\}$$



## Poisson Distribution: Log-Likelihood Function

$$L(\lambda \mid \mathbf{y}) = \prod_{i=1}^n \frac{e^{-\lambda} \lambda^{y_i}}{y_i!}$$

## Poisson Distribution: Log-Likelihood Function

$$L(\lambda \mid \mathbf{y}) = \prod_{i=1}^n \frac{e^{-\lambda} \lambda^{y_i}}{y_i!}$$

$$\log(L(\lambda \mid \mathbf{y})) = \sum_{i=1}^n \log\left(\frac{e^{-\lambda} \lambda^{y_i}}{y_i!}\right)$$



## Poisson Distribution: Log-Likelihood Function

$$\begin{aligned} L(\lambda \mid \mathbf{y}) &= \prod_{i=1}^n \frac{e^{-\lambda} \lambda^{y_i}}{y_i!} \\ \log(L(\lambda \mid \mathbf{y})) &= \sum_{i=1}^n \log\left(\frac{e^{-\lambda} \lambda^{y_i}}{y_i!}\right) \\ &= -\sum_{i=1}^n \lambda + \log(\lambda) \sum_{i=1}^n y_i - \sum_{i=1}^n \log(y_i!) \end{aligned}$$

## Poisson Distribution: Log-Likelihood Function

$$\begin{aligned}L(\lambda \mid \mathbf{y}) &= \prod_{i=1}^n \frac{e^{-\lambda} \lambda^{y_i}}{y_i!} \\ \log(L(\lambda \mid \mathbf{y})) &= \sum_{i=1}^n \log\left(\frac{e^{-\lambda} \lambda^{y_i}}{y_i!}\right) \\ &= -\sum_{i=1}^n \lambda + \log(\lambda) \sum_{i=1}^n y_i - \sum_{i=1}^n \log(y_i!) \\ &= -n\lambda + \log(\lambda) \sum_{i=1}^n y_i - \sum_{i=1}^n \log(y_i!)\end{aligned}$$

## Poisson Distribution: Log-Likelihood Function

$$\begin{aligned}L(\lambda \mid \mathbf{y}) &= \prod_{i=1}^n \frac{e^{-\lambda} \lambda^{y_i}}{y_i!} \\ \log(L(\lambda \mid \mathbf{y})) &= \sum_{i=1}^n \log\left(\frac{e^{-\lambda} \lambda^{y_i}}{y_i!}\right) \\ &= -\sum_{i=1}^n \lambda + \log(\lambda) \sum_{i=1}^n y_i - \sum_{i=1}^n \log(y_i!) \\ &= -n\lambda + \log(\lambda) \sum_{i=1}^n y_i - \sum_{i=1}^n \log(y_i!)\end{aligned}$$

**Note:**  $\sum_{i=1}^n \log(y_i!)$  does not depend on  $\lambda$

## Poisson Distribution: Log-Likelihood Function

$$\begin{aligned}L(\lambda \mid \mathbf{y}) &= \prod_{i=1}^n \frac{e^{-\lambda} \lambda^{y_i}}{y_i!} \\ \log(L(\lambda \mid \mathbf{y})) &= \sum_{i=1}^n \log\left(\frac{e^{-\lambda} \lambda^{y_i}}{y_i!}\right) \\ &= -\sum_{i=1}^n \lambda + \log(\lambda) \sum_{i=1}^n y_i - \sum_{i=1}^n \log(y_i!) \\ &= -n\lambda + \log(\lambda) \sum_{i=1}^n y_i - \sum_{i=1}^n \log(y_i!)\end{aligned}$$

**Note:**  $\sum_{i=1}^n \log(y_i!)$  does not depend on  $\lambda$

$$\frac{\partial \log(L(\lambda \mid \mathbf{y}))}{\partial \lambda} = -n + \frac{\sum_{i=1}^n y_i}{\lambda}$$

## Poisson Distribution: Log-Likelihood Function

$$\begin{aligned}L(\lambda \mid \mathbf{y}) &= \prod_{i=1}^n \frac{e^{-\lambda} \lambda^{y_i}}{y_i!} \\ \log(L(\lambda \mid \mathbf{y})) &= \sum_{i=1}^n \log\left(\frac{e^{-\lambda} \lambda^{y_i}}{y_i!}\right) \\ &= -\sum_{i=1}^n \lambda + \log(\lambda) \sum_{i=1}^n y_i - \sum_{i=1}^n \log(y_i!) \\ &= -n\lambda + \log(\lambda) \sum_{i=1}^n y_i - \sum_{i=1}^n \log(y_i!)\end{aligned}$$

**Note:**  $\sum_{i=1}^n \log(y_i!)$  does not depend on  $\lambda$

$$\begin{aligned}\frac{\partial \log(L(\lambda \mid \mathbf{y}))}{\partial \lambda} &= -n + \frac{\sum_{i=1}^n y_i}{\lambda} \\ \frac{\partial^2 \log(L(\lambda \mid \mathbf{y}))}{\partial \lambda^2} &= -\frac{\sum_{i=1}^n y_i}{\lambda^2}\end{aligned}$$

## The variance of the MLE estimator

The estimator of the MLE variance is:

$$\begin{aligned}\widehat{Var}(\hat{\lambda}) &= \left\{ -\frac{\partial^2 \log(L(\lambda | \mathbf{y}))}{\partial \lambda^2} \right\}^{-1} \\ &= \left\{ \frac{\sum_{i=1}^n y_i}{\lambda^2} \right\}^{-1}\end{aligned}$$

# The Logit Model:


We want to model the probability of a binary outcome,  $Y \in \{0, 1\}$

- Assume that  $P(Y = y_i | x)$  depends on a single predictor,  $x$ .
- Thus,  $P(Y = y_i | x_i) = F(\beta_0 + \beta_1 x)$

# The Logit Model:

We want to model the probability of a binary outcome,  $Y \in \{0, 1\}$

- Assume that  $P(Y = y_i | x)$  depends on a single predictor,  $x$ .
- Thus,  $P(Y = y_i | x_i) = F(\beta_0 + \beta_1 x)$





## The Logit Model:

$$\begin{aligned} P(y = 1 \mid x) &= F(\beta_0 + \beta_1 x) \\ &= \frac{e^{\beta_0 + \beta_1 x}}{1 + e^{\beta_0 + \beta_1 x}} \end{aligned}$$

## The Logit Model:

$$\begin{aligned}P(y = 1 \mid x) &= F(\beta_0 + \beta_1 x) \\&= \frac{e^{\beta_0 + \beta_1 x}}{1 + e^{\beta_0 + \beta_1 x}}\end{aligned}$$

$$\begin{aligned}P(y = 1 \mid x) &= 1 - F(\beta_0 + \beta_1 x) \\&= 1 - \frac{e^{\beta_0 + \beta_1 x}}{1 + e^{\beta_0 + \beta_1 x}} \\&= \frac{1}{1 + e^{\beta_0 + \beta_1 x}}\end{aligned}$$

## The Logit Model:

Suppose we want to know the probability of  $Y = y_i$ , where  $y_i \in \{0, 1\}$ .

$$P(Y = y_i | x_i) = P(Y = y_i | x)^{y_i} P(Y = 1 - y_i | x)^{1 - y_i}$$

## The Logit Model:

Suppose we want to know the probability of  $Y = y_i$ , where  $y_i \in \{0, 1\}$ .

$$P(Y = y_i | x_i) = P(Y = y_i | x)^{y_i} P(Y = 1 - y_i | x)^{1 - y_i}$$

Recall that in the Bernoulli case this was simply:

$$P(Y = y_i) = p^{y_i} (1 - p)^{1 - y_i}$$

# The Logit Model:

Suppose we want to know the probability of  $Y = y_i$ , where  $y_i \in \{0, 1\}$ .

$$P(Y = y_i | x_i) = P(Y = y_i | x)^{y_i} P(Y = 1 - y_i | x)^{1-y_i}$$

Recall that in the Bernoulli case this was simply:

$$P(Y = y_i) = p^{y_i} (1 - p)^{1-y_i}$$

In logit model:

$$P(Y = y_i | x_i) = \left( \frac{e^{\beta_0 + \beta_1 x_i}}{1 + e^{\beta_0 + \beta_1 x_i}} \right)^{y_i} \left( \frac{1}{1 + e^{\beta_0 + \beta_1 x_i}} \right)^{1-y_i}$$

# The Logit Model:

Suppose we want to know the probability of  $Y = y_i$ , where  $y_i \in \{0, 1\}$ .

$$P(Y = y_i | x_i) = P(Y = y_i | x)^{y_i} P(Y = 1 - y_i | x)^{1-y_i}$$

Recall that in the Bernoulli case this was simply:

$$P(Y = y_i) = p^{y_i} (1 - p)^{1-y_i}$$

In logit model:

$$P(Y = y_i | x_i) = \left( \frac{e^{\beta_0 + \beta_1 x_i}}{1 + e^{\beta_0 + \beta_1 x_i}} \right)^{y_i} \left( \frac{1}{1 + e^{\beta_0 + \beta_1 x_i}} \right)^{1-y_i}$$

The likelihood function is:

$$L(\beta_0, \beta_1 | \mathbf{y}, \mathbf{x}) = \prod_{i=1}^N \left( \frac{e^{\beta_0 + \beta_1 x_i}}{1 + e^{\beta_0 + \beta_1 x_i}} \right)^{y_i} \left( \frac{1}{1 + e^{\beta_0 + \beta_1 x_i}} \right)^{1-y_i}$$

## The likelihood function:

$$L(\beta_0, \beta_1 \mid \mathbf{y}, \mathbf{x}) = \prod_{i=1}^N \left( \frac{e^{\beta_0 + \beta_1 x_i}}{1 + e^{\beta_0 + \beta_1 x_i}} \right)^{y_i} \left( \frac{1}{1 + e^{\beta_0 + \beta_1 x_i}} \right)^{1-y_i}$$

## The likelihood function:

$$L(\beta_0, \beta_1 \mid \mathbf{y}, \mathbf{x}) = \prod_{i=1}^N \left( \frac{e^{\beta_0 + \beta_1 x_i}}{1 + e^{\beta_0 + \beta_1 x_i}} \right)^{y_i} \left( \frac{1}{1 + e^{\beta_0 + \beta_1 x_i}} \right)^{1-y_i}$$

For example, suppose we have 3 observations:

- $y = [1, 0, 1]$
- $x = [5, 2, 1]$



## The likelihood function:

$$L(\beta_0, \beta_1 \mid \mathbf{y}, \mathbf{x}) = \prod_{i=1}^N \left( \frac{e^{\beta_0 + \beta_1 x_i}}{1 + e^{\beta_0 + \beta_1 x_i}} \right)^{y_i} \left( \frac{1}{1 + e^{\beta_0 + \beta_1 x_i}} \right)^{1-y_i}$$

For example, suppose we have 3 observations:

- $y = [1, 0, 1]$
- $x = [5, 2, 1]$

$$L(\beta_0, \beta_1 \mid \mathbf{y}, \mathbf{x}) = \left( \frac{e^{\beta_0 + 5\beta_1}}{1 + e^{\beta_0 + 5\beta_1}} \right) \left( \frac{1}{1 + e^{\beta_0 + 2\beta_1}} \right) \left( \frac{e^{\beta_0 + \beta_1}}{1 + e^{\beta_0 + \beta_1}} \right)$$

## The log likelihood function:

Given:

$$L(\beta_0, \beta_1 \mid \mathbf{y}, \mathbf{x}) = \prod_{i=1}^N \left( \frac{e^{\beta_0 + \beta_1 x_i}}{1 + e^{\beta_0 + \beta_1 x_i}} \right)^{y_i} \left( \frac{1}{1 + e^{\beta_0 + \beta_1 x_i}} \right)^{1-y_i}$$

The log likelihood function is:

$$\begin{aligned} \log(L(\beta_0, \beta_1 \mid \mathbf{y}, \mathbf{x})) &= \sum_{i=1}^N y_i \log \left( \frac{e^{\beta_0 + \beta_1 x_i}}{1 + e^{\beta_0 + \beta_1 x_i}} \right) \\ &\quad + \sum_{i=1}^N (1 - y_i) \log \left( \frac{1}{1 + e^{\beta_0 + \beta_1 x_i}} \right) \end{aligned}$$