

Emory University
MATH 411 & 2 Real Analysis
Learning Notes

Jiuru Lyu

April 29, 2025

Contents

1	The Real Line and Euclidean Space	4
1.1	Algebraic Properties of \mathbb{R} (as a Ordered Field)	4
1.2	Construction of \mathbb{R} and Completeness of \mathbb{R}	7
1.3	Another Approach: Least Upper Bound	11
1.4	Cauchy Sequence and Cauchy Completeness	13
	Summary I: Completeness on Ordered Field	15
1.5	\liminf and \limsup	16
1.6	Euclidean Space \mathbb{R}^n and General Metric Space	18
2	Topology of Euclidean Space	23
2.1	Open Set	23
2.2	Closed Sets	26
	Summary II: Definitions of Point Set Topology	28
2.3	Convergence	29
2.4	Completeness	30
3	Compactness and Connectedness	32
3.1	Compactness	32
3.2	Compactness in \mathbb{R}^n	37
3.3	Nested Set Property	38
3.4	Connectedness	39
4	Continuous Mappings	43
4.1	Continuity	43
4.2	Properties of Continuous Mappings	45
4.3	Uniform Continuity (UC)	48

4.4	Differentiability	50
4.5	Integration	55
5	Uniform Convergence	63
5.1	Definition of Convergence	63
5.2	Integration and Differentiation of Series	67
5.3	The Space of Continuous Functions	68
5.4	The Contraction Mapping Principle (CMP)	71
6	Differential Mappings	75
6.1	Definition and Matrix Representation of a Differential	75
6.2	Necessary and Sufficient Conditions for Differentiability	80
	Summary III: Relations among Properties of $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$	85
6.3	Differentiation Rules	85
6.3.1	Chain Rule	85
6.3.2	Other Differentiation Rules	87
6.4	Geometric Interpretation of Gradient	88
6.5	Mean Value Theorem (MVT)	89
6.6	Taylor's Theorem & Higher Order Differentials	90
6.6.1	One Dimensional Case	90
	Summary IV: Taylor's Formula & Taylor's Approximation	92
6.6.2	Taylor Series	92
6.6.3	Higher Dimensional Case	94
	Summary V: Higher Order Differentials	96
6.7	Minima & Maxima in \mathbb{R}^n	98
7	Inverse and Implicit Function Theorem	105
7.1	Inverse Function Theorem	105
7.2	Implicit Function Thm and Applications	110
7.3	Constrained Extrema	113
7.3.1	Morse Theory: Local Behavior Near a Critical Point	113
7.3.2	Constrained Extremal Problem	114
8	Integration	117
8.1	Definition of Integration	117
8.2	Criterion for Integrability	122
8.2.1	Measure Zero: How to Measure the Size of a Set	122
8.2.2	Lebesgue's Theorem	124
8.3	Improper Integrals	130
8.4	Lebesgue Convergence Theorem	134

9	Computing Integrals	138
9.1	Fubini's Theorem	138
9.2	Change of Variable	141
10	Fourier Analysis	146
10.1	Introduction	146
10.2	Inner Product Space of Functions	146
10.2.1	Basic Concepts	146
10.2.2	The Space \mathcal{C} and L^2	148
10.3	Fourier Analysis on Inner Product Space	149
10.3.1	Geometry of an Inner Product Space	149
10.3.2	Fourier Series and Complete Family	150
10.4	Completeness and Convergence in L^2	153

1 The Real Line and Euclidean Space

1.1 Algebraic Properties of \mathbb{R} (as a Ordered Field)

Axiom 1.1.1 Field Axioms: Recall the following properties

- Addition Axioms

- (1) *commutativity*: $x + y = y + x$
- (2) *associativity*: $(x + y) + z = x + (y + z)$
- (3) *the zero element*: $x + 0 = x$
- (4) *the negative element*: $x + (-x) = 0$

This further gives the definition of *subtraction*: $y - x = y + (-x)$.

- Multiplication Axioms

- (5) *commutativity*: $xy = yx$
- (6) *associativity*: $(xy)z = x(yz)$
- (7) *the one element/unit vector*: $x \cdot 1 = x$
- (8) *inverse*: for each $x \neq 0$, $\exists x^{-1}$ s.t. $x \cdot x^{-1} = 1$

This further gives the definition of *division*: $y/x = y \cdot x^{-1}$ when $x \neq 0$.

- (9) *distribution*: $x(y + z) = xy + xz$

- (10) $1 \neq 0$

- Order Axioms

- (11) *reflexivity*: $x \leq x$
- (12) *anti-symmetry*: If $x \leq y$ and $y \leq x \implies x = y$.
- (13) *transitivity*: If $x \leq y$ and $y \leq z \implies x \leq z$
- (14) *linear relation*: For each pair x, y , either $x \leq y$ or $y \leq x$.
- (15) *compatibility with addition*: If $x \leq y \implies x + z \leq y + z \quad \forall z$
- (16) *compatibility with multiplication*: If $0 \leq x$ and $0 \leq y \implies 0 \leq xy$.

Definition 1.1.2 (Ordered Field). A system (or a set) \mathcal{F} is called an *ordered field* if it satisfies all the above 16 properties.

Remark 1.1 (Examples of Ordered Field) \mathbb{R} and \mathbb{Q} .

Definition 1.1.3 (Field). A set is called a *field* if satisfies all the addition and multiplication axioms.

Definition 1.1.4 (Ring). A set is a *ring* if it satisfies (1) – (9) except (5) and (8).

Example 1.1.5 \mathbb{Z} as a Ring

$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$, the set of integers, is a commutative ring, but not a field.

Remark 1.2 *There is no division operation in a ring as multiplicative inverse is not defined.*

Definition 1.1.6 (Group). A set is a *group* if it satisfies (1) – (4).

Theorem 1.1.7 Law of Trichotomy

If x and y are elements of an ordered field, then exactly one of the relations $x < y$, $x = y$, or $x > y$ holds.

Proposition 1.1.8 Other Algebraic Properties of \mathbb{R} (as an Ordered Field):

1. *unique identities:* If $a + x = a$ for every a , then $x = 0$. If $a \cdot x = a$ for every a , then $x = 1$.
2. *unique inverses:* If $a + x = 0$, then $x = -a$. If $ax = 1$, then $x = a^{-1}$.
3. *no divisors of zero:* If $xy = 0$, then $x = 0$ or $y = 0$.
4. *cancellation laws for addition:* If $a + x = b + x$, then $a = b$. If $a + x \leq b + x$, then $a \leq b$.
5. *cancellation for multiplication:* If $ax = bx$ and $x \neq 0$, then $a = b$. If $ax \geq bx$ and $x > 0$, then $a \geq b$.
6. $0 \cdot x = 0$ for every x .
7. $-(-x) = x$ for every x .
8. $-x = (-1)x$ for every x .
9. If $x \neq 0$, then $x^{-1} \neq 0$ and $(x^{-1})^{-1} = x$.
10. If $x \neq 0$ and $y \neq 0$, then $xy \neq 0$ and $(xy)^{-1} = x^{-1}y^{-1}$.
11. If $x \leq y$ and $0 \leq z$, then $xz \leq yz$. If $x \leq y$ and $z \leq 0$, then $yz \leq xz$.
12. If $x \leq 0$ and $y \leq 0$, then $xy \geq 0$. If $x \leq 0$ and $y \geq 0$, then $xy \leq 0$.
13. $0 < 1$.
14. For any x , $x^2 \geq 0$.

Proof 1. (Of No. 14)

Case I If $x \geq 0$, then $x^2 = x \cdot x \geq 0$, by property (16) of Axiom 1.1.

Case II If $x < 0$, then

$$\begin{aligned} x^2 &= x \cdot x = (-1)(-x) \cdot (-1)(-x) \quad [\text{by property 7 of Proposition 1.7}] \\ &= (-1)^2 \cdot (-x)^2. \end{aligned}$$

Note that $0 = (-1)(-1 + 1) = (-1)^2 + (-1)$ if we distribute (-1) . Then, adding 1 on both sides, we have

$$1 = (-1)^2 + (-1) + 1 = (-1)^2 \quad [\text{by additive inverse}]$$

That is, $(-1)^2 = 1$. So, $x^2 = (-1)^2 \cdot (-x)^2 = 1 \cdot (-x)^2 = (-x)^2 \geq 0$ by Case I.

Q.E.D. ■

Proposition 1.1.9: $ab \leq \frac{a^2 + b^2}{2}$.

Proof 2.

$$(a - b)^2 \geq 0 \quad [\text{By property 14 of Proposition 1.7}]$$

$$a^2 + b^2 - 2ab \geq 0$$

$$2ab \leq a^2 + b^2$$

$$ab \leq \frac{a^2 + b^2}{2}.$$

Q.E.D. ■

Definition 1.1.10 (Absolute Value (Norm) and Distance (Metric)). For $x, y \in \mathbb{R}$, $|x| = \begin{cases} x, & x \geq 0 \\ -x, & x < 0 \end{cases}$

is the *absolute value*, and $d(x, y) = |x - y|$ is the *distance*.

Proposition 1.1.11 Properties of Absolute Value and Distance:

- $|x| \geq 0$ for every x .
- $|x| = 0$ if and only if $x = 0$.
- $|xy| = |x||y|$.
- $d(x, y) \geq 0$
- $d(x, y) = 0$ if and only if $x = y$.
- $d(x, y) = d(y, x)$.

Theorem 1.1.12 Triangle Inequalities

$\forall x, y, z \in \mathbb{R}$

1. $|x + y| \leq |x| + |y|$
2. $||x| - |y|| \leq |x - y|$
3. $d(x, y) \leq d(x, z) + d(z, y)$

Proof 3. (Of No. 1)

Case I Suppose $x \geq 0$ and $y \geq 0$. Then, $x + y \geq 0$, and

$$|x + y| = x + y = |x| + |y|. \quad \square$$

Case II WLOG, suppose $x \geq 0$ and $y < 0$.

- Suppose $x + y \geq 0$, then

$$|x + y| = x + y = |x| - (-y) = |x| - |y| \leq |x| + |y|. \quad \square$$

- Suppose $x + y < 0$, then

$$|x + y| = -(x + y) = -x - y = -|x| + |y| \leq |x| + |y|. \quad \square$$

Case III Suppose $x < 0$ and $y < 0$. Then, $x + y < 0$, and

$$|x + y| = -(x + y) = -x + (-y) = |x| + |y|$$

Q.E.D. ■

1.2 Construction of \mathbb{R} and Completeness of \mathbb{R}

Notation 1.1. Recall the following number systems:

$$\begin{array}{l|l} \mathbb{N} = \mathbb{Z}^+ = \{0, 1, 2, 3, \dots\} & \text{non-negative integers} \\ \mathbb{Z} & \text{integers} \\ \mathbb{Q} = \left\{ \frac{m}{n} : m, n \in \mathbb{Z}, n \neq 0 \right\} & \text{rational numbers} \end{array}$$

Proposition 1.2.2 Important Properties of Number Systems:

- For \mathbb{N} :
 - **Definition 1.2.3 (Principle of mathematical induction).** If S is a subset of \mathbb{Z}^+ s.t. $0 \in S$ and $k \in S \implies k + 1 \in S$, then $S = \mathbb{Z}^+$.
 - **Definition 1.2.4 (Well-Ordered Property).** Each subset $S \neq \emptyset$ has a smallest element.
As a consequence of well-ordering property, we have the principle of complete induction:
 - **Definition 1.2.5 (Principle of Complete Induction).** If $S \subset \mathbb{Z}^+$ is a subset s.t. $\{x \in \mathbb{Z}^+ \mid x < n\} \subset S \implies n \in S$, then $S = \mathbb{Z}^+$.
- For \mathbb{Z} :
 - Commutative ring with identity
- For \mathbb{Q} :
 - **Definition 1.2.6 (Countable).** \mathbb{Q} can be placed in one-to-one correspondence with \mathbb{N} (or a subset of it). The whole \mathbb{Q} can be displayed as a list or sequence.

Remark 1.3 A simple way to prove it is to consider the points in the plane with integer coordinates, say (p, q) . After assigning fraction $\frac{p}{q}$ (simplified to lowest terms and leave out cases when $q = 0$) to this point, we achieve a one-to-one correspondence.

- **Definition 1.2.7 (Dense in Itself).** If $x, y \in \mathbb{Q}$ and $x < y \implies \exists z \in \mathbb{Q} \text{ s.t. } x < z < y$.
- **Proposition 1.2.8 Archimedean Property:**

$$\forall x \in \mathbb{Q}, \exists n \in \mathbb{Z} \text{ s.t. } n > x.$$

Proof 1. If $x \leq 0$, take $n = 1$. If $x = \frac{p}{q}$ with $p, q > 0$, take $n = p + 1$.

Q.E.D. ■

Remark 1.4 *Equivalent formulation of the Archimedean Property:*

- * If $x \in \mathbb{Q}$, then \exists integer n s.t. $x < n$.
- * If $x, y \in \mathbb{Q}$ and $0 < x < y$, then \exists integer k s.t. $kx > y$.
- * If $x > 0 \in \mathbb{Q}$, then \exists integer $n > 0$ s.t. $0 < \frac{1}{n} < x$.

- Ordered field.

\mathbb{Q} is already an ordered field, why do we bother to define \mathbb{R} for analysis?

The big idea: \mathbb{Q} is not *quite complete*

- Evidence 1 (Analysis POV): There is no rational whose square is 2. That is, $x^2 = 2$ has no solution in \mathbb{R} .

Proof 2. We will use proof by contradiction. Assume \exists solution $x = \frac{m}{n}$ with $m, n \in \mathbb{Z}$ and they have no common factors. Then,

$$\left(\frac{m}{n}\right)^2 = 2 \implies m^2 = 2n^2.$$

So, m^2 is even, then m is even as well. Suppose $m = 2k$, $k \in \mathbb{Z}$. Then,

$$\begin{aligned} m^2 &= (2k)^2 = 4k^2 = 2n^2 \\ n^2 &= 2k^2. \end{aligned}$$

So, n^2 is even, and n is even.

* m, n both even, so they have a common factor of 2. This contradict with our assumption.

So, \nexists a solution $x \in \mathbb{Q} \text{ s.t. } x^2 = 2$.

Q.E.D. ■

- Evidence 2 (Geometry POV): There is no rational representation of the diagonal of a square of size 1.

Remark 1.5 (Informal Definition of Sequence Limit) *A sequence is said to converge to a limit x if we can guarantee that the points in the sequence are as close as we wish to x by going far enough out in the sequence.*

Definition 1.2.9 (Limit of a Sequence). A sequence $\{x_n\}$ is said to *converge* to x if $\forall \varepsilon > 0, \exists$ integer N s.t. $|x_n - x| < \varepsilon$ whenever $n \geq N$. (Alternatively, $n \geq N \implies |x_n - x| < \varepsilon$). We denote the *limit* as

$$\lim_{n \rightarrow \infty} x_n = x \quad \text{or, simply} \quad x_n \rightarrow x \text{ as } n \rightarrow \infty.$$

Remark 1.6 N depends on ε , and the smaller the ε , the bigger the N .

Example 1.2.10 Show $\lim_{n \rightarrow \infty} \frac{n+1}{n+2} = 1$.

Proof 3. Given $\varepsilon > 0$ [fix ε], we need to find N s.t. $n \geq N \implies |x_n - 1| < \varepsilon$, where $x_n = \frac{n+1}{n+2}$. Consider

$$|x_n - 1| = \left| \frac{n+1}{n+2} - 1 \right| = \left| \frac{n+1 - n - 2}{n+2} \right| = \left| \frac{-1}{n+2} \right| = \frac{1}{n+2}.$$

Then, we want

$$\frac{1}{n+2} < \varepsilon \iff n+2 > \frac{1}{\varepsilon} \iff n > \frac{1}{\varepsilon} - 2.$$

By the Archimedean property, choose integer $N > \frac{1}{\varepsilon} - 2$. [N is fixed and is what we want to find]
Then, based on the arguments, when $n \geq N$ [n is changing], we have

$$|x_n - 1| = \frac{1}{n+2} \leq \frac{1}{N+2} < \varepsilon.$$

That is,

$$\lim_{n \rightarrow \infty} x_n = 1.$$

Q.E.D. ■

Theorem 1.2.11 Basic Properties of Limits

- **Sandwich Lemma/Squeeze Theorem:** Suppose $x_n \rightarrow L$, $y_n \rightarrow L$, and $x_n \leq z_n \leq y_n$ for all n . Then, $z_n \rightarrow L$. It is also enough to assume that $\exists N_0$ s.t. $n > N_0 \implies x_n \leq z_n \leq y_n$
- If $a \leq x_n \leq b$ for every n and $x_n \rightarrow x$, then $a \leq x \leq b$.
- **Uniqueness:** If x_n is a sequence in an ordered field and $x_n \rightarrow x$ and $x_n \rightarrow y$, then $x = y$.
- **Boundedness:** A convergent sequence is bounded.
- **Arithmetic of Sequence and Limits:** Suppose $x_n \rightarrow x$ and $y_n \rightarrow y$. Then,

$$\{x_n\} + \{y_n\} = \{x_n + y_n\} \implies x_n + y_n \rightarrow x + y$$

$$\lambda \{x_n\} = \{\lambda x_n\} \implies \lambda x_n \rightarrow \lambda x$$

$$\{x_n\} \{y_n\} = \{x_n y_n\} \implies x_n y_n \rightarrow xy$$

$$\{x_n\} / \{y_n\} = \{x_n / y_n\} \implies x_n / y_n \rightarrow x / y$$

Definition 1.2.12 (Monotone Sequence Property/MSP). Every *monotone increasing sequence* that is *bounded (bdd) above* converges.

Remark 1.7 “monotone increasing sequence” refers to a sequence where $x_n \leq x_{n+1} \quad \forall n$; “bdd above” refers to $\exists x$ s.t. $x_n \leq x \quad \forall n$, and we call this x an upper bound.

Definition 1.2.13 (Completeness). An ordered field \mathcal{F} is said to be *complete* if it has the MSP.

Construction of \mathbb{R} (from \mathbb{Q})

Consider set S of sequences,

$$S = \{(x_1, x_2, \dots) \mid x_n \in \mathbb{Q}, x_n \uparrow \text{ (monotone increasing)}, x_n \text{ bdd above}\}.$$

Define *equivalence relation* (reflexive, transitive, symmetric) \sim on S :

$$\{x_n\} \sim \{y_n\} \iff x_n \text{ and } y_n \text{ have the same upper bounds.}$$

Then, each equivalence class defines a unique real number (as the limit of the representing sequence). Let

$$\mathbb{R} = \{x \mid x \text{ is an equivalence class in } S\}.$$

If $r \in \mathbb{Q}$, then r is represented by the sequence r itself $(\{r\})$. So, $\mathbb{Q} \subseteq \mathbb{R}$.

Claim 1.2.14 \mathbb{R} is a complete ordered field under the following operations: For $x = [\{x_n\}]$ and $y = [\{y_n\}]$,

- Addition: $x + y = [\{x_n + y_n\}]$
- Multiplication: $x \cdot y = [\{x_n \cdot y_n\}]$
- Order: $x \leq y \iff \exists \text{ upper bd of } \{x_n\} \text{ that is } \leq \text{all upper bd of } \{y_n\}.$

Theorem 1.2.15

\mathbb{R} is the “unique” complete ordered field.

Remark 1.8 By unique, we mean isomorphism. That is, if \exists another complete ordered field \mathcal{F} , we can put \mathcal{F} and \mathbb{R} into a one-to-one relationship.

Proposition 1.2.16 Properties of \mathbb{R} :

- \mathbb{R} is Archimedean: $\forall x \in \mathbb{R}, \exists \text{ integer } n > x.$
- \mathbb{Q} is dense in \mathbb{R} :
 - If $x, y \in \mathbb{R}$ and $x < y \implies \exists r \in \mathbb{Q} \text{ s.t. } x < r < y.$
 - If $x \in \mathbb{R}$ and $\varepsilon > 0 \implies \exists r \in \mathbb{Q} \text{ s.t. } |x - r| < \varepsilon.$

- The interval $(0, 1)$ is uncountable. (Hence, \mathbb{R} is uncountable).

Proof 4. (of uncountability)

Assume $(0, 1)$ is countable. Then, it can be put into a one-to-one relationship with \mathbb{N} . Say the following list exhausts elements of \mathbb{R} :

$$x_1 = 0.a_{11}a_{12}\cdots a_{1n}\cdots, x_2 = 0.a_{21}a_{22}\cdots a_{2n}\cdots, \dots, x_k = 0.a_{k1}a_{k2}\cdots a_{kn}\cdots, \dots$$

[Goal: find a new number that is not in the list] Define a new number:

$$x = 0.x'_1x'_2\cdots x'_k\cdots,$$

where for each k , $x'_k = \begin{cases} 4 & \text{if } a_{kk} \neq 4 \\ 3 & \text{if } a_{kk} = 4. \end{cases}$ [This construction ensures $x'_k \neq a_{kk}$] Then, $x \in (0, 1)$ and $x \neq x_k \quad \forall k$. * We have constructed a number that is not in the list. So, $(0, 1)$ is not countable.

Q.E.D. ■

1.3 Another Approach: Least Upper Bound

Definition 1.3.1 (Upper Bound/Least Upper Bound). Let $S \subset \mathbb{R}$.

- We say b is an *upper bd* for S if $x \leq b \quad \forall x \in S$.
- We say b is a *least upper bd* for S if b is an upper bd and \leq any upper bd of S .

We use $\text{lub}(S) = \sup(S)$ to denote the least upper bd. (\sup stands for supremum). For sets without an upper bound, we define $\sup(S) = +\infty$.

Remark 1.9 $b = \text{lub}(S) \iff (1) b \text{ is an upper bound, and } (2) b \leq \text{any upper bound of } S$.

Example 1.3.2

Suppose $S_1 = (0, 2)$; $S_2 = [0, 2]$; $S_3 = \emptyset$; $S_4 = (0, \infty)$. Then, $\text{lub}(S_1) = 2$, $\text{lub}(S_2) = 2$, $\text{lub}(S_3) = +\infty$, $\text{lub}(S_4) = +\infty$.

Definition 1.3.3 (Greatest Lower Bound). We use $\text{glb}(S) = \inf(S)$ to denote the greatest lower bound. It is the largest lower bound of S . For sets without a lower bound, we define $\inf(S) = -\infty$.

Example 1.3.4

Example 1.3.2: $\inf(S_1) = 0$, $\inf(S_2) = 0$, $\inf(S_3) = -\infty$, $\inf(S_4) = 0$, $\inf((-\infty, 4)) = -\infty$.

Proposition 1.3.5 : Let $S \subset \mathbb{R}$, $S \neq \emptyset$, then

- $b = \text{lub}(S) \iff b \text{ is an upper bound and } \forall \varepsilon > 0, \exists x \in S \text{ s.t. } x > b - \varepsilon$. This implies that an element slightly smaller than b is not an upper bound any more.

- $a = \inf(S) \iff a$ is a lower bound and $\forall \varepsilon > 0, \exists x \in S$ s.t. $x < a + \varepsilon$.

Proposition 1.3.6 : Suppose $\emptyset \neq A \subset B \subset \mathbb{R}$. Then,

$$\inf(B) \leq \inf(A) \leq \sup(A) \leq \sup(B).$$

Theorem 1.3.7 Equivalent Condition for Completeness: Least Upper Bound Condition

\mathbb{R} has the following properties:

- LUB property: Every non-empty subset bounded above has the least upper bound.
- GLB property: Every non-empty subset bounded below has the greatest lower bound.

Proof 1. (of the LUB Property)

Set-up: Fix any $S \in \mathbb{R}$ that is bounded above and $S \neq \emptyset$.

[WTS: the existence of $\text{lub}(S) \iff$ Tool: MSP (but we need to construct monotone sequence first.)

Step 1 Construction of a Monotone Sequence

Fix an upper bound M for S . For each fixed integer $n \geq 1$, consider $a_k = M - \frac{k}{2^n}, k = 1, 2, \dots$. By the well-ordering property, we can choose an integer k_n who is the 1st integer k s.t. a_k is not an upper bound.

Let $b_n = M - \frac{k_n}{2^n}$. Then, b_n is not an upper bound, but $b_n + \frac{1}{2^n}$ is an upper bound (by construction).

Step 2 Apply MSP to $\{b_n\}$

- b_n is monotone increasing:

Note that

$$b_{n+1} - b_n = \left(M - \frac{k_{n+1}}{2^{n+1}}\right) - \left(M - \frac{k_n}{2^n}\right) = \frac{2k_n - k_{n+1}}{2^{n+1}}$$

Suppose, for the sake of contradiction, that $b_{n+1} - b_n < 0$. Then, $b_{n+1} - b_n \leq -\frac{1}{2^{n+1}}$. That is,

$$b_n \geq b_{n+1} + \frac{1}{2^{n+1}}.$$

* However, by construction, b_n is not an upper bound, but $b_{n+1} + \frac{1}{2^{n+1}}$ is an upper bound. So, there is a contradiction, and thus $b_{n+1} - b_n > 0$. This contradictions shows that b_n is a monotone increasing sequence.

- b_n is bounded above:

Note that $b_n \leq M$. So, b_n is bounded above.

By MSP, suppose $b_n \rightarrow b$ for some $b \in \mathbb{R}$.

Step 3 Show $b = \text{lub}(S)$

- b is an upper bound:

Fix $x \in S$, we have $x \leq b_n + \frac{1}{2^n} \quad \forall n$. When $x \rightarrow \infty$, $x \leq b + 0$. So, $x \leq b$.

- **b is the least upper bound:** [WTS: $\forall \varepsilon > 0, \exists x \in S$ s.t. $b - \varepsilon < x$.]

As b is the limit, we can always find a b_n s.t. $|b_n - b| < \varepsilon$. That is, $b - b_n < \varepsilon$, or $b_n > b - \varepsilon$. Hence, b is the least upper bound.

Q.E.D. ■

1.4 Cauchy Sequence and Cauchy Completeness

Definition 1.4.1 (Cauchy Sequence). A sequence $x_n \in \mathbb{R}$ is a *Cauchy Sequence* if $\forall \varepsilon > 0, \exists N$ s.t. $n, m \geq N \implies |x_n - x_m| < \varepsilon$.

Proposition 1.4.2 : Every convergent sequence is Cauchy.

Proof 1. Suppose $x_n \rightarrow x \in \mathbb{R}$. Given $\varepsilon > 0$. Consider

$$\begin{aligned} |x_n - x_m| &= |x_n - x + x - x_m| \\ &\leq |x_n - x| + |x - x_m| \\ &\leq \frac{\varepsilon}{2} + \frac{\varepsilon}{2}. \end{aligned}$$

Q.E.D. ■

Theorem 1.4.3 Cauchy Completeness

Every Cauchy sequence in \mathbb{R} converges.

Remark 1.10 (Strategy of the Proof) $\text{Cauchy Sequence} \xrightarrow{\text{Lemma 1.4.4}} \text{Bounded Sequence} \xrightarrow{\text{Theorem 1.4.5}} \exists \text{ convergent subsequence} + \text{Cauchy sequence} \xrightarrow{\text{Lemma 1.4.6}} \text{Sequence converges.}$

Lemma 1.4.4 : Every Cauchy sequence is Bounded.

Theorem 1.4.5

Every bounded sequence in \mathbb{R} has a subsequence that converges to some point in \mathbb{R} .

Proof 2. Let $\{x_n\}$ be a bounded sequence in \mathbb{R} . Fix M s.t. $-M < x_n < M \quad \forall n$.

Divide $[-M, M]$ into subintervals $[-M, 0]$ and $[0, M]$. One of them, called I_0 , must contain infinitely many terms of $\{x_n\}$. Choose n_0 s.t. $x_{n_0} \in I_0$.

Divide I_0 into two equal subintervals. One of them, denoted I_1 , contains infinitely many elements. Choose $n_1 > n_0$ s.t. $x_{n_1} \in I_1$.

Continuing this process, we obtain subintervals $I_k = [a_k, b_k]$ for $k = 0, 1, \dots$, and includes n_k with the following properties:

- $I_0 \supset I_1 \supset I_2 \supset \dots$
- $b_k - a_k = \frac{M}{2^k}$

- $x_{n_k} \in I_k$

[To prove $\{x_{n_k}\}$ converges, we prove $\{a_k\}$ and $\{b_k\}$ converge, and apply the Squeeze Theorem.]

- **Show $\{a_k\}$ converges:** a_k is monotone increasing and bounded. By MSP, $a_k \rightarrow a \in \mathbb{R}$.
- **Show $\{b_k\}$ converges:** Note that $b_k = a_k + \frac{M}{2^k}$. When $k \rightarrow \infty$,

$$a_k + \frac{M}{2^k} = a + 0 = a.$$

So, $b_k \rightarrow a$ when $k \rightarrow \infty$.

Hence, as $a_k \leq x_{n_k} \leq b_k$, $a_k \rightarrow a$, $b_k \rightarrow a$, it must be that $x_{n_k} \rightarrow a$ as well.

Q.E.D. ■

Lemma 1.4.6 : If a subsequence of a Cauchy sequence converges to x , then the sequence itself converges to x .

Proof 3. Given $\{x_n\}$ is Cauchy and $x_{n_k} \rightarrow x$, [WTS: $x_n \rightarrow x$]. Consider

$$\begin{aligned} |x_n - x| &= |x_n - x_{n_k} + x_{n_k} - x| \\ &\leq \underbrace{|x_n - x_{n_k}|}_{\text{Cauchy} \Rightarrow \text{small}} + \underbrace{|x_{n_k} - x|}_{\text{Convergent} \Rightarrow \text{small}} \end{aligned}$$

Q.E.D. ■

Summary I: Completeness on Ordered Field

Let \mathcal{F} be an ordered field.

Definitions

- **Archimedean Property:** $\forall x \in \mathcal{F}, \exists \text{ integer } N \text{ s.t. } x < N$.
(Equivalently, $\forall \varepsilon > 0, \exists \text{ integer } n \text{ s.t. } 0 < \frac{1}{n} < \varepsilon$).
- **Monotone Sequence Property (MSP):** Every monotone increasing sequence bounded above converges.
- **Completeness:** We say \mathcal{F} is complete if it has the MSP.
- **LUB Property:** Every set $S \neq \emptyset$ bounded above has a least upper bound.
- **Cauchy Property:** Every Cauchy sequence converges.

Facts in any ordered field

- $\text{MSP} \implies \text{Archimedean Property}$

Remark 1.11 *In general, the converse is not true. For example, \mathbb{Q} has the Archimedean property but not MSP.*

- $\text{MSP} \iff \text{LUB Property}$.
- $\text{MSP} \implies \text{Cauchy Property}$

Remark 1.12 *The converse is true when Archimedean property is true.*

Facts in \mathbb{R}

- $\text{MSP} \iff \text{LUB Property} \iff \text{Cauchy Property}$

1.5 \liminf and \limsup **Example 1.5.1 Cluster Points of a Sequence**

Consider the sequence

$$a_n = (-1)^n + \frac{1}{n}.$$

Then, $a_1 = 0$, $a_2 = 1 + \frac{1}{2}$, $a_3 = -1 + \frac{1}{3}$, $a_4 = 1 + \frac{1}{4}$, \dots . This sequence does not converge. However, its terms seem to “cluster” around 1 and -1 .

Definition 1.5.2 (Cluster Points). A point x is called a *cluster point* of a sequence $\{x_n\}$ if $\forall \varepsilon > 0$, \exists infinitely many values of n s.t. $|x_n - x| < \varepsilon$.

Remark 1.13 *This definition is weaker than that of limits.*

Proposition 1.5.3 Relation Between Limits and Cluster Points: Suppose $x_n \in \mathbb{R}$ and $x \in \mathbb{R}$. Then,

1. x is a cluster point of $\{x_n\} \iff \forall \varepsilon > 0$ and \forall integer N , $\exists n > N$ s.t. $|x_n - x| < \varepsilon$.
2. x is a cluster point of $\{x_n\} \iff \exists$ subsequence $x_{n_k} \rightarrow x$.
3. $x_n \rightarrow x \iff$ every subsequence converges to x .
4. $x_n \rightarrow x \iff$ the sequence is bounded and x is the only cluster point.
5. $x_n \rightarrow x \iff$ every subsequence has a further sequence that converges to x .

Proof 1. (of some claims)

1. Follows from Definition.
2. (\Rightarrow) Assume x is a cluster point. [WTS: \exists subsequence $x_{n_k} \rightarrow x$].

Given $\varepsilon_1 = 1$ and $N = 1$, by (1), $\exists n_1 > 1$ s.t. $|x_{n_1} - x| < \varepsilon = 1$.

Given $\varepsilon_2 = \frac{1}{2}$ and $N = n_1$, by (1), $\exists n_2 > n_1$ s.t. $|x_{n_2} - x| < \varepsilon = \frac{1}{2}$.

So, in general, given $\varepsilon_k = \frac{1}{k}$ and $N = n_{k-1}$,

$$\exists n_k > n_{k-1} = N_k \text{ s.t. } |x_{n_k} - x| < \varepsilon_k = \frac{1}{k}.$$

Then, $x_{n_k} \rightarrow x$ as $k \rightarrow \infty$.

3. (\Leftarrow) [Prove by contrapositive/contradiction] Assume every subsequence converges. For the sake of contradiction, assume x_n does not converge to x . Then we need to construct a subsequence x_{n_k} s.t. $x_{n_k} \not\rightarrow x$.
4. (\Leftarrow) [Prove by contrapositive/contradiction]

5. (\Leftarrow) Use (4). Every subsequence has its own subsequence that converges to x . So, x is a cluster point of every subsequence. Then, we just need to show x is the only cluster point of $\{x_n\}$.

Q.E.D. ■

Definition 1.5.4 (\liminf **and** \limsup). Given a sequence $x_n \in \mathbb{R}$. For each integer $k \geq 1$, let

$$a_k = \inf \underbrace{\{x_{k+1}, x_{k+2}, \dots\}}_{\text{Set } S_k} \quad \text{and} \quad b_k = \sup \{x_{k+1}, x_{k+2}, \dots\} = \sup S_k.$$

Then,

$$\liminf x_n = \sup \{a_k\} \quad \text{and} \quad \limsup x_n = \inf \{b_k\}.$$

Remark 1.14

- $a_k \leq b_k$, a_k is monotone increasing sequence, and b_k is monotone decreasing sequence. Thus,

$$\liminf x_n = \lim_{k \rightarrow \infty} a_k \quad \text{and} \quad \limsup x_n = \lim_{k \rightarrow \infty} b_k.$$

Also, $\liminf x_n \leq \limsup x_n$.

- $\limsup x_n = +\infty \iff b_k = +\infty \quad \forall k \iff x_n$ is not bounded above.

$\liminf x_n = -\infty \iff a_k = -\infty \quad \forall k \iff x_n$ is not bounded below.

Proposition 1.5.5 : $\limsup x_n = b \in \mathbb{R} \iff \forall \varepsilon > 0$,

1. $\exists N$ s.t. $n \geq N \implies x_n < b + \varepsilon$, and
2. $\forall M, \exists n \geq M$ s.t. $x_n > b - \varepsilon$.

Proof2. (of forward direction) By definition, we know $\lim_{k \rightarrow \infty} b_k = b$, which implies $\forall \varepsilon > 0, \exists N$ s.t. $k \geq N \implies |b_k - b| < \varepsilon$. That is, $-\varepsilon < b_k - b < \varepsilon$. As b_k is monotone decreasing, $b_k - b \geq 0$. So, $\boxed{0 \leq b_k - b < \varepsilon}$.

1. Note that $b_k = \sup \{x_{k+1}, x_{k+2}, \dots\}$. So, if $n > k$, $x_n \leq b_k < b + \varepsilon \quad \forall k \geq N$. Therefore,

$$n \geq N + 1 \implies x_n < b + \varepsilon.$$

2. We have $0 \leq b_k - b$, or $b_k \geq b \quad \forall k$. Given any integer M . [We need to find $n \geq M$ s.t. $x_n > b - \varepsilon$]
Then,

$$b_M = \sup \{x_{M+1}, x_{M+2}, \dots\} \geq b.$$

So, by definition of supremum, we can find $n > M$ s.t. $x_n > b_M - \varepsilon \geq b - \varepsilon$.

Q.E.D. ■

Proposition 1.5.6 : $\limsup x_n = b \in \mathbb{R} \implies \exists$ subsequence $x_{n_k} \rightarrow b$.

Proof3. We will construct a subsequence n_k inductively such that

$$b - \varepsilon_k < x_{n_k} < b + \varepsilon_k, \quad \varepsilon_k = \frac{1}{k}.$$

Given $\varepsilon = 1$, by Proposition 1.5.5(1), $\exists N_1$ s.t. $n \geq N_1 \implies x_n < b + \varepsilon_1$. Further, by Proposition 1.5.5(2), for $M = N_1$, $\exists n_1 > N_1$ s.t. $x_{n_1} > b - \varepsilon_1$. Therefore,

$$b - \varepsilon_1 < x_{n_1} < b + \varepsilon_1.$$

Claim Given k_n , we can find n_{k+1} s.t. $n_{k+1} > n_k$, and

$$b - \frac{1}{k+1} < x_{n_{k+1}} < b + \frac{1}{k+1}.$$

After $\{x_{n_k}\}$ is constructed, use the sandwich lemma to prove $x_{n_k} \rightarrow b$.

Q.E.D. ■

Remark 1.15 Similar arguments hold for $\liminf x_n = a$.

Proposition 1.5.7 Relation Between Cluster Points and Limit: Let $x_n \in \mathbb{R}$ be a given sequence.

1. If x is a cluster point $\implies \liminf x_n \leq x \leq \limsup x_n$.
2. If $a = \liminf x_n$ is finite $\implies a$ is the smallest cluster point.
3. If $b = \limsup x_n$ is finite $\implies b$ is the largest cluster point.
4. $x_n \rightarrow x \in \mathbb{R} \iff \liminf x_n = \limsup x_n = x$.

Proof4. (of (1)) Suppose x is a cluster point. Then, \exists subsequence $x_{n_k} \rightarrow x$ as $k \rightarrow \infty$.

[WTS: $a_n \leq x \leq b_n \quad \forall n$]

For each n , $b_n = \sup \{x_{n+1}, x_{n+2}, \dots\} \geq x_{n_k}$ for large enough k . Let $k \rightarrow \infty$, we have $b_n \geq x$. Similarly, $a_n = \inf \{x_{n+1}, x_{n+2}, \dots\} \leq x_{n_k}$ for large enough k . As $k \rightarrow \infty$, $a_n \leq x$.

So, $a_n \leq x \leq b_n$. Take the limit as $n \rightarrow \infty$:

$$\lim_{n \rightarrow \infty} a_n \leq x \leq \lim_{n \rightarrow \infty} b_n \implies \liminf x_n \leq x \leq \limsup x_n.$$

Q.E.D. ■

1.6 Euclidean Space \mathbb{R}^n and General Metric Space

Notation 1.1. $\mathbb{R}^n = \{(x_1, x_2, \dots, x_n) \mid x_1, \dots, x_n \in \mathbb{R}\}$.

Remark 1.16 (\mathbb{R}^n is a Vector Space) We can write its standard bases as $\{e_1, e_2, \dots, e_n\}$, and the general representation of x will be

$$x = \sum_{j=1}^n x_j e_j.$$

Definition 1.6.2 (Norm and Metric). For $x, y \in \mathbb{R}^n$, define *norm* (or length) as

$$\|x\| = \sqrt{\sum_{i=1}^n x_i^2}$$

and the *metric* (distance) as

$$d(x, y) = \|x - y\| = \sqrt{\sum_{i=1}^n (x_i - y_i)^2}.$$

Definition 1.6.3 (Inner Product). We define the *inner product* (or dot product) as

$$\langle x, y \rangle = \sum_{i=1}^n x_i y_i.$$

Geometrically, if θ is the angle between x and y , then

$$\langle x, y \rangle = \|x\| \cdot \|y\| \cdot \cos \theta.$$

So, if $x \perp y$, $\langle x, y \rangle = 0$.

Proposition 1.6.4 Properties of Inner Product: Suppose $\langle \cdot, \cdot \rangle$ is an inner product, then

- Positive definite: $\langle x, x \rangle \geq 0$ and $\langle x, x \rangle = 0 \iff x = 0$.
- Linearity: $\langle x, y + z \rangle = \langle x, y \rangle + \langle x, z \rangle$ and $\langle \alpha x, y \rangle = \alpha \langle x, y \rangle$.
- Symmetry: $\langle x, y \rangle = \langle y, x \rangle$.

Proposition 1.6.5 Properties of Norm: Suppose $\|\cdot\|$ is a norm, then

- Positive definite: $\|x\| \geq 0$ and $\|x\| = 0 \iff x = 0$.
- Linearity: $\|\alpha x\| = |\alpha| \cdot \|x\|$.
- Triangle Inequality: $\|x + y\| \leq \|x\| + \|y\|$.

Proposition 1.6.6 Properties of Metric: Suppose $d(\cdot, \cdot)$ is a metric, then

- Positive definite: $d(x, y) \geq 0$ and $d(x, y) = 0 \iff x = y$.
- Symmetry: $d(x, y) = d(y, x)$.
- Triangle Inequality: $d(x, y) \leq d(x, z) + d(z, y)$.

Remark 1.17 Inner product always induces a norm. Norm always induced a metric.

Theorem 1.6.7 Cauchy-Schwarz Inequality

$$|\langle x, y \rangle| \leq \|x\| \cdot \|y\|.$$

Example 1.6.8 Use Cauchy-Schwarz Inequality to Prove Triangle Inequality of Norms**Proof 1.** Note that

$$\begin{aligned}
\|x + y\|^2 &= \langle x + y, x + y \rangle && \text{[Definition]} \\
&= \langle x + y, x \rangle + \langle x + y, y \rangle && \text{[Distribution]} \\
&= \langle x, x \rangle + \langle y, x \rangle + \langle x, y \rangle + \langle y, y \rangle && \text{[Distribution]} \\
&= \|x\|^2 + \|y\|^2 + 2\langle x, y \rangle && \text{[Symmetry]} \\
&\leq \|x\|^2 + \|y\|^2 + 2 \cdot \|x\| \cdot \|y\| && \text{[Cauchy-Schwarz]} \\
&= (\|x\| + \|y\|)^2.
\end{aligned}$$

Q.E.D. ■

Definition 1.6.9 (General Metric Space). A *metric space* (M, d) is a set M and a function $d : M \times M \rightarrow \mathbb{R}$ s.t. $\forall x, y, z \in M$, the following conditions hold:

- Positive definite: $d(x, y) \geq 0$ and $d(x, y) = 0 \iff x = y$.
- Symmetry: $d(x, y) = d(y, x)$.
- Triangle Inequality: $d(x, y) \leq d(x, z) + d(z, y)$.

Definition 1.6.10 (General Normed Space). A *normed space* $(V, \|\cdot\|)$ is a vector space V together with a function $\|\cdot\| : V \rightarrow \mathbb{R}$ s.t. $\forall x, y \in V$ and $\forall \alpha \in \mathbb{R}$,

- Positive definite: $\|x\| \geq 0$ and $\|x\| = 0 \iff x = 0$.
- Linearity: $\|\alpha x\| = |\alpha| \cdot \|x\|$
- Triangle Inequality: $\|x + y\| \leq \|x\| + \|y\|$

Definition 1.6.11 (General Inner Product Space). An *inner product space* $(V, \langle \cdot, \cdot \rangle)$ is a vector space V and a function $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{R}$ s.t. $\forall x, y, z \in V$ and $\forall \alpha \in \mathbb{R}$:

- Positive definite: $\langle x, x \rangle \geq 0$ and $\langle x, x \rangle = 0 \iff x = 0$.
- Symmetry: $\langle x, y \rangle = \langle y, x \rangle$.
- Linearity: $\langle \alpha x, y \rangle = \alpha \langle x, y \rangle$ and $\langle x + y, z \rangle = \langle x, z \rangle + \langle y, z \rangle$.

Example 1.6.12

- \mathbb{R}^n is a metric space with $d(x, y) = \|x - y\|$.
- *Discrete Metric*: Given any set M , define

$$d(x, y) = \begin{cases} 0, & x = y \\ 1, & x \neq y. \end{cases}$$

- *Bounded Metric*: Given metric space (M, d) , define $\rho : M \times M \rightarrow \mathbb{R}$:

$$\rho(x, y) = \frac{d(x, y)}{1 + d(x, y)}.$$

Claim 1.6.13 (M, ρ) is also a metric space.

- \mathbb{R}^2 is a metric space under the taxicab metric $d_1 : \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow \mathbb{R}$:

$$d_1((x_1, y_1), (x_2, y_2)) = |x_2 - x_1| + |y_2 - y_1|.$$

- Let $\mathcal{C}([0, 1])$ be the collection of all continuous function $f : [0, 1] \rightarrow \mathbb{R}$. Define

$$\langle f, g \rangle = \int_0^1 f(x)g(x) \, dx.$$

Then, \mathcal{C} is an inner product space.

Remark 1.18 (Relation Among Inner Product, Normed, and Metric Space)

$$\text{Inner Product} \implies \text{Norm} \implies \text{Metric}$$

- An inner product $\langle \cdot, \cdot \rangle$ induces a norm:

$$\|x\| = \sqrt{\langle x, x \rangle}.$$

- A norm $\|\cdot\|$ always induces a metric:

$$d(x, y) = \|x - y\|.$$

Theorem 1.6.14 General Cauchy-Schwarz Inequality

In an inner product space $(V, \langle \cdot, \cdot \rangle)$, we have $\forall v, w \in V$,

$$|\langle v, w \rangle| \leq \langle v, v \rangle^{\frac{1}{2}} \cdot \langle w, w \rangle^{\frac{1}{2}}.$$

Proof2. If $v = 0$ or $w = 0$, it is trivial.

Assume $v \neq 0$ and $w \neq 0$. For any $t \in \mathbb{R}$, consider

$$\langle tv + w, tv + w \rangle$$

Then,

$$0 \leq \langle tv + w, tv + w \rangle = t^2 \underbrace{\langle v, v \rangle}_a + 2t \underbrace{\langle v, w \rangle}_b + \underbrace{\langle w, w \rangle}_c$$

Let $f(t) = at^2 + 2bt + c$ be a 2nd order polynomial of t . Note that $f(t) \geq 0 \quad \forall t \in \mathbb{R}$. On the other hand (OTOH), since $a = \langle v, v \rangle > 0$, $f(t)$ has minimum where $f'(t) = 0$.

$$f'(t) = 2at + 2b = 0$$

$$t = -\frac{b}{a}.$$

So, $f\left(-\frac{b}{a}\right) \geq 0$, or

$$\left(-\frac{b}{a}\right)^2 a + 2b\left(-\frac{b}{a}\right) + c \geq 0$$

$$\frac{b^2}{a} - 2\frac{b^2}{a} + c \geq 0$$

$$c \geq \frac{b^2}{a}$$

$$b^2 \leq ac$$

$$(\langle v, w \rangle)^2 \leq \langle v, v \rangle \cdot \langle w, w \rangle$$

$$|\langle v, w \rangle| \leq \langle v, v \rangle^{\frac{1}{2}} \cdot \langle w, w \rangle^{\frac{1}{2}}.$$

Q.E.D. ■

2 Topology of Euclidean Space

2.1 Open Set

Definition 2.1.1 (Neighborhood & Open Set). Let (M, d) be a metric space. Fix $x \in M$ and $\varepsilon > 0$.

- *Neighborhood (nbdd):*

$$D(x, \varepsilon) = \{y \in M \mid d(x, y) < \varepsilon\}.$$

It is also referred as ε -nbdd, ε -disk, or ε -ball.

- *Open Set:* A set $A \subset M$ is *open* if $\forall x \in A, \exists \varepsilon > 0$ s.t. $D(x, \varepsilon) \subset A$.

Example 2.1.2 Open Set

- The unit disk $D = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 < 1\}$ is open in \mathbb{R}^2 .
- The interval $(0, 1) \subset \mathbb{R}^1$ is open.
- Given any metric space (M, d) and $x_0 \in M$. The disk

$$D(x_0, r) = \{x \in M \mid d(x, x_0) < r\}$$

is open $\forall r > 0$.

Proof 1. Fix $x \in D(x_0, r)$. [WTS: $\exists \varepsilon > 0$ s.t. $D(x, \varepsilon) \subset D(x_0, r)$.]

Since $x \in D(x_0, r)$, by definition, $d(x, x_0) < r$. Hence, $\varepsilon = r - d(x, x_0) > 0$.

Claim 2.1.3 $D(x, \varepsilon) \subset D(x_0, r)$.

Proof. Let $y \in D(x, \varepsilon)$. Then,

$$\begin{aligned} d(y, x_0) &\leq d(y, x) + d(x, x_0) \\ &< \varepsilon + d(x, x_0) \\ &= r - \cancel{d(x_0, x)} + \cancel{d(x_0, x)} \\ &= r. \end{aligned}$$

So, $d(y, x_0) < r$. By definition, $y \in D(x_0, r)$. \square

So, $D(x, \varepsilon) \subset D(x_0, r)$. By definition, $D(x_0, r)$ is open.

Q.E.D. \blacksquare

- The set $S = \{(x, y) \in \mathbb{R}^2 \mid xy > 1\}$ is open.

Proof 2. Given $(x, y) \in S$. [WTS: $\exists \varepsilon > 0$ s.t. $D((x, y), \varepsilon) \subset S$.]

Since $xy > 1$, $\lambda = \frac{1}{2} \left(1 - \frac{1}{xy} \right) > 0$.

WLOG, assume $x > 0$ and $y > 0$.

Let $\varepsilon = \min \{\lambda x, \lambda y\}$. Then, for $(u, v) \in D((x, y), \varepsilon)$, we have

$$\begin{aligned} d((u, v), (x, y)) &< \varepsilon \\ \sqrt{(x - u)^2 + (y - v)^2} &< \varepsilon. \end{aligned}$$

So, $|x - u| < \varepsilon$ and $|y - v| < \varepsilon$. Then,

$$\begin{aligned} x \left| 1 - \frac{u}{x} \right| &< \varepsilon \\ \frac{u}{x} &> 1 - \frac{\varepsilon}{x} \geq 1 - \frac{\lambda x}{x} = 1 - \lambda. \end{aligned}$$

Similarly,

$$\frac{v}{y} > 1 - \lambda.$$

Then,

$$\begin{aligned} u \cdot v &= \frac{u}{x} \cdot \frac{v}{y} \cdot (xy) > (1 - \lambda)^2 (xy) \\ &> (1 - 2\lambda)(xy) = 1. \end{aligned}$$

So, as $uv > 1$, $(u, v) \in S$. Hence, S is open.

Sketch. Given $xy > 0$; Want $uv > 1$. Note that

$$\begin{aligned} uv &= \underbrace{\frac{u}{x}}_{(1-\lambda)} \cdot \underbrace{\frac{v}{y}}_{(1-\lambda)} \cdot xy \\ &= (1 - \lambda)^2 (xy) \\ &> (1 - 2\lambda + \lambda^2)(xy) \\ &> (1 - 2\lambda)(xy) \\ &\geq 1 \\ \implies 1 - 2\lambda &\geq \frac{1}{xy}. \end{aligned}$$

Q.E.D. ■

Remark 2.1

- In the above definition, ε depends on the point x .
- The open set is defined w.r.t. the underline metric space.

Example 2.1.4

$A = (0, 1)$. Then, A is an open set as a subset in \mathbb{R}^1 . However, A is not open as a subset in \mathbb{R}^2 .

Proposition 2.1.5 Properties of Open Set: Let (M, d) be a metric space. Then,

- The intersection of a finite number of open sets is open.
- The union of any number of open sets is open.
- \emptyset and M are open.

Proof 3. (of ①) Suppose $A = \bigcap_{j=1}^n A_j$. Fix $x \in A$. By definition, $x \in A_j \quad \forall j = 1, \dots, n$. Then, we can find $\varepsilon_j > 0$ s.t. $D(x, \varepsilon_j) \in A_j$. As A_j is open. Take $\varepsilon = \min \{\varepsilon_1, \varepsilon_2, \dots, \varepsilon_n\}$. We know

$$D(x, \varepsilon) \in A_j \quad \forall j = 1, \dots, n.$$

Hence, $D(x, \varepsilon) \in \bigcap_{j=1}^n A_j$. So, A is open.

Q.E.D. ■

Remark 2.2 The intersection of infinitely many number of open sets may not be open.

Definition 2.1.6 (Interior Point). Let $A \subset M$. A point $x \in A$ is called an *interior point* of A if $\exists \varepsilon > 0$ s.t. $D(x, \varepsilon) \subset A$. The *interior* of A is the collection of all interior points, denoted by $\text{int}(A)$.

Example 2.1.7

- $A = \{x_0\} \subset \mathbb{R}^n$, $\text{int}(A) = \emptyset$ as there is no nbdd around the point x_0 .
- $A = (0, 1) \subset \mathbb{R}^1$, $\text{int}(A) = A$.

Remark 2.3 A set is open if every point in A is an interior point of A .

- $B = [0, 1] \subset \mathbb{R}^1$, $\text{int}(B) = (0, 1)$.

Proposition 2.1.8 Properties of $\text{int}(A)$:

- $\text{int}(A)$ is open.
- $\text{int}(A)$ is the union of all open subsets of A .

Remark 2.4 Or, $\text{int}(A)$ is the largest open subset of A .

- A is open $\iff A = \text{int}(A)$.

2.2 Closed Sets

Definition 2.2.1 (Closed Set). A set $A \subset M$ is *closed* if its complement, $A^C = M \setminus A$, is open.

Example 2.2.2

- $A = [0, 1] \subset \mathbb{R}^1$.
 $A^C = (-\infty, 0) \cup (1, +\infty)$.
 A^C is open $\implies A$ is closed.
- $B = \{(x, y) \in \mathbb{R}^2 \mid 1 < x^2 + y^2 \leq 4\}$.
 $B^C = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 \leq 1 \text{ or } x^2 + y^2 > 4\}$.
 B is not open and not closed.
- A single point set is closed.
- $B(x, \varepsilon) = \{y \in M \mid d(y, x) \leq \varepsilon\}$ is closed.

Proposition 2.2.3 Basic Properties of Closed Sets: Given (M, d) , then

- Union of finite number of closed set is closed.
- Intersection of any number of closed set is closed.
- \emptyset and M are always closed.

Remark 2.5 In property ①, one cannot replace “finite number” by “countably many.”

Definition 2.2.4 (Accumulation Point). A point $x \in M$ is an *accumulation point* of the set A if $\forall \varepsilon > 0, \exists y \in A$ s.t. $y \neq x$ and $y \in D(x, \varepsilon)$. The collection of accumulation points of A is denoted as $\text{ac}(A)$.

Remark 2.6 x does not need to be in A .

Definition 2.2.5 (Closure/ $\text{cl}(A)$).

$$\begin{aligned} \text{cl}(A) &= \text{intersection of all closed sets containing } A \\ &= A \cup \text{ac}(A). \end{aligned}$$

Definition 2.2.6 (Boundary of $A/\partial A/\text{bd}(A)$).

$$\begin{aligned} \text{bd}(A) &= \partial A = \text{cl}(A) \cap \text{cl}(M \setminus A) \\ &= \text{cl}(A) \setminus \text{int}(A). \end{aligned}$$

Theorem 2.2.7 Equivalent Conditions of Closed Sets

Let $A \subset M$, the following are equivalent (TFAE):

- A is closed.
- $\text{ac}(A) \subset A$.
- $A = \text{cl}(A)$.
- $\text{bd}(A) \subset A$.

Proof 1.

(① \implies ②): Let $A \subset M$ be closed and $x \in \text{ac}(A)$. [WTS: $x \in A$.] Assume $x \notin A$. Then, $x \in M \setminus A$.

[Proof by contradiction.]

Since A is closed, $M \setminus A$ is open, which means $x \in \text{int}(M \setminus A)$. That is, $\exists \varepsilon > 0$ s.t. $D(x, \varepsilon) \subset M \setminus A$. Hence, $D(x, \varepsilon) \cap A = \emptyset$. * This contradicts with the assumption that $x \in \text{ac}(A)$. As $D(x, \varepsilon) \cap A = \emptyset$, $\nexists y \in A$ s.t. $y \in D(x, \varepsilon)$. Hence, $x \in A$. \square

(② \iff ③): We have $\text{cl}(A) = A \cup \text{ac}(A)$.

(\implies): If ② is true, $\text{ac}(A) \subset A$. Then, $\text{cl}(A) = A$.

(\impliedby): If ③ is true $\text{cl}(A) = A$. Then, $A \cup \text{ac}(A) = A$, so $\text{ac}(A) \subset A$. \square

(③ \implies ④): Note that $\text{bd}(A) = \text{cl}(A) \cap \text{cl}(M \setminus A)$. Then, $\text{bd}(A) \subset \text{cl}(A)$. If $A = \text{cl}(A)$, then $\text{bd}(A) \subset \text{cl}(A) = A$. \square

(④ \implies ①): Suppose $\text{bd}(A) \subset A$. Assume A is not closed, then $M \setminus A$ is not open. [Proof by contradiction.] So, $\exists x_0 \in M \setminus A$ that is not an interior point. Hence, $\forall \varepsilon > 0$, $D(x_0, \varepsilon) \not\subset M \setminus A$. So, $D(x_0, \varepsilon) \cap A \neq \emptyset$. Hence, $\exists y \in D(x_0, \varepsilon) \cap A$. Note that $x_0 \in M \setminus A$ but $y \in D(x_0, \varepsilon) \cap A$. So, $x_0 \neq y$. By definition, $x_0 \in \text{ac}(A)$. * As $x_0 \in \text{ac}(A) \subset \text{bd}(A)$, but $x_0 \notin A$, this contradicts with the assumption that $\text{bd}(A) \subset A$. Hence, A must be closed.

Q.E.D. \blacksquare

Proposition 2.2.8 :

- $\text{cl}(A) \cap A = A$.
- If A is open, then $\text{bd}(A) \subset M \setminus A$.

Definition 2.2.9 (Limit Point of a Set). A point $x \in M$ is called a limit point of A if $U \cap A \neq \{x\}$ for every open set U containing x .

Proposition 2.2.10 :

- If $x \in \text{ac}(A)$, then x is a limit point.
- If x is a limit point of A and $x \notin A$, then $x \in \text{ac}(A)$.
- If x is a limit point of A , \exists a sequence $x_n \in A$ with $x_n \rightarrow x$.
- A is closed $\iff A$ contains all of its limit points.

Summary II: Definitions on Point Set Topology

Let M be a metric space and $A \subset M$.

- $x \in A$ is an *interior point* of A if $\exists \varepsilon > 0$ with $D(x, \varepsilon) \subset A$.
- A is said to be *open* if every point of A is an interior point, or equivalently, $\text{int}(A) = A$.
- A *neighborhood* of a point x is any open set U containing x .
- A is *closed* if its complement $M \setminus A$ is open.
- A point $x \in M$ is an *accumulation point* of A is $\forall \varepsilon > 0, \exists y \in A$ with $y \neq x$ and $y \in D(x, \varepsilon)$.
- *Closure* of A : $\text{cl}(A) = A \cup \text{ac}(A)$.
- *Boundary* of A : $\partial A = \text{bd}(A) = \text{cl}(A) \cap \text{cl}(A \setminus M) = \text{cl}(A) \setminus \text{int}(A)$.

2.3 Convergence

Definition 2.3.1 (Convergence of a Sequence). Let (M, d) be a metric space. Let $x_k \in M$ be a sequence and $x \in M$. We say that x_k *converges* to x (write $x_k \rightarrow x$) if $\forall \varepsilon > 0, \exists N$ s.t. $d(x_k, x) < \varepsilon \quad \forall k \geq N$.

Theorem 2.3.2 Equivalent Definitions of Convergence

- $x_k \rightarrow x \iff \forall$ open set U containing $x, \exists N$ s.t. $x_k \in U \quad \forall k \geq N$.

Remark 2.7 This definition replaces ε -neighborhood by an arbitrary neighborhood.

- $x_k \rightarrow x \iff d(x_k, x) \rightarrow 0$.

Theorem 2.3.3 Equivalent Definition of Convergence in \mathbb{R}^n

In \mathbb{R}^n , write

$$v_k = (v_k^{(1)}, v_k^{(2)}, \dots, v_k^{(n)}) \quad \text{and} \quad v = (v^{(1)}, v^{(2)}, \dots, v^{(n)}).$$

Then,

$$d(v_k, v)^2 = \|v_k - v\|^2 = \sum_{i=1}^n |v_k^{(i)} - v^{(i)}|^2.$$

Thus, $v_k \rightarrow v \iff v_k^{(i)} \rightarrow v^{(i)} \quad \forall i = 1, \dots, n$

Proposition 2.3.4 : Let $v_k, w_k \in \mathbb{R}^n$ and $\lambda_k, \lambda \in \mathbb{R}$ with $v_k \rightarrow v, w_k \rightarrow w, \lambda_k \rightarrow \lambda$. Then,

- $v_k + w_k \rightarrow v + w$
- $\lambda v_k \rightarrow \lambda v$
- $\lambda_k v_k \rightarrow \lambda v$

Theorem 2.3.5 Convergence and Closedness

Let (M, d) be a metric space and $A \subset M$.

- A is closed \iff for every sequence $x_k \in A$ that converges in M , the limit lies in A .
- $x \in \text{cl}(A) \iff \exists x_k \in A$ s.t. $x_k \rightarrow x$.

Proof1. (of ①, sketch):

(\Rightarrow) Assume $A \subset M$ is closed. Let $x_k \in A$ be a sequence with $x_k \rightarrow x \in M$. [WTS: $x \in A$.] Suppose $x \notin A$. Then, $x \in M \setminus A$. A is closed $\implies M \setminus A$ is open $\implies \exists \varepsilon > 0$ with $D(x, \varepsilon) \subset M \setminus A$. As $x_k \rightarrow x$, some $x_k \in D(x, \varepsilon) \subset M \setminus A$. ✖ This contradicts with our assumption that $x_k \in A$. So, $x \in A$. \square

(\Leftarrow): Suppose $x_k \in A$ with $x_k \rightarrow x \in A$. Assume $A \subset M$ is not closed. Then, $M \setminus A$ is not open $\implies \exists x \in M \setminus A$ s.t. $\forall \varepsilon > 0, D(x, \varepsilon) \not\subset M \setminus A$. For $\varepsilon = \frac{1}{k}, \exists x_k \in D(x, \frac{1}{k}) \cap A$. Then, ✖ $x_k \rightarrow x \notin A$, contradicting with the assumption $x_k \rightarrow x \in A$. Hence, A must be closed.

Q.E.D. \blacksquare

2.4 Completeness

Definition 2.4.1 (Cauchy Sequence). $\{x_k\} \in M$ is a *Cauchy sequence* if $\forall \varepsilon > 0, \exists N$ s.t. $\forall m, n \geq N, d(x_n, x_m) < \varepsilon$.

Definition 2.4.2 (Bounded Sequence). A sequence $\{x_k\} \in M$ is *bounded* if $\exists x_0 \in M$ and $\exists R > 0$ s.t.

$$d(x_0, x_k) \leq R \quad \forall k.$$

Or, $x_k \in B(x_0, R) \quad \forall k$, where $B(x_0, R)$ denotes a closed ball centered at x_0 with radius R .

Definition 2.4.3 (Completeness). (M, d) is *complete* if every Cauchy sequence in M converges.

Example 2.4.4

- \mathbb{R}^1 and \mathbb{R}^n are complete
- $M = \mathbb{R}^1 \setminus \{0\}$ is not complete. For example, $x_k = \frac{1}{k}$ does not converge in $\mathbb{R}^1 \setminus \{0\}$.
- \mathbb{Q} is not complete.

Proposition 2.4.5 Basic Properties of Cauchy Sequence:

- Cauchy sequence is always bounded.
- Any converging sequence is always Cauchy.
- If a subsequence of a Cauchy sequence converges, then the original sequence converges.

Proof 1. (of ①): Suppose $\{x_k\}$ is Cauchy sequence. [WTS: $\exists x_0$ and $\exists R$ s.t. $x_k \in B(x_0, R) \quad \forall k$.]

Then, fix $\varepsilon = 1$. By Cauchy sequence, $\exists N$ s.t. $m, n \geq N \implies d(x_m, x_n) < \varepsilon = 1$. Define

$$\begin{aligned} R &= \max \{ \varepsilon, d(x_N, x_1), d(x_N, x_2), \dots, d(x_N, x_{N-1}) \} \\ &= \max \{ 1, d(x_N, x_k) : k = 1, \dots, N-1 \} \end{aligned}$$

Then, we have $d(x_k, x_N) \leq R \quad \forall k$, which implies that Cauchy sequence is bounded.

Q.E.D. ■

Theorem 2.4.6 Closedness and Completeness

Let (M, d) be a metric space.

- $N \subset M$ is complete $\implies N$ is closed. [Completeness is stronger than closedness]
- $N \subset M$ is closed and M is complete $\implies N$ is complete.

Remark 2.8 If (M, d) is a metric space and $N \subset M$, then (N, d) is also a metric space.

Proof 2.

- (of ①): Suppose $N \subset M$ is complete. [WTS: every sequence $x_k \in N$ that converges, the limit is in N .]

Given $\{x_k\} \in N$ with $x_k \rightarrow x \in M$. [WTS: $x \in N$.]

Since $\{x_k\} \in M$ converges, it is Cauchy. Further, as $N \subset M$ is complete, by definition, $x_k \rightarrow x \in N$ as desired. \square

- (of ②): Suppose $N \subset M$ is closed and M is complete. [WTS: Cauchy sequence $x_k \rightarrow x \in N$.]

Given $x_k \in N$ is a Cauchy sequence. Then, $x_k \in M$ as $N \subset M$. Since M is complete, we know $x_k \rightarrow x \in M$. Further, as N is closed, we know $x_k \rightarrow x \in N$. Hence, every Cauchy sequence converges in N . By definition, N is complete.

Q.E.D. \blacksquare

Definition 2.4.7 (Cluster Point). x is a *cluster point* of $\{x_k\}$ if $\forall \varepsilon > 0$, \exists infinitely many indices k s.t. $d(x_k, x) < \varepsilon$.

Proposition 2.4.8 Properties of Cluster Points:

- x is a cluster point $\iff \forall \varepsilon > 0, \forall N, \exists k > N$ s.t. $d(x_k, x) < \varepsilon$.
- x is a cluster point $\iff \exists$ subsequence $x_{n_k} \rightarrow x$.
- $x_k \rightarrow x \iff$ each subsequence $x_{n_k} \rightarrow x$.
- $x_k \rightarrow x \iff$ each subsequence has a further subsequence that converges to x .

3 Compactness and Connectedness

3.1 Compactness

Definition 3.1.1 (Cover and Subcover). Let $A \subset M$.

- A *cover* of a set $A \subset M$ is a collection $\{U_i\}$ of sets $U_i \subset M$ such that

$$\bigcup_i U_i \supset A.$$

- We say $\{U_i\}$ of A is an *open cover* if each U_i is open.
- A *subcover* of a given cover is a subcollection of $\{U_i\}$ whose union contains A .
- We say a cover is a *finite cover* if the subcollection contains finite number of sets.

Example 3.1.2

Suppose $A = [0, 1] \subset \mathbb{R}^1$. Consider

$$U_1 = (-1, 0.1), \quad U_2 = (0, 0.5), \quad U_3 = (0.5, 1).$$

$$U_4 = (0.2, 0.6), \quad U_5 = (0.8, 2), \quad U_6 = (0, 1).$$

Then,

- $\{U_1, \dots, U_6\}$ is a finite cover of A .
- It is also an open cover.
- $\{U_1, U_5, U_6\}$ is a subcover.

Definition 3.1.3 (Compactness). A set $A \subset M$ is called *compact* if every open cover of A has a finite subcover.

Definition 3.1.4 (Sequentially Compact). A set $A \subset M$ is *sequentially compact* if every sequence in A has a subsequence that converges to a point in A .

Definition 3.1.5 (Totally Bounded). A set $A \subset M$ is *totally bounded* if $\forall \varepsilon > 0, \exists$ finite set $\{x_1, x_2, \dots, x_N\} \subset M$ s.t.

$$A \subset \bigcup_{i=1}^N D(x_i, \varepsilon).$$

Remark 3.1

- A is sequentially compact $\implies A$ is closed and bounded.

Proof 1. Suppose A is unbounded. Fix $x_0 \in M$. For any $n \geq 1, \exists x_n \in A$ s.t.

$$d(x_n, x_0) \geq n.$$

By sequential compactness, \exists subsequence $x_{n_k} \rightarrow x \in A$ such that

$$\begin{aligned} d(x_{n_k}, x_0) &\leq d(x_{n_k}, x) + d(x, x_0) \\ &< \varepsilon + d(x, x_0). \end{aligned}$$

Take $\varepsilon = 1$, $d(x_{n_k}, x_0) < 1 + d(x, x_0)$ is a finite number. However, $d(x_{n_k}, x_0) \geq n_k$. \times As $n_k \rightarrow \infty$, $1 + d(x, x_0)$ is a finite number, we reach a contradiction. Hence, A must be bounded.

Q.E.D. ■

- A is totally bounded $\implies A$ is bounded.

Theorem 3.1.6 Bolzano-Weirstrass Theorem (B-W Thm.)

$A \subset M$ is compact $\iff A$ is sequentially compact.

Proof2.

Lemma 3.1.7: $A \subset M$ is compact $\implies A$ is closed.

Proof. [WTS: $M \setminus A$ is open.]

Fix $x \in M \setminus A$. For $n = 1, 2, \dots$, let $U_n = \left\{ y \mid d(x, y) > \frac{1}{n} \right\}$.

Claim $\{U_n \mid n = 1, 2, \dots\}$ is an open cover of A .

Proof. In fact, let $a \in A$. Then, $d(a, x) > 0$. By Archimedean, $\exists n$ s.t.

$$\frac{1}{n} < d(a, x).$$

This implies that $a \in U_n$. So, $a \in \bigcup_{i=1}^{\infty} U_i$. That is, $A \subset \bigcup_{i=1}^{\infty} U_i$. □

By the compactness, \exists finite subcover, say $\{U_1, \dots, U_N\}$. Thus,

$$A \subset \bigcup_{i=1}^N U_i = U_N = \left\{ y \mid d(y, x) > \frac{1}{N} \right\}.$$

Therefore,

$$D\left(x, \frac{1}{N}\right) = \left\{ y \mid d(y, x) < \frac{1}{N} \right\} \subset M \setminus A.$$

Hence, by definition, $M \setminus A$ is open, and so A must be closed. □

Lemma 3.1.8 (When is the converse of Lemma 3.1.7 true?):

$B \subset M$ is closed and M is compact $\implies B$ is compact.

Proof. Given an open cover $\{V_i \mid i \in I\}$ of B . [WTS: \exists a finite subcover of B .]

Since B is closed, $M \setminus B$ is open. Then,

$$\{V_i \mid i \in I\} \cup \{M \setminus B\} \text{ is an open cover of } M.$$

Since M is compact, \exists a finite subcover of M :

$$\{V_1, V_2, \dots, V_N\} \cup \{M \setminus B\}.$$

Note that

$$\bigcup_{i=1}^N V_i \supset B,$$

we know

$$\{V_1, V_2, \dots, V_N\} \text{ is a finite subcover of } B.$$

Hence, by definition, B is compact. \square

(\Rightarrow): Now, we prove the forward direction of the B-W Theorem. Let $A \subset M$ be compact. [WTS: A is sequentially compact]

- Set Up: Given a sequence $\{x_k\} \in A$. [WTS: $\exists x_{n_k} \rightarrow x \in A$]

By Lemma 3.1.7, compactness \implies closedness. Since A is closed, all converging sequence converges to some point in A . Hence, we only need to show \exists converging subsequence.

- Reduction: To this end, we may assume that $\{x_k\}$ contains a subsequence of distinct terms. Denote this subsequence by $\{y_k\}$. [WTS: $\{y_k\}$ has a convergent subsequence]

If $\{x_k\}$ does not contain subsequence of distinct terms, then $\{x_k\}$ is a constant sequence after sufficient terms. Therefore, it must converge and is trivial in this discussion.

- Suppose, for the sake of contradiction, $\{y_k\}$ does not have a convergent subsequence.
- **Claim** y_k 's are "isolated:" For each $k = 1, 2, \dots, \exists$ neighborhood U_k of y_k s.t. $y_j \notin U_k$ for any $j \neq k$.

Proof. Suppose, for the sake of contradiction, that the claim does not hold. Then, $\exists k$ with the property $\forall \varepsilon > 0, \exists j \neq k$ s.t. $y_j \in U_k = D(y_k, \varepsilon)$. Take $\varepsilon = \frac{1}{m}$. We obtain subsequence $y_{j_m} \in D\left(y_k, \frac{1}{m}\right), m = 1, 2, \dots$. Hence, when $m \rightarrow \infty, y_{j_m} \rightarrow y_k$.

This implies $\{y_k\}$ has a convergent subsequence. \times This contradicts with our assumption that $\{y_k\}$ does not have a convergent subsequence. Hence, the claim must be true. \square

- Now, proceed with the assumption that this claim is true.

Consider the set formed by elements in $\{y_k\}$:

$$B = \{y_1, y_2, \dots\}$$

Since $\{y_n\}$ has no convergent subsequence, B has no accumulation point, and so $\text{cl}(B) = B$, which implies B is closed.

By Lemma 3.1.8, B is compact.

On the other hand, $\{U_k\}$ is an open cover of B . But by claim, \exists no finite subcover. * This contradicts with the fact that B is compact. Thus, $\{y_k\}$ has a convergent subsequence, which converges to a point because A is closed.

□

(\Leftarrow): Now, let's consider the backward direction. Suppose $A \subset M$ is sequentially compact. [WTS: A is compact]

Let $\{u_i\}$ be an open cover of A . [WTS: \exists a finite subcover]

Claim (1) $\exists r > 0$ s.t. for each $y \in A$, $D(y, r) \subset U_i$ for some i . \implies Each point has a neighborhood of fixed size that is contained in some U_i .

Proof. Suppose otherwise. Then,

$$\forall r = \frac{1}{n} > 0, \exists y_n \in A \text{ s.t. } D\left(y_n, \frac{1}{n}\right) \text{ is not contained in any } U_i.$$

By assumption, A is sequentially compact. Then, $\{y_n\}$ has a convergent subsequence $z_n \rightarrow z \in A$.

On the other hand, U_i is an open cover of A , then $z_n \in U_{i_0}$ for some i_0 . Further, since U_{i_0} is open, $\exists \varepsilon > 0$ s.t. $D(z, \varepsilon) \subset U_{i_0}$.

Fix large N s.t.

$$d(z_N, z) < \frac{\varepsilon}{2}.$$

So,

$$D\left(z, \frac{\varepsilon}{2}\right) \subset D(z, \varepsilon) \subset U_{i_0}.$$

* This is a contradiction with our assumption that $D\left(y_n, \frac{1}{n}\right)$ is not contained in any U_i . Hence, the original claim is true. □

Claim (2) A is totally bounded.

Proof. Suppose otherwise. Then, $\exists \varepsilon > 0$ s.t. A cannot be covered by finite number of balls of radius ε . Choose $y_1 \in A$ and $y_2 \in A \setminus D(y_1, \varepsilon)$. Then, choose $y_3 \in A \setminus (D(y_1, \varepsilon) \cup D(y_2, \varepsilon))$. This process can go forever as A cannot be covered by finite number of balls of radius ε . So, we get sequence

$$y_n \in A \setminus (D(y_1, \varepsilon) \cup \dots \cup D(y_{n-1}, \varepsilon)).$$

We have a sequence $\{y_n\}$ with the property that

$$d(x_n, x_m) > \varepsilon \quad \forall n \neq m.$$

So, $\{y_n\}$ does not have a convergent subsequence.

Everything convergent must be Cauchy. $d(x_n, x_m) > \varepsilon$ implies not Cauchy, so it must be non-convergent. * This contradicts with the assumption that A is sequentially compact (has a subsequence converges to some point in A). Hence, this claim must be true. □

Now, let $r > 0$ be as in Claim (1). By Claim (2), $\exists y_1, y_2, \dots, y_N \in A$ s.t.

$$A \subset \bigcup_{j=1}^N D(y_j, r).$$

Then, further by Claim (1), we get $D(y_j, r) \subset U_{i_j}$. So,

$$A \subset \bigcup_{j=1}^N D(y_j, r) \subset \bigcup_{j=1}^N U_{i_j}.$$

Therefore, A can be covered by a finite subcover. Hence, A is compact.

Q.E.D. ■

Theorem 3.1.9

$A \subset M$ is compact $\iff A$ is complete and totally bounded.

Remark 3.2 So, if a set is not bounded/totally bounded, it cannot be compact.

Proof 3. (\Rightarrow): Done when proving B-W Thm. □

(\Leftarrow): Assume A is complete and totally bounded. [WTS: A is compact/sequentially compact]

Let $\{y_n\}$ be a sequence in A . [WTS: \exists subsequence y_{n_k} converges in A]

WLOG, we may assume $\{y_n\}$ is formed by distinct terms. If we don't get distinct terms, we will have a constant sequence when n gets sufficiently large. Hence, it converges in A and is trivial to discuss.

Since A is totally bounded, for $\varepsilon_1 = 1$, A is covered by finite number of balls:

$$D(x_1^{(1)}, \varepsilon_1), \dots, D(x_{L_1}^{(1)}, \varepsilon_1).$$

We can choose a subsequence $\{y_{1n}\}_{n=1}^\infty$ of $\{y_n\}$ that is contained one of the balls.

Repeat that for $\varepsilon_2 = \frac{1}{2}$, we have

$$A \subset D(x_1^{(2)}, \varepsilon_2) \cup \dots \cup D(x_{L_2}^{(2)}, \varepsilon_2).$$

We can choose a subsequence $\{y_{2n}\}_{n=1}^\infty$ of $\{y_n\}$ that is contained in one of the balls.

Continuing this process with $\varepsilon_m = \frac{1}{m}$, $m = 1, 2, \dots$. We obtain a subsequence $\{y_{mn}\}_{n=1}^\infty$ that is contained in a ball of radius $\varepsilon_m = \frac{1}{m}$. Then, we have the following subsequence:

$$\begin{array}{ccccccc} y_{11}, & y_{12}, & y_{13}, & \cdots, & y_{1n}, & \cdots \\ y_{21}, & y_{22}, & y_{23}, & \cdots, & y_{2n}, & \cdots \\ \vdots & & & & & \\ y_{m1}, & y_{m2}, & y_{m3}, & \cdots, & y_{mn}, & \cdots \\ \vdots & & & & & \end{array}$$

Each subsequence is a subsubsequence of the proceeding subsequence.

Select $y_{11}, y_{22}, y_{33}, \dots, y_{nn}, \dots$ to form a subsequence of $\{y_n\}$.

Denote this subsequence as $\{z_n\} = \{y_{nn}\}$.

A is complete. To show z_n converge in A , we only need to show z_n is Cauchy.

Claim $\{z_n\}$ is Cauchy.

Proof. Assume $n > m$:

$$d(z_n, z_m) < \frac{2}{m}.$$

When $m \rightarrow \infty$, $d(z_n, z_m) \rightarrow 0$. So, $\{z_n\}$ is Cauchy. \square

Since A is complete, $\{z_n\}$ is Cauchy, we have $z_n \rightarrow z \in A$. Hence, A is sequentially compact. By B-W Theorem, A is compact.

Q.E.D. \blacksquare

3.2 Compactness in \mathbb{R}^n

Theorem 3.2.1 Heine-Borel Theorem

A set $A \subset \mathbb{R}^n$ is compact $\iff A$ is bounded and closed.

Proof 1. (\implies) : True in general metric space. \square

(\impliedby) : Assume $A \subset \mathbb{R}^n$ is closed and bounded. [WTS: A is sequentially compact]

Given sequence $\{x_k\}$ in A , write

$$x_k = (x_k^{(1)}, x_k^{(2)}, \dots, x_k^{(n)}) \in A \subset \mathbb{R}^n.$$

A is bounded $\implies \{x_k\}$ is bounded $\implies \{x_k^{(1)}\}$ is bounded in \mathbb{R} .

$\implies \exists$ converging subsequence $\{x_{f_1(k)}^{(1)}\}_{k=1}^\infty$.

Similarly, $\{x_{f_1(k)}^{(2)}\}_{k=1}^\infty$ is bounded in \mathbb{R} . $\implies \exists$ converging subsequence $\{x_{f_2(k)}^{(2)}\}_{k=1}^\infty$.

In this way, we obtain subsequence

$$x_{f_n(k)} = (x_{f_n(k)}^{(1)}, x_{f_n(k)}^{(2)}, \dots, x_{f_n(k)}^{(n)})$$

with $x_{f_n(k)}^{(i)} \xrightarrow{k \rightarrow \infty} x^{(i)}$ for $i = 1, 2, \dots, n$. Hence,

$$x_{f_n(k)} \rightarrow (x^{(1)}, x^{(2)}, \dots, x^{(n)}) \in A.$$

Therefore, A is sequentially compact.

Q.E.D. \blacksquare

Remark 3.3 In Heine-Borel Theorem, (\impliedby) does not hold in general metric space. That is, A metric space that is closed and bounded does not imply compactness. For example, let $M =$ infinite set with discrete

metric

$$d(x, y) = \begin{cases} 0, & x = y \\ 1, & x \neq y. \end{cases}$$

M is closed and bounded, but M is not compact.

Example 3.2.2

- $A \subset \mathbb{R}^n$ is bounded $\implies \text{cl}(A)$ is compact.
- $A = [0, 1] \subset \mathbb{R}^1$ is compact.
- $A = (0, 1] \subset \mathbb{R}$ is not compact.
- \mathbb{R} is not compact because it is not totally bounded.

3.3 Nested Set Property

Theorem 3.3.1 Nested Set Property

Let F_k be a set of non-empty compact sets in M s.t.

$$F_{k+1} \subset F_k \quad \forall k = 1, 2, \dots$$

Then,

$$\bigcap_{k=1}^{\infty} F_k \neq \emptyset.$$

Proof 1. For each $k = 1, 2, \dots$, choose $x_k \in F_k$. Then, $\{x_k\} \subset F_1$. Since F_1 is compact, \exists subsequence

$$x_{f(k)} \xrightarrow{k \rightarrow \infty} x \in F_1.$$

Claim $x \in F_n \quad \forall n$.

Proof. Fix $n > 1$. Then, for large k ($\exists N$ s.t. $k \geq N$), we have $f(k) \geq n$. Then, $F_{f(k)} \subset F_n$. Recall that $x_{f(k)} \in F_{f(k)}$ and $x_{f(k)} \xrightarrow{k \rightarrow \infty} x$, then

$$x \in F_n$$

as F_n is closed. \square

Hence, $x \in \bigcap_{k=1}^{\infty} F_k \neq \emptyset$.


Q.E.D. \blacksquare

Remark 3.4 “Compact” cannot be replaced by “open,” “closed,” or “bounded open.”

3.4 Connectedness


Definition 3.4.1 (Path-Connected, Geometric Point of View). A set $A \subset M$ is *path-connected* if each pair of points $x, y \in A$ can be joined by a continuous path given by a continuous map

$$\varphi : [0, 1] \rightarrow A \quad \text{s.t.} \quad \varphi(0) = x \quad \text{and} \quad \varphi(1) = y$$



Example 3.4.2


- This is not path-connected:



- Let $\varphi : [0, 1] \rightarrow M$ be a continuous map. Then, $C = \varphi([0, 1]) \subset M$ (the image of this mapping) is path-connected.

Definition 3.4.3 (Disconnected Set, Topological Point of View). A set $A \subset M$ is said to be *disconnected* if \exists open sets $U, V \subset M$ that separate A :

- $U \cap V \cap A = \emptyset$
- $U \cap A \neq \emptyset$ and $V \cap A \neq \emptyset$
- $A \subset U \cup V$



Definition 3.4.4 (Connected Set). If a set is not disconnected, then it is *connected*.

Remark 3.5 It is easy to prove disconnectedness since we only need to find one pair of open sets satisfying the 4 conditions. To prove connectedness, we need to show \forall open sets $U, V \subset M$, they cannot satisfy the 4 conditions at the same time.

Theorem 3.4.5


Path-connectedness \implies connectedness

Proof 1. We will start the proof with the following claim (The proof is trivial, and so we omit the proof):

Claim 3.4.6 The interval $[a, b] \subset \mathbb{R}^1$ is connected.

Suppose, for the sake of contradiction, that $A \subset M$ is path-connected but not connected. Then, \exists open sets U, V that separates A as defined in Definition 3.4.3.

Fix $x \in U \cap A$ and $y \in V \cap A$.



Since A is path-connected, \exists a continuous map $\varphi : [0, 1] \rightarrow A$ with $\varphi(0) = x$ and $\varphi(1) = y$. Let

$$C = \varphi^{-1}(A \cap U) \subset [0, 1]$$

$$:= \{t \in [0, 1] \mid \varphi(t) \in A \cap U\}.$$

Similarly, we can define $D = \varphi^{-1}(A \cap V)$. Then, $0 \in C$ and $1 \in D$.

Claim 3.4.7 C is closed.

Proof. Let $t_k \in C$ s.t. $t_k \rightarrow t$. Then, by continuity of φ , $\varphi(t_k) \rightarrow \varphi(t) \in A$. Suppose, for the sake of contradiction, $\varphi(t) \notin U$. Then, $\varphi(t) \in V$. Since V is open, $\varphi(t_k) \in V$ for large k . Hence,

$$\varphi(t_k) \in A \cap U \cap V = \emptyset.$$

✱ We reach a contradiction. So, $\varphi(t) \in U$, which implies $t \in C$. As $t_k \rightarrow t \in C$, we have shown that C is closed. \square

Corollary 3.4.8 : By symmetry of C and D , D is also closed.

To derive a contradiction with Claim 3.4.6, note that

$$A \cap U \cap V = \emptyset,$$

which implies $C \cap D = \emptyset$. Therefore, the two open sets $(\mathbb{R} \setminus C)$ and $(\mathbb{R} \setminus D)$ separates $[0, 1]$. ✱ This contradicts with Claim 3.4.6 that $[0, 1]$ is connected. Hence, our assumption was wrong, and A must be path-connected and connected. In other words, path-connectedness \implies connectedness.

Q.E.D. \blacksquare

Remark 3.6 *The converse is not true.*

Example 3.4.9

$$\text{Suppose } A = \underbrace{\left\{ \left(x, \sin \frac{1}{x} \right) \mid x > 0 \right\}}_{\text{graph of } f(x) = \sin\left(\frac{1}{x}\right)} \cup \underbrace{\{(0, y) \mid -1 \leq y \leq 1\}}_{\text{segment of } y\text{-axis}} \subset \mathbb{R}^2.$$

Then, A is connected but not path-connected.

Proposition 3.4.10 : $A \subset \mathbb{R}^n$ open and connected \implies path-connected.

Proof2. (Sketch) Fix a point $x_0 \in A$ s.t.

$$B = \{y \in A \mid x_0 \text{ and } y \text{ can be joined by a continuous path } \in A\}.$$

Show:

- $B \neq \emptyset$ [$x_0 \in B$]
- B is open.
- B is closed in A .

Then, $B = A$. [If $B \neq A$, then $U = B$ and $V = A \setminus B$ separates $A \implies A$ is disconnected \implies contradiction, it must be $A = B$.]

Q.E.D. \blacksquare

Theorem 3.4.11 Equivalent Ways to Describe Connectedness

- In Definition 3.4.3, one can replace “open sets” by “closed sets.”

$A \subset M$ is disconnected $\iff \exists$ closed sets E, F s.t.

- $E \cap F \cap A = \emptyset$
- $E \cap A \neq \emptyset$ and $F \cap A \neq \emptyset$
- $A \subset E \cup F$

[Take complement of open sets, we get closed sets]

- In Definition 3.4.3, one can replace “ U, V ” by “disjoint open sets.”

$A \subset M$ is disconnected $\iff \exists$ disjoint open sets U_1 and V_1 s.t.

- $U_1 \cap V_1 \cap A = \emptyset$
- $U_1 \cap A \neq \emptyset$ and $V_1 \cap A \neq \emptyset$
- $A \subset U_1 \cup V_1$

Proof3. (Hint of ②): Consider the distance function $d(x, A \cap V)$ given fixed $x \in U \cap A$.

Claim $\forall x \in A \cap U$, *define* $d(x) = d(x, A \cap V) = \inf \{d(x, a) \mid a \in A \cap V\}$. *Then*, $d(x) > 0$. *Similarly*, $\forall y \in A \cap V$, *define* $d(y) = d(y, A \cap U) = \inf \{d(y, a) \mid a \in A \cap U\}$. *Then*, $d(y) > 0$.

Define open sets U_1, V_1 as follows:

$$U_1 = \left\{ D\left(x, \frac{1}{2}d(x)\right) \mid x \in A \cap U \right\} \quad \text{and} \quad V_1 = \left\{ D\left(y, \frac{1}{2}d(y)\right) \mid y \in A \cap V \right\}$$


We have the desired disjoint U_1 and V_1 .

Q.E.D. ■

4 Continuous Mappings

4.1 Continuity

Definition 4.1.1 (Maps). Suppose (M, d) and (N, ρ) are metric spaces. Let $A \subset M$. Then, $f : A \rightarrow N$ is a *map* (or a function)



Definition 4.1.2 (Continuous Maps). f is *continuous at a point* $x_0 \in A$ if


$$\lim_{\substack{x \rightarrow x_0 \\ x \in A}} f(x) = f(x_0).$$

f is *continuous in A* if it is continuous at each point in A .

Definition 4.1.3 (Limit of a Function). $b \in N$ is the limit of $f(x)$ at x_0 , written as

$$\lim_{x \rightarrow x_0} f(x) = b,$$

if $\forall \varepsilon > 0, \exists \delta > 0$ s.t. $x \in A$ and $d(x, x_0) < \delta \implies \rho(f(x), b) < \varepsilon$.



Definition 4.1.4 (Isolated Points). $x_0 \in A$ is an *isolated point* in A if $\exists \delta > 0$ s.t. $D(x_0, \delta) \cap A = \{x_0\}$.

Remark 4.1

- The continuous definition implies three things: the function is defined, the limit exists, and the limit value equals the function value.
- A point is either an isolated point or an accumulation point.
- For the limit definition, x_0 is not required to be in A . For example,

$$f(x) = \frac{\sin(x)}{x}, \quad x \in (0, 1) \quad \lim_{x \rightarrow 0} f(x) = 0 \notin (0, 1).$$

- If x_0 is an isolated point in A , then $\lim_{x \rightarrow x_0} f(x) = f(x_0)$ is always true. Therefore, any function $f(x)$ is continuous at isolated points.

Example 4.1.5

- $f(x) = x : \mathbb{R}^n \rightarrow \mathbb{R}^n$ (identity function) is continuous
- $g(x) = \begin{cases} x, & 0 \leq x \leq 1 \\ 2x, & 1 < x \leq 3 \end{cases} : [0, 3] \rightarrow \mathbb{R}^1$ is continuous at every point except for $x = 1$.
- $h(x) = \begin{cases} x, & x \neq 1 \\ 3, & x = 1 \end{cases} : \mathbb{R} \rightarrow \mathbb{R}$ is continuous at every point except $x = 1$.

Theorem 4.1.6 Equivalent Conditions for Continuity

Let $f : A \subset M \rightarrow N$. The following are equivalent:

- f is continuous on A .
- For each converging sequence $x_k \rightarrow x \in A$, $f(x_k) \rightarrow f(x)$.


Remark 4.2 Continuous map preserves the convergence of sequences

- For each open set $U \subset N$, the pre-image $f^{-1}(U) \subset A$ is open relative to A . That is

$$f^{-1}(U) = \{x \in A \mid f(x) \in U\} = A \cap V, \quad \text{where } V \subset M \text{ is open.}$$

- For each close set $F \subset N$, the pre-image $f^{-1}(F) \subset A$ is closed relative to A . That is,

$$f^{-1}(F) = A \cap E, \quad \text{where } E \subset M \text{ is closed.}$$



Proof 1. We will prove equivalence by the following cycle: ① \implies ② \implies ④ \implies ③ \implies ①.

(① \implies ②): Given sequence $x_k \in A$ with $x_k \rightarrow x \in A$. [WTS: $\lim_{k \rightarrow \infty} f(x_k) = f(x)$]

(② \implies ④): Fix closed set $F \in N$. [WTS: $f^{-1}(F) = A \cap \text{cl}(f^{-1}(F))$] It is trivial that $f^{-1}(F) \subset A \cap \text{cl}(f^{-1}(F))$. So, we only need to prove the “ \supset ” direction. Given $x \in A \cap \text{cl}(f^{-1}(F))$, \exists sequence $x_n \in f^{-1}(F) \subset A$ s.t. $x_n \rightarrow x$. Then, $y_n = f(x_n) \rightarrow f(x) \in F$ by ② and closedness. So, $x \in f^{-1}(F)$. That is, $A \cap \text{cl}(f^{-1}(F)) \supset f^{-1}(F)$. Hence, $f^{-1}(F) = A \cap \text{cl}(f^{-1}(F))$, implying $f^{-1}(F)$ is closed in A .

(④ \implies ③): [Use complement: $U \subset N$ is open $\iff F = N \setminus U$ is closed]

(③ \implies ①): Given $x_0 \in A$. [WTS: $\lim_{x \rightarrow x_0} f(x) = f(x_0)$] Fix any $\varepsilon > 0$. [WTS: $\exists \delta > 0$ s.t. $d(x, x_0) < \delta \implies \rho(f(x), f(x_0)) < \varepsilon$] Let $U = D(f(x_0), \varepsilon) < r$ is open. By ③, $f^{-1}(U)$ is open in A . i.e.,

$$f^{-1}(U) = A \cap V, \quad V \subset M \text{ is open.}$$

Note that $x_0 \in f^{-1}(U) \implies x_0 \in V$. Since V is open, $\exists \delta > 0$ s.t. $D(x, \delta) \subset V$. [WTS: $x \in A$, $d(x, x_0) < \delta \implies \rho(f(x), f(x_0)) < \varepsilon$] Suppose $x \in A$ with $d(x, x_0) < \delta$. Then, $x \in A$ and $x \in V$. That is, $x \in A \cap V = f^{-1}(U)$. Hence, $f(x) \in U$. By definition of U , we get $\rho(f(x), f(x_0)) < \varepsilon$ as desired.

Q.E.D. ■

4.2 Properties of Continuous Mappings

Theorem 4.2.1 Images of Compact and Connected Sets

Suppose $f : M \rightarrow N$ is continuous. Then,

- If $K \subset M$ is compact, then $f(K)$ is also compact.
- If $B \subset M$ is connected, then $f(B)$ is also connected.

Proof 1.

- Let x_k be a sequence in K . Then, $y_k = f(x_k)$ is a sequence in $f(K)$. [WTS: $f(K)$ is sequentially compact.] Suppose K is compact, $\exists x_{k_j} \rightarrow x_0 \in K$ when $j \rightarrow \infty$. By continuity of f , $f(x_{k_j}) \rightarrow f(x_0) \in f(K)$ when $k \rightarrow \infty$. So, for sequence $y_k = f(x_k)$, we find a subsequence $f(x_{k_j}) \rightarrow f(x_0) \in f(K)$. So, $f(K)$ is sequentially compact. □
- Given connected set $B \subset M$. Assume, for the sake of contradiction, that $f(B)$ is disconnected. Then, \exists open sets U, V s.t. $f(B) \cap U \cap V = \emptyset$ and $f(B) \cap U \neq \emptyset$, $f(B) \cap V \neq \emptyset$, $f(B) \subset U \cup V$. [We can derive that B is also disconnected, which is a contradiction.] So, it must be that $f(B)$ is also connected.

Q.E.D. ■

Theorem 4.2.2 Operations on Continuous Mapping

Addition, multiplication, divisions, and compositions of continuous functions (if they are well-defined) are also continuous.

Example 4.2.3

If $f(x) = \mathbb{R} \rightarrow \mathbb{R}$, $g : \mathbb{R} \rightarrow \mathbb{R}$ are continuous, then, $f(x)g(x)$ is also continuous.

Proof 2. Denote $F(x) = f(x)g(x)$. Then,

$$\begin{aligned} |F(x) - F(x_0)| &= |f(x)g(x) - f(x_0)g(x_0)| \\ &\leq |f(x)g(x) - f(x)g(x_0)| + |f(x)g(x_0) - f(x_0)g(x_0)| \\ &= |f(x)||g(x) - g(x_0)| + |g(x_0)||f(x) - f(x_0)| \\ &\vdots \\ &< \varepsilon \end{aligned}$$

Q.E.D. ■

Theorem 4.2.4 Maximum/Minimum Property

Let $K \subset M$ be compact and $f : K \rightarrow \mathbb{R}$ be continuous. Then,

- f is bounded on K (i.e., $f(K)$ is a bounded set)
- $\exists x_0, x_1 \in K$ s.t.

$$f(x_1) = \max_{x \in K} f(x) \quad \text{and} \quad f(x_0) = \min_{x \in K} f(x).$$

That is, $f(x_0) \leq f(x) \leq f(x_1) \quad \forall x \in K$.

Proof 3.

- Since K is compact and f is continuous, $f(K)$ is compact. Since $f(K) \subset \mathbb{R}$ is compact, $f(K)$ is closed and bounded.
- Since $f(K)$ is bounded, we know $\inf(f(K))$ and $\sup(f(K))$ exist and are finite. Further since $f(K)$ is closed, $\inf(f(K)), \sup(f(K)) \in f(K)$. Hence, $\exists x_0 = \inf(f(K))$ and $x_1 = \sup(f(K))$ s.t.

$$f(x_0) \leq f(x) \leq f(x_1) \quad \forall x \in K.$$

Q.E.D. ■

Remark 4.3

- The condition “compact” cannot be removed.

Example 4.2.5

$$f(x) = \frac{1}{x} : (0, 1) \rightarrow \mathbb{R} \text{ is continuous but not bounded}$$

$f(x) = x : (0, 1) \rightarrow \mathbb{R}$ is bounded, but does not have max/min values

- The condition “continuity” cannot be removed.


Example 4.2.6

Consider function $f : [0, 1] \rightarrow \mathbb{R}$ by $f(x) = \begin{cases} \frac{1}{x}, & x > 0 \\ 2, & x = 0. \end{cases}$ Although $[0, 1]$ is compact, $f(x)$ is not continuous, and f is not bounded and does not have max/min values on $[0, 1]$.

- We don't need differentiability here.

Theorem 4.2.7 Intermediate Value Theorem (IVT)

Let $K \subset M$ be connected and $f : K \rightarrow \mathbb{R}$ be continuous. Suppose $x, y \in K$ with $f(x) < f(y)$. Then, for any intermediate value c s.t. $f(x) < c < f(y)$, $\exists z \in K$ with $x < z < y$ s.t. $f(z) = c$.



Proof 4. Let $K \subset M$ be connected and $f : K \rightarrow \mathbb{R}$ be continuous. Suppose $x, y \in K$ with $f(x) < f(y)$. Assume, for the sake of contradiction, $\exists c$ with $f(x) < c < f(y)$ s.t. $c \notin f(K)$.

Since K is connected and f is continuous, $f(K)$ is also connected. However, $U = (-\infty, c)$ and $V = (c, +\infty)$ separate $f(K)$, implying $f(K)$ is not connected. ✱ We reach a contradiction. So, such a c does not exist.

Q.E.D. ■

Example 4.2.8 Application of IVT I

Let $f(x)$ be a polynomial of odd degree. Then, f has at least one real root.

Proof 5. Suppose $f(x) : \mathbb{R} \rightarrow \mathbb{R}$ is continuous. Write $f(x)$ as

$$f(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0,$$

where $a_n \neq 0$ and $n = 2k + 1$ is odd.

WLOG, suppose $a_n > 0$. Then,

$$\lim_{x \rightarrow -\infty} f(x) = -\infty \quad \text{and} \quad \lim_{x \rightarrow \infty} f(x) = \infty.$$

So, $\exists x, y \in \mathbb{R}$ s.t. $f(x) < 0$ and $f(y) > 0$. Therefore, by IVT, $\exists x_0 \in \mathbb{R}$ s.t. $f(x_0) = c = 0$.

Q.E.D. ■

Definition 4.2.9 (Fixed Point). x is a *fixed point* of f if $f(x) = x$.

Example 4.2.10 Application of IVT II

Let $f : [1, 2] \rightarrow [0, 3]$ be continuous with $f(1) = 0$, $f(2) = 3$. Then, f has a fixed point.

Proof 6. Apply IVT to a new function: $F(x) = f(x) - x$. Take $c = 0$ as the intermediate value.

Q.E.D. ■

4.3 Uniform Continuity (UC)

Definition 4.3.1 (Uniform Continuity (UC)). A function $f : A \subset M \rightarrow N$ is *uniformly continuous* on A if $\forall \varepsilon > 0, \exists \delta > 0$ s.t. $x, y \in A$ and $d(x, y) < \delta \implies \rho(f(x), f(y)) < \varepsilon$.

Remark 4.4

- For uniform continuity, the δ depends only on ε not on points.
- For continuity (at x_0), the δ may depend on ε and the point x_0 .

Example 4.3.2

Consider $f(x) = \frac{1}{x} : (0, 1) \rightarrow \mathbb{R}$. f is continuous at any point $x_0 \in (0, 1)$. But to satisfy

$$|x - x_0| < \delta \implies |f(x) - f(x_0)| = \left| \frac{1}{x} - \frac{1}{x_0} \right| = \frac{|x - x_0|}{|x \cdot x_0|} < \varepsilon$$

we need to pick

$$|x - x_0| = \delta = \min \left\{ \frac{1}{2} x_0^2 \varepsilon, \frac{1}{2} x_0 \right\}.$$

Theorem 4.3.3 Uniform Continuity on Compact Set

Let $f : K \subset M \rightarrow N$ be continuous and K be compact. Then, f is uniformly continuous on K .

Proof 1. Fix $\varepsilon > 0$. For each $x \in K$, since f is continuous at x , $\exists \delta_x$ s.t. for $y \in K$ with $d(x, y) < \delta_x$, we have $\rho(f(x), f(y)) < \frac{\varepsilon}{2}$.

Consider the open cover of K : $\left\{ D\left(x, \frac{\delta_x}{2}\right) \mid x \in K \right\}$. Since K is compact, \exists subcover:

$$D\left(x_i, \frac{\delta_{x_i}}{2}\right), \quad i = 1, 2, \dots, L$$

Finally, let

$$\delta = \min_{1 \leq i \leq L} \left\{ \frac{\delta_{x_i}}{2} \right\}.$$

Claim $x, y \in K$ with $d(x, y) < \delta \implies \rho(f(x), f(y)) < \varepsilon$.

Proof. Note that

$$\begin{aligned} d(y, x_i) &\leq d(y, x) + d(x, x_i) \\ &< \delta + \frac{\delta_{x_i}}{2} \\ &< \delta_{x_i}. \end{aligned}$$

One can continue to show that $\rho(f(x), f(y)) < \varepsilon$. \square

Q.E.D. \blacksquare

Definition 4.3.4 (Lipschitz Continuity). A function $f : A \subset M \rightarrow N$ is called *Lipschitz* if \exists constant L s.t.

$$\rho(f(x), f(y)) \leq L \cdot d(x, y) \quad \forall x, y \in A.$$

Theorem 4.3.5 Lipschitz and Uniform Continuity

If $f : A \subset M \rightarrow N$ is Lipschitz, then f is uniformly continuous in A .

Corollary 4.3.6 : Suppose $f : (a, b) \rightarrow \mathbb{R}$ is differentiable and $\exists M > 0$ s.t. $|f'(x)| \leq M \quad \forall x \in (a, b)$. Then, f is Lipschitz.

Proof2. Given $x, y \in (a, b)$. Then,

$$\begin{aligned} |f(y) - f(x)| &= |f'(z)(y - x)| && \text{[Mean Value Theorem]} \\ &\leq M|y - x|. \end{aligned}$$

Q.E.D. \blacksquare

Example 4.3.7 Lipschitz Functions

$f(x) = x$ and $f(x) = \sin x$ are Lipschitz functions.

Remark 4.5

- If f has bounded derivative (or slope), then f is uniformly continuous.

- But if f is differentiable and uniformly continuous, f may not have bounded derivative.
- Open End-ed Questions:
 - $f : \mathbb{R} \rightarrow \mathbb{R}$ is bounded and continuous, f may not be uniformly continuous.
 - $f, g : \mathbb{R} \rightarrow \mathbb{R}$ are uniformly continuous, $f \cdot g$ is not uniformly continuous in general.
 - But if f , or g , or both are bounded and uniformly continuous, is $f \cdot g$ uniformly continuous?

4.4 Differentiability

Remark 4.6 Starting from this section, we will only consider functions $f : \text{an interval} \rightarrow \mathbb{R}$.

Definition 4.4.1 (Differentiability). A function f is *differentiable* at a point x_0 if it is defined in an open interval that contains x_0 and its derivative exists:

$$f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0}, \quad (\text{D})$$

or equivalently, set $h = x - x_0$,

$$f'(x_0) = \lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}.$$

Remark 4.7 (Interpretation)

- Rewrite (D) as

$$\lim_{x \rightarrow x_0} \left[\frac{f(x) - f(x_0) - f'(x_0)(x - x_0)}{x - x_0} \right] = 0.$$

This implies the function $y = f(x)$ can be approximated by the linear function

$$y = f(x_0) + f'(x_0)(x - x_0)$$

in a neighborhood of x_0 .

- Rewrite (D) as

$$\lim_{\Delta x \rightarrow 0} \left[\frac{f(x_0 + \Delta x) - f(x_0)}{\Delta x} - f'(x_0) \right] = 0.$$

this implies the slope of tangent line is the limit of the slope of secant lines.

Theorem 4.4.2 Continuity of Differentiable Functions

Suppose $f : A \subset M \rightarrow N$ is differentiable at x_0 . Then, it is continuous at x_0 .

Proof 1. Given $\varepsilon > 0$. Since

$$\lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0} = f'(x_0),$$

$\exists \delta_1 > 0$ s.t. $|x - x_0| < \delta_1 \implies \left| \frac{f(x) - f(x_0)}{x - x_0} \right| < |f'(x)| + 1$. Choose

$$\delta = \min \left\{ \frac{\varepsilon}{|f'(x)| + 1}, \delta_1 \right\}.$$

So, when $|x - x_0| < \delta$, we have

$$\begin{aligned} |f(x) - f(x_0)| &= \frac{|f(x) - f(x_0)|}{|x - x_0|} \cdot |x - x_0| \\ &< (|f'(x)| + 1) \cdot \frac{\varepsilon}{|f'(x)| + 1} \\ &= \varepsilon. \end{aligned}$$

Q.E.D. ■

Remark 4.8 *The converse is not true: continuity $\not\Rightarrow$ differentiability. Counterexample: $f(x) = |x|$.*

Proof 2. (Another Approach) Note that

$$\begin{aligned} \lim_{x \rightarrow x_0} (f(x) - f(x_0)) &= \lim_{x \rightarrow x_0} \left[\frac{f(x) - f(x_0)}{x - x_0} \right] (x - x_0) \\ &= \lim_{x \rightarrow x_0} \left[\frac{f(x) - f(x_0)}{x - x_0} \right] \lim_{x \rightarrow x_0} (x - x_0) \quad [\text{Product Rule of Limit}] \\ &= f'(x) \cdot 0 \\ &= 0. \end{aligned}$$

So, the function is continuous.

Q.E.D. ■

Theorem 4.4.3 Rules of Differentiation

- Constant multiple rule:

$$(kf)'(x_0) = k \cdot f'(x_0).$$

- Sum rule:

$$(f + g)'(x_0) = f'(x_0) + g'(x_0)$$

- Product rule:

$$(fg)'(x_0) = f'(x_0)g(x_0) + f(x_0)g'(x_0).$$

- Quotient rule:

$$\left(\frac{f}{g} \right)'(x_0) = \frac{f'(x_0)g(x_0) - f(x_0)g'(x_0)}{(g(x_0))^2}$$

- Chain rule:

$$(g \circ f)'(x_0) = g'(f(x_0)) \cdot f'(x_0).$$

Lemma 4.4.4 : If $f : (a, b) \rightarrow \mathbb{R}$ is differentiable and f has a max (or min) at $c \in (a, b)$, then $f'(c) = 0$.

Proof 3. Assume f has a max at $c \in (a, b)$. Then,

$$f'(c) = \lim_{h \rightarrow 0} \frac{f(h + c) - f(c)}{h}.$$

[WTS: $f'(c) \geq 0$ and $f'(c) \leq 0$.]

As f has a max at c , $f(h+c) \leq f(c)$, and so

$$f(h+c) - f(c) \leq 0.$$

Case I $h > 0$:

$$f'(c) = \lim_{h^+ \rightarrow 0} \frac{f(h+c) - f(c)}{h} \leq 0.$$

Case II $h < 0$:

$$f'(c) = \lim_{h^- \rightarrow 0} \frac{f(h+c) - f(c)}{h} \geq 0.$$

As $f'(c) \geq 0$ and $f'(c) \leq 0$, it must be that $f'(c) = 0$.

Q.E.D. ■

Theorem 4.4.5 Rolle's Theorem

Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous and f be differentiable on (a, b) . If $f(a) = f(b) = 0$, then $\exists c \in (a, b)$ s.t. $f'(c) = 0$.

Proof 4. f has max and min on $[a, b]$ as $[a, b]$ is compact. [WTS: This max/min occur in (a, b) .]

Since $f(a) = f(b) = 0$, then max and min cannot both occur at the endpoint (i.e., either max or min occur in (a, b)) unless f is the constant function $f(x) = 0$.

Now, by Lemma 4.4.4, $\exists c \in (a, b)$ s.t. $f'(c) = 0$, where c is either the max or min.

Q.E.D. ■


Theorem 4.4.6 Mean Value Theorem (MVT)

Suppose f is continuous on $[a, b]$ and differentiable on (a, b) . Then, $\exists c \in (a, b)$ s.t.

$$f(b) - f(a) = f'(c)(b - a) \quad \text{or} \quad f'(c) = \frac{f(b) - f(a)}{b - a}.$$

Remark 4.9 Rolle's Theorem is a special case of MVT. We will use the special case to prove the general case.

Proof 5.



Construct $\varphi(x)$:

$$\varphi(x) = f(x) - \left[f(a) + \frac{f(b) - f(a)}{b - a}(x - a) \right].$$

One can verify the following:

- $\varphi(a) = 0$;
- $\varphi(b) = 0$; and
- φ is continuous and differentiable.

Then, apply Rolle's Theorem to $\varphi(x)$: $\exists c \in (a, b)$ s.t.

$$\varphi'(c) = 0.$$

Note that $\varphi'(c) = f'(c) - \frac{f(b) - f(a)}{b - a} = 0$, we have

$$f'(c) = \frac{f(b) - f(a)}{b - a}.$$

Q.E.D. ■

Remark 4.10 (Geometric Interpretation) *There is at least one point where the instant change of rate is the same as the average change of rate.*

Definition 4.4.7 (Monotonecity).

- We say $f(x)$ is *increasing* (or *strictly increasing*) at a point x_0 if \exists open interval (a, b) containing x_0 with:
 - $a < x < x_0 \implies f(x) \leq f(x_0)$ (or $f(x) < f(x_0)$);
 - $x_0 < x < b \implies f(x) \geq f(x_0)$ (or $f(x) > f(x_0)$).
- Similar definition for decreasing (or strictly decreasing) at a point x_0 .
- $f(x)$ is increasing (or strictly increasing) on an interval I if for $x_1, x_2 \in I$

$$x_1 < x_2 \implies f(x_1) \leq f(x_2) \quad (\text{or } f(x_1) < f(x_2)).$$

- Similar definition for decreasing (or strictly decreasing) on an interval.

Theorem 4.4.8 Local Monotonecity and Derivative

Let f be differentiable at x_0 . Then,

- f increasing at $x_0 \implies f'(x_0) \geq 0$; f decreasing at $x_0 \implies f'(x_0) \leq 0$.
- $f'(x_0) > 0 \implies f$ strictly increasing at x_0 ; $f'(x_0) < 0 \implies f$ strictly decreasing at x_0 .

Proof 6. (of ①): Suppose f is increasing at x_0 . Then

$$\begin{aligned} f(x_0 + h) - f(x_0) &\geq 0 \quad \text{when } h > 0 \\ &\leq 0 \quad \text{when } h < 0. \end{aligned}$$

Then,

$$f'(x_0) = \lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h} \geq 0.$$

(of ②): Suppose $f'(x_0) > 0$. Then, for $\varepsilon = \frac{1}{2}f'(x_0) > 0$, $\exists \delta > 0$ s.t.

$$0 < |h| < \delta \implies \left| \frac{f(x_0 + h) - f(x_0)}{h} - f'(x_0) \right| < \varepsilon = \frac{1}{2}f'(x_0).$$

$$-\frac{1}{2}f'(x_0) < \frac{f(x_0 + h) - f(x_0)}{h} - f'(x_0) < \frac{1}{2}f'(x_0) \implies 0 < \frac{1}{2}f'(x_0) < \frac{f(x_0 + h) - f(x_0)}{h} < \frac{3}{2}f'(x_0).$$

When $x < x_0$, $h = x - x_0 < 0$. As $\frac{f(x_0 + h) - f(x_0)}{h} > 0$,

$$f(x_0 + h) - f(x_0) = f(x) - f(x_0) < 0 \implies f(x) < f(x_0)$$

When $x > x_0$, $h = x - x_0 > 0$,

$$f(x_0 + h) - f(x_0) = f(x) - f(x_0) > 0 \implies f(x) > f(x_0).$$

Hence, f is strictly increasing.

Q.E.D. ■

Theorem 4.4.9 Global Monotonecity and Derivative

Let f be continuous on $[a, b]$ and differentiable on (a, b) . Then,

- $f'(x) \geq 0 \quad \forall x \in (a, b) \implies f$ increasing on $[a, b]$.
- $f'(x) \leq 0 \quad \forall x \in (a, b) \implies f$ decreasing on $[a, b]$.
- $f'(x) > 0 \quad \forall x \in (a, b) \implies f$ strictly increasing on $[a, b]$.
- $f'(x) < 0 \quad \forall x \in (a, b) \implies f$ strictly decreasing on $[a, b]$.

Theorem 4.4.10 Local Max/Min and Derivative

Suppose f is continuous on $[a, b]$ and twice differentiable on (a, b) . Let $x_0 \in (a, b)$.

- $f'(x_0) = 0$ and $f''(x_0) > 0 \implies x_0$ is a strict local min of f .
- $f'(x_0) = 0$ and $f''(x_0) < 0 \implies x_0$ is a strict local max of f .

Proof 7. (of ①) By Theorem 3.3.8(2), $f''(x_0) > 0 \implies f'(x)$ is strictly increasing at x_0 . Then,

- $f'(x) < f'(x_0) = 0 \quad \forall x \in (x_0 - \delta, x_0) \implies f(x)$ strictly decreasing on $(x_0 - \delta, x_0)$
- $f'(x) > f'(x_0) = 0 \quad \forall x \in (x_0, x_0 + \delta) \implies f(x)$ strictly increasing on $(x_0, x_0 + \delta)$.

Q.E.D. ■

Theorem 4.4.11 Inverse Function Theorem (IFT)

Suppose $f'(x) > 0 \quad \forall x \in (a, b)$ (or, $f'(x) < 0 \quad \forall x \in (a, b)$). Then,

- $f : (a, b) \rightarrow \mathbb{R}$ is a bijection onto its image
- Inverse f^{-1} is differentiable on its domain.
- $(f^{-1})'(y) = \frac{1}{f'(x)}$, where $y = f(x)$.

Proof 8. Assume $f'(x) > 0 \quad \forall x \in (a, b)$. Then, f is strictly increasing. Then, f is 1-to-1 function $\implies f$ is a bijection $\implies f^{-1}$ exists. [WTS: f^{-1} is continuous.]

Let U be an open set in (a, b) . [WTS: $(f^{-1})^{-1}(U) = f(U)$ is open.]

Finally, write $y = f(x)$. Then, $x = f^{-1}(y)$. Let $y_0 = f(x_0)$. Then,

$$\begin{aligned}
 (f^{-1})'(y_0) &= \lim_{y \rightarrow y_0} \frac{f^{-1}(y) - f^{-1}(y_0)}{y - y_0} \\
 &= \lim_{y \rightarrow y_0} \frac{x - x_0}{f(x) - f(x_0)} \\
 &= \lim_{x \rightarrow x_0} \frac{1}{\frac{f(x) - f(x_0)}{x - x_0}} \\
 &= \frac{1}{\lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0}} \\
 &= \frac{1}{f'(x_0)}.
 \end{aligned}$$

Q.E.D. ■

4.5 Integration

Definition 4.5.1 (Riemann Integrable). Let $A \subset \mathbb{R}$ be bounded and $f : A \rightarrow \mathbb{R}$ be a bounded function.

[We want to make sense $\int_A f(x)dx$.]


- Partition the interval:

If interval $[a, b] \supset A$ and extend function $f(x)$ to $[a, b]$ by letting $f(x) = 0 \quad \forall x \notin A$.

Partition the interval $[a, b]$ by points: $a = x_0 < x_1 < \dots < x_n = b$. Denote P by

$$P = \{x_0, x_1, x_2, \dots, x_n\}.$$

- Form Upper and Lower Sum of f .



For any fixed partition, let

$$U(f, P) = \sum_{i=1}^n \sup \{f(x) \mid x \in [x_{i-1}, x_i]\} (x_i - x_{i-1})$$

is the upper sum, and

$$L(f, P) = \sum_{i=1}^n \inf \{f(x) \mid x \in [x_{i-1}, x_i]\} (x_i - x_{i-1})$$

is the lower sum.

Claim Suppose $m \leq f(x) \leq M$. Then,

$$m(b - a) \leq L(f, P) \leq U(f, P) \leq M(b - a).$$

- Upper integral and Lower integral are defined as

$$\int_A^{\bar{}} f = \inf \{U(f, P) : P \text{ is a partition}\} \quad (\text{Upper Integral})$$

$$\int_A^{\underline{}} f = \sup \{L(f, P) : P \text{ is a partition}\} \quad (\text{Lower Integral})$$

- We say a function f is *Riemann integrable* if

$$\int_A^{\bar{}} f = \int_A^{\underline{}} f,$$

and we write

$$\int_A f = \int_A^{\bar{}} f = \int_A^{\underline{}} f.$$

Example 4.5.2 Riemann Integrable

- Define $f : [0, 1] \rightarrow \mathbb{R}$ by

$$f(x) = \begin{cases} 0 & \text{if } x \text{ is rational} \\ 1 & \text{if } x \text{ is irrational.} \end{cases}$$

Then, for any partition P ,

$$U(f, P) = \sum_{i=1}^n 1 \cdot (x_i - x_{i-1}) = 1$$

and

$$L(f, P) = \sum_{i=1}^n 0 \cdot (x_i - x_{i-1}) = 0.$$

So,

$$\int_A^{\bar{}} f \neq \int_A^{\underline{}} f \implies f \text{ is not integrable}$$


- Compute $\int_0^1 x \, dx$ and $\int_0^1 x \, dx$.

Hint: Consider partition $P_n = \left\{ 0 < \frac{1}{n} < \frac{2}{n} < \dots < \frac{n}{n} \right\}$.

Lemma 4.5.3 : Let $f : [a, b] \rightarrow \mathbb{R}$ be bounded. If P, P' are partitions of $[a, b]$ with $P \subset P'$ (P' is a refinement of P), then

$$L(f, P) \leq L(f, P') \leq U(f, P') \leq U(f, P).$$

Remark 4.11 *In words, when the partition gets finer, lower sum increases but upper sum decreases.*



Proposition 4.5.4 :

$$\int_a^b f \leq \int_a^b f$$

Proof 1. For any fixed partition P and Q . As $P \subset P \cup Q$ and $Q \subset P \cup Q$, by Lemma 4.5.4, we have

$$L(f, P) \leq L(f, P \cup Q) \leq U(f, P \cup Q) \leq U(f, Q).$$

Then,

$$\int_a^b f = \sup_P L(f, P) \leq U(f, Q) \quad \text{for any } Q$$

So,

$$\int_a^b f \leq \inf_Q U(f, Q) = \int_a^b f.$$

Q.E.D. ■

Theorem 4.5.5

- If $f : [a, b] \rightarrow \mathbb{R}$ is bounded and is continuous at all but finite many points, then f is integrable.
- If f is increasing or decreasing on $[a, b]$, then f is integrable.

Proof 2.

- (Proof of ①): Observe that \forall partition P , $L(f, P) \leq \int_a^b f \leq \int_a^b f \leq U(f, P)$. [To prove a function is integrable, it's sufficient to show that $\forall \varepsilon > 0$, \exists partition P s.t. $U(f, P) - L(f, P) < \varepsilon$.]

- Suppose f is continuous on $[a, b]$ except at a_1, a_2, \dots, a_k . Since f is bounded, $\exists m, M$ s.t. $m \leq f(x) \leq M \quad \forall x \in [a, b]$. Choose partition P_1 s.t. each subinterval containing some a_i has length $\leq \frac{\varepsilon}{2} \cdot \frac{1}{2k(M-m)}$.

Let K be the union of the remaining subinterval in P_1 . Then, K is compact and f is continuous on K . So, f is uniformly continuous on K . That is,

$$\exists \delta > 0 \text{ s.t. } x_1, x_2 \in K \text{ s.t. } |x_1 - x_2| < \delta \implies |f(x_1) - f(x_2)| < \frac{\varepsilon}{2(b-a)}.$$

- Construct the refinement P of P_1 so that each subinterval in P not containing some a_i has length $< \delta$. So,

$$P = \{a = x_0 < x_1 < \dots < x_n = b\} \quad \text{and} \quad I_j = [x_{j-1}, x_j].$$

Denote

$$M_j = \sup_{I_j} f(x) \quad \text{and} \quad m_j = \inf_{I_j} f(x).$$

If I_j contains some a_i , then $m \leq m_j \leq M_j \leq M$.

If I_j contains no discontinuous points, then $I_j \subset K$, and

$$M_j - m_j = \max - \min < \frac{\varepsilon}{2(b-a)}.$$

– Finally, we have

$$\begin{aligned}
 U(f, P) - L(f, P) &= \sum_{j=1}^n (M_j - m_j)(x_j - x_{j-1}) \\
 &= \sum_{a_i \in I_j} (M_j - m_j)(x_j - x_{j-1}) + \sum_{a_i \notin I_j} (M_j - m_j)(x_j - x_{j-1}) \\
 &< \underbrace{2k}_{\text{worse case: } 2k \text{ such intervals}} \cdot \underbrace{(M - m)}_{\text{estimate of } M_j - m_j} \cdot \underbrace{\frac{\varepsilon}{2} \cdot \frac{1}{2k(M - m)}}_{\text{length of } I_j} + \underbrace{\frac{\varepsilon}{2(b-a)}}_{\text{estimate of } M_j - m_j} \underbrace{(b-a)}_{\text{total length}} \\
 &= \frac{\varepsilon}{2} + \frac{\varepsilon}{2} \\
 &= \varepsilon.
 \end{aligned}$$

Therefore,

$$\int_a^b f = \int_a^b f \implies f \text{ is integrable.}$$

- (Proof of ②): Assume f is increasing. Given $\varepsilon > 0$. Consider an equal partition

$$P_n = \left\{ a = x_0, x_1 = x_0 + \frac{b-a}{n}, x_2, \dots, x_n = b \right\}.$$

Then, by equal partition and f is increasing, we have

$$U(f, P_n) = \sum_{j=1}^n f(x_j)(x_j - x_{j-1}) = \frac{b-a}{n} \sum_{j=1}^n f(x_j)$$

and

$$L(f, P_n) = \sum_{j=1}^n f(x_{j-1})(x_j - x_{j-1}) = \frac{b-a}{n} \sum_{j=1}^n f(x_{j-1}).$$

So,

$$\begin{aligned}
 U(f, P_n) - L(f, P_n) &= \frac{b-a}{n} \sum_{j=1}^n f(x_j) - f(x_{j-1}) \\
 &= \frac{b-a}{n} (f(x_n) - f(x_1)) \quad \text{[Intermediate terms cancel]} \\
 &= \frac{b-a}{n} (f(b) - f(a)).
 \end{aligned}$$

When $n \rightarrow \infty$, $U(f, P_n) - L(f, P_n) = \frac{b-a}{n} (f(b) - f(a)) \rightarrow 0$. Therefore,

$$U(f, P_n) - L(f, P_n) < \varepsilon \quad \text{for large } n \implies f \text{ is integrable.}$$

Q.E.D. ■

Remark 4.12 To prove a function f is integrable, it is sufficient to show that $\forall \varepsilon > 0, \exists$ partition P s.t.

$$U(f, P) = L(f, P) < \varepsilon.$$

Theorem 4.5.6 Rules of Integration

- $k \int_a^b f(x) dx = \int_a^b k f(x) dx$, k is a constant.
- $\int_a^b (f(x) \pm g(x)) dx = \int_a^b f(x) dx \pm \int_a^b g(x) dx$
- $\int_a^c f(x) dx = \int_a^b f(x) dx + \int_b^c f(x) dx$, for $a \leq b \leq c$.
- If $f \leq g$, then $\int_a^b f(x) dx \leq \int_a^b g(x) dx$.

In particular, $-|f| \leq f \leq |f|$, so

$$-\int_a^b |f| \leq \int_a^b f \leq \int_a^b |f|.$$

That is,

$$\left| \int_a^b f \right| \leq \int_a^b |f|.$$

Definition 4.5.7 (Antiderivative). Let $f(x) : [a, b] \rightarrow \mathbb{R}$. An *antiderivative* of f is a continuous function $F(x) : [a, b] \rightarrow \mathbb{R}$ s.t. $F'(x) = f(x) \quad \forall x \in (a, b)$.

Remark 4.13 (Antiderivative is not Unique) Suppose $F(x)$ is an antiderivative of $f(x)$. If G is another antiderivative, then

$$\frac{d}{dx}[G(x) - F(x)] = G'(x) - F'(x) = f(x) - f(x) = 0 \quad \forall x \in (a, b).$$

So, by MVT, $G(x) - F(x) = C$, where C is some constant, or

$$G(x) = F(x) + C.$$

Theorem 4.5.8 Fundamental Theorem of Calculus (FTC)

Let $f(x) : [a, b] \rightarrow \mathbb{R}$ be continuous. Then, f has an antiderivative F , and

$$\int_a^b f(x) dx = F(b) - F(a) \left[= F(x) \Big|_a^b \right].$$


Proof 3. Define $F(x)$ by

$$F(x) = \int_a^x f(t) dt$$

for $x \in [a, b]$.

Claim $F(x)$ is an antiderivative of $f(x)$.

Proof.



Fix $x \in (a, b)$. Let $h > 0$ s.t. $(x - h, x + h) \subset (a, b)$. Then,

$$\begin{aligned} \frac{F(x+h) - F(x)}{h} &= \frac{1}{h} \left(\int_a^{x+h} f(t) dt - \int_a^x f(t) dt \right) \\ &= \frac{1}{h} \left(\cancel{\int_a^x f(t) dt} + \int_x^{x+h} f(t) dt - \cancel{\int_a^x f(t) dt} \right) = \frac{1}{h} \int_x^{x+h} f(t) dt. \end{aligned}$$

Note that

$$f(x) = \frac{1}{h} \int_x^{x+h} \underbrace{f(x)}_{\text{constant w.r.t. } t} dt$$

So,

$$\begin{aligned} \frac{F(x+h) - F(x)}{h} - f(x) &= \frac{1}{h} \int_x^{x+h} f(t) dt - \frac{1}{h} \int_x^{x+h} f(x) dt \\ &= \frac{1}{h} \int_x^{x+h} f(t) - f(x) dt \end{aligned}$$

Given $\varepsilon > 0$, f is continuous at x . So, $\exists \delta > 0$ s.t.

$$|t - x| < \delta \implies |f(t) - f(x)| < \varepsilon.$$

Then, when $|h| < \delta$, we have

$$\begin{aligned} \left| \frac{F(x+h) - F(x)}{h} \right| &\leq \left| \frac{1}{h} \int_x^{x+h} f(t) - f(x) dt \right| \\ &\leq \frac{1}{|h|} \int_x^{x+h} |f(t) - f(x)| dt \\ &< \frac{1}{|h|} \int_x^{x+h} \varepsilon dt \\ &= \frac{1}{|h|} \cdot \varepsilon \cdot |h| \\ &= \varepsilon. \end{aligned}$$

So,

$$\lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = f(x) \quad \text{i.e., } F'(x) = f(x).$$

Furthermore, one can show that $F(x)$ is continuous on $[a, b]$. [As $F(x)$ is differentiable on (a, b) , it is continuous on (a, b) . We only need to check for the endpoints.] \square

Finally, by definition,

$$F(b) = \int_a^b f(t) \, dt \quad \text{and} \quad F(a) = \int_a^b f(t) \, dt = 0.$$

So,

$$\int_a^b f(t) \, dt = F(b) - F(a).$$

Q.E.D. \blacksquare

Remark 4.14 In FTC, the continuity assumption of $f(x)$ cannot be removed. More specifically, it cannot be replaced by integrability. For example,

$$f(x) = \begin{cases} 0, & 0 \leq x \leq 1 \\ 1, & 1 < x \leq 2. \end{cases}$$

f is integrable, and its antiderivative

$$F(x) = \int_0^x f(t) \, dt \quad \text{is well-defined.}$$

However, $F'(x) = f(x)$ for $1 < x \leq 2$. When $x = 1$, $F'(x)$ does not even exist.

5 Uniform Convergence

5.1 Definition of Convergence

Definition 5.1.1 (Pointwise Convergence). Given a sequence of functions $f_k(x) : A \subset M \rightarrow N$ for $k = 1, 2, \dots$. We say $f_k(x) \rightarrow f(x)$ *converges pointwise* on A if $\forall x \in A$, the sequence of points $\{f_k(x)\}$ converges to $f(x)$. That is, $\forall x, \forall \varepsilon > 0, \exists K \text{ s.t. } k \geq K \implies \rho(f_k(x), f(x)) < \varepsilon$.

Definition 5.1.2 (Uniform Convergence). $f_k(x) \rightarrow f(x)$ *converges uniformly* on A if $\forall \varepsilon > 0, \exists K \text{ s.t. } k \geq K \implies \rho(f_k(x), f(x)) < \varepsilon \quad \forall x \in A$. We write $f_k \rightarrow f$ UC on A .

Remark 5.1 For pointwise convergence, the choice of K depends both on ε and the point x . However, for uniform convergence, K only depends on ε but not specific point x .

Definition 5.1.3 (Convergence of Series). Assume N is a normed space. Suppose $g_k : A \subset M \rightarrow N$. Then, $\sum_{k=1}^{\infty} g_k(x)$ converges to $g(x)$ *pointwise* or *uniformly*. Using sequence of partial sums, we have

$$f_n(x) = \sum_{k=1}^n g_k(x).$$

Remark 5.2 UC is stronger: $UC \implies$ pointwise convergence.

However, pointwise convergence $\not\Rightarrow$ UC in general.


Example 5.1.4

Consider $A = [0, 1]$ and

$$f_k(x) = \begin{cases} 0 & \text{if } \frac{1}{k} \leq x \leq 1 \\ 1 & \text{if } 0 \leq x \leq \frac{1}{k} \end{cases}$$

Note that $f_k(x) \rightarrow f(x)$ pointwise, where

$$f(x) = \begin{cases} 0 & x > 0 \\ 1 & x = 0. \end{cases}$$



However, this convergence is not uniform: $\exists \varepsilon_0 > 0 \text{ s.t. } \forall K, \exists k \geq K \text{ s.t. } \rho(f_k(x), f(x)) > \varepsilon_0$ for some $x \in A$. For example, take $\varepsilon_0 = \frac{1}{2}$ and $0 < x < \frac{1}{k}$.

Theorem 5.1.5 Continuity of Uniform Limit

Let $f_k : A \subset M \rightarrow N$ be a sequence of continuous functions and $f_k \rightarrow f$ uniformly converges on A . Then, f is also continuous.

Proof 1. Fix $x_0 \in A$. Given $\varepsilon > 0$. By UC, $\exists K$ s.t. $\rho(f_K(x), f(x)) < \frac{\varepsilon}{3} \quad \forall x \in A$. Since f_K is continuous, $\exists \delta > 0$ s.t.

$$x \in A, d(x, x_0) < \delta \implies \rho(f_K(x), f(x_0)) < \frac{\varepsilon}{3}.$$

Therefore, by triangle inequality, we have

$$\begin{aligned} \rho(f(x), f(x_0)) &\leq \rho(f(x), f_K(x)) + \rho(f_K(x), f_K(x_0)) + \rho(f_K(x_0), f(x_0)) \\ &< \frac{\varepsilon}{3} + \frac{\varepsilon}{3} + \frac{\varepsilon}{3} \\ &= \varepsilon. \end{aligned}$$

So, f is continuous at x_0 .

Q.E.D. ■

Remark 5.3 This result can be used to show that a convergence is not uniform.

Example 5.1.6

- $f_n(x) = \frac{x^n}{1+x^n}$, with $A = [0, 2]$.

– Find pointwise limit

$$f_n(x) \rightarrow f(x) = \begin{cases} 0, & 0 \leq x < 1 \\ \frac{1}{2}, & x = 1 \\ 1, & 1 < x \leq 2. \end{cases}$$

– Determine uniform convergence:

The convergence is not uniform because f is not continuous.

- Geometric Series: *Counterexample to the converse of Theorem 5.1.5*

$$\sum_{k=0}^{\infty} x^k \quad \text{with } A = (-1, 1).$$

– Converge pointwise to $g(x) = \frac{1}{1-x}$.

Find partial sum:

$$S_n(x) = \sum_{k=0}^n x^k = \frac{1-x^{n+1}}{1-x}.$$

Since $x \in (-1, 1)$, as $n \rightarrow \infty$, $x^{n+1} \rightarrow 0$. So,

$$S_n(x) = \frac{1 - x^{n+1}}{1 - x} \xrightarrow{n \rightarrow \infty} \frac{1}{1 - x} \quad \text{for } x \in (-1, 1).$$

- Uniform convergence on subinterval $[-a, a]$ for any $0 < a < 1$.

Estimate the error term:

$$|S_n(x) - g(x)| = \frac{|x|^{n+1}}{|1 - x|}.$$

When $x \rightarrow 1$, $|S_n(x) - g(x)| \rightarrow \infty$ as $|1 - x| \rightarrow 0$. However, if we restrict $x \in [-a, a]$ for some $0 < a < 1$, then $|1 - x| \geq 1 - a$, and we have

$$\forall \varepsilon > 0, \quad \exists N \text{ s.t. } n \geq N \implies \frac{a^{n+1}}{1 - a} < \varepsilon.$$

$$\implies |S_n(x) - g(x)| \leq \frac{a^{n+1}}{1 - a} < \varepsilon \quad \forall x \in [-a, a].$$

- Convergence is not uniform on $(-1, 1)$.

Observe that for any fixed N , we have $\frac{|x|^{N+1}}{|1 - x|} \xrightarrow{x \rightarrow 1^-} \infty$. Therefore,

$$\exists x_0 < 1 \text{ s.t. } \frac{|x_0|^{N+1}}{|1 - x_0|} = |S_N(x_0) - g(x_0)| \geq 1 = \varepsilon_0.$$

Definition 5.1.7 (Uniformly Cauchy Sequence). A sequence of functions $f_k : A \subset M \rightarrow N$ is *uniformly Cauchy sequence* if $\forall \varepsilon > 0, \exists L > 0 \text{ s.t. } j, k \geq L \implies \rho(f_k(x), f_j(x)) < \varepsilon \quad \forall x \in A$.

Theorem 5.1.8 Cauchy Criterion

Let (N, ρ) be a *complete* metric space and $f_k : A \subset M \rightarrow N$ be a sequence of functions. Then, f_k converges uniformly on $A \iff \forall \varepsilon > 0, \exists L > 0 \text{ s.t.}$

$$j, k \geq L \implies \rho(f_k(x), f_j(x)) < \varepsilon \quad \forall x \in A.$$

Proof2. (\implies) Assume $f_k \rightarrow f$ uniformly. [WTS: f_k is uniformly Cauchy.]

$$\rho(f_k(x), f_j(x)) \leq \rho(f_k(x), f(x)) + \rho(f(x), f_j(x)). \quad \square$$

(\Leftarrow) Assume $\{f_k\}$ is uniformly Cauchy.

- Find the limit function (pointwise)

For each fixed $x \in A$, the sequence of points $\{f_k(x)\}$ is Cauchy in N . By completeness of N , $f_k(x)$ converges to some point in N . Denoted by $f(x)$.

• **Show $f_k(x) \rightarrow f(x)$ UC**

Given $\varepsilon > 0$, $\exists L_1$ s.t. $j, k \geq L_1 \implies \rho(f_k(x), f_j(x)) < \frac{\varepsilon}{2} \quad \forall x \in A$. Furthermore, as $f_k(x) \rightarrow f(x)$ pointwise, for any $x \in A$, $\exists L_x \geq L_1$ s.t. $j \geq L_x \implies \rho(f_j(x), f(x)) < \frac{\varepsilon}{2}$.

Now, let $K = L_1$. Then, when $k \geq K$ we have

$$\begin{aligned} \rho(f_k(x), f(x)) &\leq \rho(f_k(x), f_{L_x}(x)) + \rho(f_{L_x}(x), f(x)) \\ &< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} \\ &= \varepsilon \quad \forall x \in A. \end{aligned}$$

Just pick $j = L_x$, we have different intermediate term for different x 's.

Q.E.D. ■

Corollary 5.1.9 Weierstrass M Test: Let N be a complete normed space and $g_k : A \rightarrow N$ be a sequence of functions s.t. \exists constants M_k with

- $\|g_k(x)\| \leq M_k$ for all $x \in A$, and
- $\sum_{k=1}^{\infty} M_k$ converges.

Then, the series $\sum_{k=1}^{\infty} g_k(x)$ converge uniformly.

Proof 3. The sequence of partial sums $\{f_n(x)\}$ is uniformly Cauchy.

$$f_n(x) = \sum_{k=1}^n g_k(x).$$

Then, apply Cauchy criterion.

Q.E.D. ■

Example 5.1.10

- $\sum_{n=1}^{\infty} \frac{(\sin nx)^2}{n^2}, \quad A = \mathbb{R}.$

Set $g_n(x) = \frac{(\sin nx)^2}{n^2}$. Then, $|g_n(x)| \leq \frac{1}{n^2}$.

As $\sum_{n=1}^{\infty} \frac{1}{n^2}$ converges, by M test, $\sum_{n=1}^{\infty} \frac{(\sin nx)^2}{n^2}$ converges uniformly.

- $\sum_{n=0}^{\infty} \left(\frac{x^n}{n!}\right)^2 \rightarrow f(x)$ on \mathbb{R} pointwise

If we limit $A = [-a, a]$, then $\sum_{n=0}^{\infty} \left(\frac{x^n}{n!}\right)^2$ uniformly converges.

5.2 Integration and Differentiation of Series

Theorem 5.2.1

Suppose $f_n : [a, b] \rightarrow \mathbb{R}$ and integrable and $f_n \rightarrow f$ uniformly on $[a, b]$. Then, f is integrable, and

$$\lim_{n \rightarrow \infty} \int_a^b f_n(x) \, dx = \int_a^b \lim_{n \rightarrow \infty} f_n(x) \, dx = \int_a^b f(x) \, dx.$$


Proof 1. Assume f is integrable. Then,

$$\begin{aligned} \left| \int_a^b f_n(x) \, dx - \int_a^b f(x) \, dx \right| &\leq \int_a^b \underbrace{|f_n(x) - f(x)|}_{< \varepsilon \quad \forall x, \text{ by UC}} \, dx \\ &< \int_a^b \varepsilon \, dx = \varepsilon(b - a). \end{aligned}$$

Q.E.D. ■

Remark 5.4 One cannot replace uniform convergence by pointwise convergence.

Example 5.2.2



Define $f_k(x) : [0, 1] \rightarrow \mathbb{R}$ s.t.

$$\int_0^1 f_k(x) \, dx = 1.$$

Observe that $f_k(x) \xrightarrow{\text{pointwise}} f(x) \equiv 0 \quad \forall x \in [0, 1]$. So,

$$\int_0^1 f_k(x) \, dx \not\rightarrow \int_0^1 f(x) \, dx$$

Remark 5.5 The same result is not true for differentiation. One cannot simply replace integrable with differentiable. For example, consider

$$f_n(x) = \frac{x^{n+1}}{n+1} \quad \text{on } [0, 1] \implies f'_n(x) = x^n.$$

We have $f_n(x) \xrightarrow{UC} f(x) \equiv 0$. However,


$$\lim_{n \rightarrow \infty} f'_n(x) \neq \lim_{n \rightarrow \infty} f'(x).$$

Theorem 5.2.3

Let $f_n : (a, b) \rightarrow \mathbb{R}$ be differentiable, converging pointwise to $f(x) : (a, b) \rightarrow \mathbb{R}$. If $f'_n(x)$ are continuous and converges uniformly to a function g , then $f'(x) = g(x)$. i.e.,

$$\lim_{n \rightarrow \infty} \frac{d}{dx}(f_n(x)) = \frac{d}{dx} \left(\lim_{n \rightarrow \infty} f_n(x) \right) = \frac{d}{dx} f(x) = g(x).$$

Proof2.



Use Fundamental Theorem of Calculus,

$$\begin{aligned} f_n(x) &= f_n(x_0) + f_n(x) - f_n(x_0) \\ &= f_n(x_0) + \int_{x_0}^x f'_n(t) dt. \end{aligned}$$

When $n \rightarrow \infty$, for fixed $x \in A$,

$$f_n(x) \rightarrow f(x), \quad f_n(x_0) \rightarrow f(x_0), \quad \int_{x_0}^x f'_n(t) dt \rightarrow \int_{x_0}^x g(t) dt.$$

So,

$$\begin{aligned} f(x) &= f(x_0) + \int_{x_0}^x g(t) dt \\ \frac{d}{dx}(f(x)) &= \frac{d}{dX}(f(x_0)) + \frac{d}{dx} \int_{x_0}^x g(t) dt \\ \lim_{n \rightarrow \infty} f'_n(x) &= f'(x) = 0 + g(x) = g(x). \end{aligned}$$

Q.E.D. ■

Example 5.2.4 One cannot replace UC with pointwise convergence

$$f_n = \frac{nx^2}{1 + nx^2}, \quad -1 \leq x \leq 1 \implies f'_n(x) \xrightarrow{\text{pointwise}} g(x)$$

However, $f'_n(x) \neq g(x)$.

5.3 The Space of Continuous Functions

Notation 5.1. Let $A \subset M$ be a metric space and N is a normal vector space. Then

- $\mathcal{C} = \mathcal{C}(A, N) = \{f \mid f: A \rightarrow N \text{ continuous}\}$: the collection of all continuous functions $f: A \rightarrow N$
- $\mathcal{C}_b = \mathcal{C}(A, N) = \{f \in \mathcal{C} \mid f \text{ is bounded}\}$: the collection of all bounded continuous functions
($\exists M$ s.t. $\|f(x)\|_N \leq M \quad \forall x \in A$)

Example 5.3.2

$A = [0, 1] \subset \mathbb{R}, N = \mathbb{R}$. Then,

$$\mathcal{C}_b = \mathcal{C}, \quad \text{the set of all continuous functions on } [0, 1].$$

Remark 5.6

- \mathcal{C}_b and \mathcal{C} are vector spaces;
- Goal: Study \mathcal{C}_b as a normed vector spaces as \mathbb{R}^n .

Definition 5.3.3 (Norm on \mathcal{C}_b). Given $f \in \mathcal{C}_b$. Define $\|f\|$ as follows:

$$\|f\| = \sup \{\|f(x)\|_N \mid x \in A\}.$$

This is called the *maximum absolute value norm*.

Theorem 5.3.4

$\|\cdot\|$ defined in Definition 5.3.3 is a norm in \mathcal{C}_b . i.e.,

- Positive definiteness: $\|f\| \geq 0$ and $\|f\| = 0 \iff f = 0$;
- Scalar multiplicity: $\|\alpha f\| = |\alpha| \|f\| \quad \forall \alpha \in \mathbb{R}$
- Triangle inequality: $\|f + g\| \leq \|f\| + \|g\|$

Proof 1. (of ③) By definition, $\|f + g\| = \sup \{\|f(x) + g(x)\|_N \mid x \in A\}$. [WTS: $\|f\| + \|g\|$ is an upper bound.] Note that

$$\begin{aligned} \|f(x) + g(x)\|_N &\leq \|f(x)\|_N + \|g(x)\|_N && \text{[triangle inequality in } N\text{]} \\ &\leq \|f\| + \|g\| && \text{[definition]} \end{aligned}$$

So, $\|f + g\| \leq \|f\| + \|g\|$.

Q.E.D. ■

Definition 5.3.5 (Convergence in \mathcal{C}_b). $f_k \rightarrow f$ in \mathcal{C}_b means that $\|f_k - f\| \rightarrow 0$ as $k \rightarrow \infty$.

Theorem 5.3.6

$f_k \rightarrow f$ in \mathcal{C}_b (convergence in norm as vectors) $\iff f_k \rightarrow f$ uniformly on A (convergence in function)

Proof2. (\Rightarrow): Assume $\|f_k - f\| \rightarrow 0$. Then, $\forall \varepsilon > 0, \exists K$ s.t. $k \geq K \implies \|f_k - f\| \leq \varepsilon$. Thus, $\forall x \in A$, by definition of norm, for $k \geq K$,

$$\|f_k(x) - f(x)\|_N \leq \|f_k - f\| < \varepsilon.$$

So, $f_k(x) \rightarrow f(x)$ uniformly on A . \square

(\Leftarrow): Assume $f_k(x) \rightarrow f(x)$ uniformly on A . Then, $\forall \varepsilon > 0, \exists K$ s.t. $k \geq K \implies \|f_k(x) - f(x)\|_N < \varepsilon$. Then, ε is an upper bound. Note that

$$\|f_k - f\| = \sup \{\|f_k(x) - f(x)\|_N \mid x \in A\}$$

is a least upper bound. So,

$$\|f_k - f\| = \sup \{\|f_k(x) - f(x)\|_N \mid x \in A\} < \varepsilon$$

So, $\|f_k - f\| \rightarrow 0$ as $k \rightarrow \infty$.

Q.E.D. \blacksquare

Theorem 5.3.7 Completeness of \mathcal{C}_b

If N is complete, so is $\mathcal{C}_b(A, N)$.

Proof3. Let $\{f_k\}$ be a Cauchy sequence in \mathcal{C}_b . Then, $\forall \varepsilon > 0, \exists K$ s.t. $j, k \geq K \implies \|f_j - f_k\| < \varepsilon$. By definition, we have

$$\|f_j(x) - f_k(x)\|_N \leq \|f_j - f_k\| < \varepsilon \quad \forall x \in A.$$

So, $\{f_k(x)\}$ is a uniform Cauchy sequence on A . By Cauchy criterion,

$$f_k(x) \rightarrow f(x) \quad \text{uniformly on } A.$$

f is also continuous since UC preserves continuity. By Theorem 5.3.6, we have $f_k \rightarrow f$ in \mathcal{C}_b . So, \mathcal{C}_b is complete.

Q.E.D. \blacksquare

Remark 5.7 (Comparison Between \mathcal{C}_b and \mathbb{R}^n) Let $A \subset M$ be compact and $N = \mathbb{R}^n$.

Properties	\mathbb{R}^n	$\mathcal{C}_b(A, N = \mathbb{R}^n)$
Normed Space	✓	✓
Completeness	✓	✓
Finite Dimension	✓	✗
Compact Subset	<u>Heine-Borel:</u> $B \subset \mathbb{R}^n$ is compact $\iff B$ is closed and bounded	<u>Arzela-Ascoli:</u> $A \subset M$ compact. Then, $B \subset \mathcal{C}_b$ is compact $\iff B$ is closed, bounded, and equicontinuous in A

Definition 5.3.8 (Equicontinuous). A family of function \mathcal{B} is equicontinuous at a point $x \in A$ if $\forall \varepsilon > 0, \exists \delta > 0$ s.t. $y \in D(x, \delta) \cap A \implies \|f(x) - f(y)\|_N < \varepsilon \quad \forall f \in \mathcal{B}$.

Remark 5.8 δ is independent of $f \in \mathcal{B}$.

Example 5.3.9

- $\mathcal{B} = \{f \in \mathcal{C}_b(\mathbb{R}, \mathbb{R}) \mid f(x) > 0 \quad \forall x \in \mathbb{R}\}$.

– Is \mathcal{B} open? No.

Suppose $f \rightarrow 0$ as $x \in \infty$. Then, no matter how small we take the δ , some part of $D(f, \delta)$ will not be contained in \mathcal{B} .

– What is $\text{cl}(\mathcal{B})$?

$$\text{cl}(\mathcal{B}) = \{f \in \mathcal{C}_b(\mathbb{R}, \mathbb{R}) \mid f(x) \geq 0 \quad \forall x \in \mathbb{R}\}.$$

– What is $\text{int}(\mathcal{B})$?

$$\text{int}(\mathcal{B}) = \{f \in \mathcal{C}_b(\mathbb{R}, \mathbb{R}) \mid \inf(f(x)) > 0 \quad \forall x \in \mathbb{R}\}.$$

Think of $\inf(f(x)) > 0$ in this way: we need a buffer zone.

- $\mathcal{B} = \{f \in \mathcal{C}_b([0, 1], \mathbb{R}) \mid f(x) > 0 \quad \forall x \in [0, 1]\}$.

5.4 The Contraction Mapping Principle (CMP)

Theorem 5.4.1 CMP

Let (M, d) be a complete metric space, and $\Phi : M \rightarrow M$ be a map. Suppose \exists constant k s.t. $0 < k < 1$ s.t.

$$d(\Phi(x), \Phi(y)) \leq k \cdot d(x, y) \quad \forall x, y \in M.$$

Then,

- Φ has a unique fixed point in M . That is, $\exists !x^* \in M$ s.t. $\Phi(x^*) = x^*$.
- The fixed point can be constructed (or approximated) as follows:

Fix any point $x_0 \in M$. Let $x_1 = \Phi(x_0), x_2 = \Phi(x_1), \dots, x_{n+1} = \Phi(x_n), \dots$. Then,

$$\lim_{n \rightarrow \infty} x_n = x^*.$$

Remark 5.9 Φ is continuous. Further, Φ is Lipschitz $\implies \Phi$ is uniform continuous.

Proof 1. Fix $x_0 \in M$. Let $x_{n+1} = \Phi(x_n)$ for $n = 0, 1, 2, \dots$.

Claim $\{x_n\}$ is Cauchy.

Note that $\forall n \geq 1$,

$$\begin{aligned} d(x_n, x_{n+1}) &= d(\Phi(x_{n-1}), \Phi(x_n)) \leq kd(x_{n-1}, x_n) \\ &\leq k^2 d(x_{n-1}, x_{n-1}) \\ &\vdots \\ &\leq k^n d(x_0, x_1). \end{aligned}$$

Thus, $\forall p \geq 1$,

$$\begin{aligned} d(x_n, x_{n+p}) &\leq d(x_n, x_{n+1}) + d(x_{n+1}, x_{n+2}) + \cdots + d(x_{n+p-1}, x_{n+p}) \\ &\leq k^n d(x_0, x_1) + k^{n+1} d(x_0, x_1) + \cdots + k^{n+p-1} d(x_0, x_1) \\ &= \underbrace{(k^n + k^{n+1} + \cdots + k^{n+p-1})}_{\text{geometric series}} d(x_0, x_1) \rightarrow 0 \quad \text{as } n \rightarrow \infty. \end{aligned}$$

As the geometric series converges, $\{x_n\}$ is Cauchy.

Since M is complete, $x_n \rightarrow x^* \in M$.

Claim x^* is a fixed point.

Since Φ is continuous,

$$\lim_{n \rightarrow \infty} \Phi(x_n) = \Phi\left(\lim_{n \rightarrow \infty} x_n\right) = \Phi(x^*).$$

Meanwhile, $\Phi(x_n) = x_{n+1}$, so

$$\lim_{n \rightarrow \infty} \Phi(x_n) = \lim_{n \rightarrow \infty} x_{n+1} = x^*.$$

Hence, $x^* = \Phi(x^*)$, implying x^* is a fixed point.

Claim The fixed point is unique.

Let $y^* \in M$ be another fixed point. One can show

$$\begin{aligned} d(x^*, y^*) &\leq d(\Phi(x^*), \Phi(y^*)) && [x^*, y^* \text{ are fixed points}] \\ &\leq kd(x^*, y^*) && [\Phi \text{ is a contraction mapping}] \end{aligned}$$

$$\implies d(x^*, y^*) = 0.$$

Q.E.D. ■

Example 5.4.2 Application in ODE

Consider the following initial value problem (IVP):

$$\frac{dx}{dt} = f(t, x) \quad x(t_0) = x_0 \quad (\text{IVP})$$

- Basic Assumptions:

1. $f(t, x)$ is continuous in a neighborhood U of $(t_0, x_0) \in \mathbb{R}^2$
2. $f(t, x)$ is Lipschitz in x : \exists constant K s.t.

$$|f(t, x_1) - f(t, x_2)| \leq K|x_1 - x_2| \quad \forall (t_1, x_1), (t_1, x_2) \in U$$

- Apply CMP:

Theorem 5.4.3

If $f(t, x)$ is continuous in U and Lipschitz in x , then (IVP) has a unique solution $x = \varphi(t)$ in the neighborhood of t_0 : $(t_0 - \delta, t_0 + \delta)$. i.e.,

$$\varphi'(t) = f(t, \varphi(t)), \quad \varphi(t_0) = x_0.$$

- Solving (IVP) is equivalent to finding a function $\varphi(t)$ s.t.

$$\varphi'(t) = f(t, \varphi(t)).$$

Or, by integration:

$$\varphi'(t) = x_0 + \int_{t_0}^t f(s, \varphi(s)) \, ds \quad [x_0 \text{ comes from plugging in the initial condition}]$$

This is just a fixed point for the following map (an integral operator):

$$\Phi : g(t) \mapsto \Phi(g) = x_0 + \int_{t_0}^t f(s, g(s)) \, ds$$

Theorem 5.4.4

We need to construct an appropriate metric space $M \subset \mathcal{C}_b$ s.t. $\Phi : M \rightarrow M$ is a contraction mapping.

Algorithm 1: Iterative Method to Approximate the Solution to (IVP)

```

1 begin
2    $\varphi_0 \equiv x_0$ ;
3   for  $n = 0, 1, 2, \dots$  do
4      $\varphi_{n+1}(t) = \Phi(\varphi_n(t)) = x_0 + \int_{t_0}^t f(s, \varphi_n(s)) \, ds$ ;
```

Example 5.4.5

Consider the IVP: $f(t, x) = tx^2 + x^3$, $x(0) = 1$.

Let $\varphi_0(t) = 1$. Then,

$$\begin{aligned}
 \varphi_1(t) &= 1 + \int_0^t s\varphi_0(s)^2 + \varphi_0(s)^3 \, ds \\
 &= 1 + \int_0^t s + 1 \, ds \\
 &= 1 + \left[\frac{1}{2}s^2 + s \right]_0^t \\
 &= 1 + \frac{1}{2}t^2 + t \\
 \varphi_2(t) &= 1 + \int_0^t s\varphi_1(s)^2 + \varphi_1(s)^3 \, ds \\
 &= 1 + \int_0^t s \left(1 + \frac{1}{2}s^2 + s \right)^2 + \left(1 + \frac{1}{2}s^2 + s \right)^3 \, ds \\
 &\vdots
 \end{aligned}$$

6 Differential Mappings

6.1 Definition and Matrix Representation of a Differential

Definition 6.1.1 (Linear Transformation). A function $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is called a *linear transformation* if $\forall x, y \in \mathbb{R}^n$ and $\lambda \in \mathbb{R}$, we have

- $T(x + y) = T(x) + T(y)$
- $T(\lambda x) = \lambda T(x)$

These two properties can be combined and written equivalently as $T(ax + by) = aT(x) + bT(y) \quad \forall x, y \in \mathbb{R}^n$ and $\forall a, b \in \mathbb{R}$.

6.1.2 Matrix Representation of T .

Observation: Given $m \times n$ matrix A , define function $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ by $T(x) = A \cdot x$. Then, T is a linear transformation.

Proof 1.

$$T(ax + by) = A(ax + by) = A(ax) + A(by) = aAx + bAy = aT(x) + bT(y).$$

Q.E.D. ■

Example 6.1.3

Suppose $A = \begin{bmatrix} 2 & 3 \\ 1 & -1 \\ 0 & 4 \end{bmatrix}$. Then,

$$T(x) = A \cdot x = \begin{bmatrix} 2 & 3 \\ 1 & -1 \\ 0 & 4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2x_1 + 3x_2 \\ x_1 - x_2 \\ 4x_2 \end{bmatrix} \in \mathbb{R}^3.$$

Theorem 6.1.4 Fact

Every linear transformation T is determined by a matrix in such a way as above (via matrix multiplication).

Proof 2. Given $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ linear, we need to find a matrix A ($m \times n$) such that

$$T(x) = A \cdot x \quad \forall x \in \mathbb{R}^n.$$

To construct A , consider the standard basis for $\mathbb{R}^n : \{e_1, e_2, \dots, e_n\}$ and for $\mathbb{R}^m : \{e'_1, e'_2, \dots, e'_m\}$. Then,

$$T(e_j) = \sum_{i=1}^m a_{ij} e'_i, \quad \forall j = 1, 2, \dots, n.$$

Let $A = (a_{ij})_{m \times n}$.

Claim $T(x) = Ax \quad \forall x \in \mathbb{R}^n$.

In fact, let $x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \in \mathbb{R}^n$. Then, we can rewrite x as a linear combination of standard basis:

$$x = \sum_{j=1}^n x_j e_j.$$

So,

$$T(x) = x_1 T(e_1) + x_2 T(e_2) + \cdots + x_n T(e_n) = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = Ax. \quad [T \text{ is Linear}]$$

Q.E.D. ■

Remark 6.1 The collection of $\{\text{linear transformation } T : \mathbb{R}^n \rightarrow \mathbb{R}^m\}$ forms a 1-to-1 correspondence with the collection of $\{m \times n \text{ matrices } A\}$.

Theorem 6.1.5 Continuity of T

If $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is linear, then it is Lipschitz, and hence continuous.

Proof 3. Recall the definition of Lipschitz: $|f(x) - f(y)| \leq L \cdot |x - y|$.

Since $T(x) - T(y) = T(x - y)$, we only need to show that

$$\|T(x)\| \leq M \cdot \|x\| \text{ for some } M \in \mathbb{R}.$$

Let $x = \sum x_j e_j$. Then, $T(x) = \sum x_j T(e_j)$. So, $\|T(x)\| \leq \sum_j |x_j| \cdot \|T(e_j)\|$.

Recall that $\|x\| = \sqrt{\sum_j x_j^2}$. So, $|x_j| \leq \|x\|$. Hence,

$$\|T(x)\| \leq \sum_j \|x\| \cdot \|T(e_j)\| = \underbrace{\left(\sum_{j=1}^n \|T(e_j)\| \right)}_{M, \text{ independent of } x} \cdot \|x\| = M \cdot \|x\|$$

Q.E.D. ■

6.1.6 Derivative (Differential) as a Linear Transformation.

- Recall one variable case: Let $f : (a, b) \rightarrow \mathbb{R}$. Then, we can rewrite $f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0}$ as

$$\lim_{x \rightarrow x_0} \left[\frac{f(x) - f(x_0) - f'(x_0)(x - x_0)}{x - x_0} \right] = 0.$$

- Definition 6.1.7 (Generalization to Higher Dimensions).** A map $f : A \subset \mathbb{R}^n \rightarrow \mathbb{R}^m$ is said to be *differentiable* at $x_0 \in A$ if there is a linear map, denoted by $\mathbb{D}f(x_0) : \mathbb{R}^n \rightarrow \mathbb{R}^m$ with

$$\lim_{x \rightarrow x_0} \frac{\|f(x) - f(x_0) - \mathbb{D}f(x_0)(x - x_0)\|}{\|x - x_0\|} = 0 \quad (\star)$$

Remark 6.2 Interpretations of (\star) :

1. Rewrite (\star) : $\forall \varepsilon > 0, \exists \delta > 0$ s.t. $\forall x \in A$,

$$\|x - x_0\| > \delta \implies \|f(x) - f(x_0) - \mathbb{D}f(x_0)(x - x_0)\| < \varepsilon \|x - x_0\|.$$

2. $f(x) \approx f(x_0) + \underbrace{\mathbb{D}f(x_0) \cdot (x - x_0)}_{\text{linear map}}$ is called the affine map.

3. *Geometric Interpretation:* $z = f(x) : \mathbb{R}^n \rightarrow \mathbb{R}^1$. Then, $z - f(x_0) = \mathbb{D}f(x_0)(x - x_0)$ represents the tangent plane of the surface $z = f(x)$.

4. For $f : \mathbb{R}^1 \rightarrow \mathbb{R}^1$, $\mathbb{D}f(x)$ is the differential, representing a linear transformation, whereas $f'(x)$ or $\frac{df}{dx}$ is the derivative, which is just a number.

For example, $f(x) = x^2$. Then, $f'(x) = 2x$. However, $\mathbb{D}f(x)$ is a linear transformation $\mathbb{R}^1 \rightarrow \mathbb{R}^1$, defined as

$$\mathbb{D}f(x)(h) = 2xh, \quad \forall h \in \mathbb{R}^1.$$

- Uniqueness of Differential

Theorem 6.1.8

Let $A \subset \mathbb{R}^n$ be open and $f : A \rightarrow \mathbb{R}^m$ be differentiable at $x_0 \in A$. Then, the differential $\mathbb{D}f(x_0)$ is uniquely determined by f .

Proof 4. Let L_1 and L_2 be two linear transformations such that

$$\lim_{x \rightarrow x_0} \frac{\|f(x) - f(x_0) - L_1(x - x_0)\|}{\|x - x_0\|} = 0 = \lim_{x \rightarrow x_0} \frac{\|f(x) - f(x_0) - L_2(x - x_0)\|}{\|x - x_0\|}.$$

We need to show that $L_1 = L_2$. i.e., $L_1(h) = L_2(h) \quad \forall h \in \mathbb{R}^n$.

Fix any unit vector $e \in \mathbb{R}^n$. Let $x = x_0 + te$, where $t \in \mathbb{R}$ and $t \neq 0$ (This makes sense because A is

open by assumption). Then,

$$\begin{aligned}
 \|L_1(e) - L_2(e)\| &= \frac{\|L_1(te) - L_2(te)\|}{|t|} \\
 &= \frac{\|L_1(x - x_0) - L_2(x - x_0)\|}{\|x - x_0\|} \quad [|t| = \|x - x_0\|] \\
 &= \frac{\|L_1(x - x_0) - (f(x) - f(x_0)) + (f(x) - f(x_0)) - L_2(x - x_0)\|}{\|x - x_0\|} \\
 &\leq \frac{\|L_1(x - x_0) - (f(x) - f(x_0))\| + \|(f(x) - f(x_0)) - L_2(x - x_0)\|}{\|x - x_0\|} \\
 &= \frac{\|L_1(x - x_0) - (f(x) - f(x_0))\|}{\|x - x_0\|} + \frac{\|(f(x) - f(x_0)) - L_2(x - x_0)\|}{\|x - x_0\|}.
 \end{aligned}$$

Note that both parts $\rightarrow 0$ as $x \rightarrow x_0$. So, $\|L_1(e) - L_2(e)\| = 0$, and thus $L_1(e) = L_2(e) \quad \forall$ unit vector e . Using linear transformation, $L_1(h) = L_2(h) \quad \forall h \in \mathbb{R}^n$.

Q.E.D. ■

Remark 6.3 Theorem 6.1.8 is not true if A is not open. A trivial example would be when $A = \{x_0\}$, the set of just one point. Then, any linear map satisfies the differential definition. That is,

$$\lim_{\substack{x \rightarrow x_0 \\ x \in A}} \frac{\|f(x) - f(x_0) - T(x - x_0)\|}{\|x - x_0\|} = 0 \quad \forall \text{ linear map } T.$$

Or, equivalently, $\|f(x) - f(x_0) - T(x - x_0)\| < \varepsilon \|x - x_0\|$.

6.1.9 Matrix Representation of the Differential $\mathbb{D}f(x)$.

Question: Given f , how do we find the linear transformation $\mathbb{D}f(x)$?

Definition 6.1.10 (Partial Derivative). Write $f(x) = (f_1(x_1, \dots, x_n), f_2(x_1, \dots, x_n), \dots, f_m(x_1, \dots, x_n)) \in \mathbb{R}^m$. Then,

$$\frac{\partial f_j}{\partial x_i} = \lim_{h \rightarrow 0} \frac{f_j(x_1, \dots, x_{i-1}, x_i + h, x_{i+1}, \dots, x_n) - f_j(x_1, \dots, x_i, \dots, x_n)}{h}.$$

Theorem 6.1.11 Relation Between Differential $\mathbb{D}f(x)$ and Partial Derivatives

Suppose $A \subset \mathbb{R}^n$ is open and $f : A \rightarrow \mathbb{R}^m$ is differentiable at $x \in A$. Then, $\frac{\partial f_j}{\partial x_i}$ exists and the matrix of the linear map $\mathbb{D}f(x)$ is given by

$$\begin{bmatrix} \frac{\partial f_1}{\partial x_1} & \frac{\partial f_1}{\partial x_2} & \cdots & \frac{\partial f_1}{\partial x_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial f_m}{\partial x_1} & \frac{\partial f_m}{\partial x_2} & \cdots & \frac{\partial f_m}{\partial x_n} \end{bmatrix} \in \mathbb{R}^{m \times n},$$

and we denote this matrix as $J_f(x)$, the *Jacobian matrix* of f at x .

Proof 5. Denote the matrix of $\mathbb{D}f(x)$ by $B = (b_{ji})_{m \times n}$. We need to show $b_{ji} = \frac{\partial f_j}{\partial x_i}$.

Recall: $b_{ji} = j$ -th component of $\mathbb{D}f(x)(e_i) = \sum_{j=1}^m b_{ji}e'_j$. Fix i, j and let $y = x + he_i$, $h \in \mathbb{R}$. Then, by definition of differential,

$$\frac{\|f(y) - f(x) - \mathbb{D}f(x)(y - x)\|}{\|y - x\|} \rightarrow 0 \quad \text{as } y - x \rightarrow 0.$$

Taking the j -th component,

$$\frac{|f_j(x_1, \dots, x_i + h, \dots, x_n) - f_j(x_1, \dots, x_n) - b_{ji} \cdot h|}{|h|} \rightarrow 0 \quad \text{as } h \rightarrow 0.$$

So,

$$\lim_{h \rightarrow 0} \frac{f_j(x_1, \dots, x_i + h, \dots, x_n) - f_j(x_1, \dots, x_n)}{h} = b_{ji}.$$

Hence,

$$\frac{\partial f_j}{\partial x_i} = b_{ji} \quad \forall i, j.$$

So, $\mathbb{D}f(x)$ is determined by the Jacobian matrix $J_f(x)$.

Q.E.D. ■

Example 6.1.12

- $f(x, y, z) = (x^4y, xe^z) : \mathbb{R}^3 \rightarrow \mathbb{R}^2$.

$$J_f(x, y, z) = \begin{bmatrix} \frac{\partial f_1}{\partial x} & \frac{\partial f_1}{\partial y} & \frac{\partial f_1}{\partial z} \\ \frac{\partial f_2}{\partial x} & \frac{\partial f_2}{\partial y} & \frac{\partial f_2}{\partial z} \end{bmatrix} = \begin{bmatrix} 4x^3y & x^4 & 0 \\ e^z & 0 & xe^z \end{bmatrix}.$$

- Special Case: $m = 1$: $f : \mathbb{R}^n \rightarrow \mathbb{R}$. Then,

$$J_f(x) = \left[\frac{\partial f}{\partial x_1} \quad \frac{\partial f}{\partial x_2} \quad \cdots \quad \frac{\partial f}{\partial x_n} \right] \quad \text{is a } 1 \times n \text{ matrix.}$$

Definition 6.1.13 (Gradient). The *gradient*, $\text{grad } f$ or ∇f , is defined by the following vector:

$$\nabla f = \left(\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_n} \right).$$

Gradient points towards the direction of fastest growth.

- $f(x, y, z) = \frac{x \sin y}{z}$. Computing $\mathbb{D}f$ and ∇f .

Solution 6.

$$\mathbb{D}f(x) = J_f(x) = \begin{bmatrix} \frac{\sin y}{z} & \frac{x \cos y}{z} & -\frac{x \sin y}{z^2} \end{bmatrix}.$$

$$\nabla f(x) = \left(\frac{\sin y}{z}, \frac{x \cos y}{z}, -\frac{x \sin y}{z^2} \right).$$

□

Remark 6.4 (Relation Between $\mathbb{D}f(x)$ and ∇f) For any $h \in \mathbb{R}^n$, we have

$$\text{matrix multiplication} \leftarrow \mathbb{D}f(x)h = \langle \nabla f, h \rangle \rightarrow \text{inner product/dot product}$$

- Special Case: $n = 1$. Consider $x = c(t) : [a, b] \subset \mathbb{R} \rightarrow \mathbb{R}^m$. Then,

$$\mathbb{D}x(t) = c'(t) = (c'_1(t), c'_2(t), \dots, c'_m(t))$$

is the tangent vector.

6.2 Necessary and Sufficient Conditions for Differentiability

Definition 6.2.1 (Locally Lipschitz). f is *locally Lipschitz* at x_0 if $\forall x_0 \in A, \exists \delta > 0$ and M s.t.

$$\|x - x_0\| < \delta \implies \|f(x) - f(x_0)\| < M \cdot \|x - x_0\|.$$

Theorem 6.2.2 Necessary Condition for Differentiability I

Suppose $A \subset \mathbb{R}^n$ is open and $f : A \rightarrow \mathbb{R}^m$ is differentiable. Then, f is locally Lipschitz.

Remark 6.5 (Ideas to Prove this Theorem)

- Linear map $\mathbb{D}f(x)$ is Lipschitz;
- $f(x)$ can be approximated by $\mathbb{D}f(x_0)$ locally.

Proof 1. Fix $x_0 \in A$. By definition,

$$\lim_{x \rightarrow x_0} \frac{\|f(x) - f(x_0) - \mathbb{D}f(x_0)(x - x_0)\|}{\|x - x_0\|} = 0.$$

For $\varepsilon = 1, \exists \delta > 0$ s.t.

$$\|x - x_0\| < \delta \implies \|f(x) - f(x_0) - \mathbb{D}f(x_0)(x - x_0)\| \leq \varepsilon \cdot \|x - x_0\| = \|x - x_0\|.$$

By triangle inequality,

$$\|f(x) - f(x_0)\| \leq \|\mathbb{D}f(x_0)(x - x_0)\| + \|x - x_0\|.$$

Since $\mathbb{D}f(x_0)$ is Lipschitz, $\exists L$ s.t.

$$\|\mathbb{D}f(x_0)(x - x_0)\| \leq L \cdot \|x - x_0\|.$$

So, $\|x - x_0\| < \delta \implies$

$$\begin{aligned} \|f(x) - f(x_0)\| &\leq L \cdot \|x - x_0\| + \|x - x_0\| \\ &= \underbrace{(L + 1)}_M \cdot \|x - x_0\| \\ &= M \cdot \|x - x_0\|. \end{aligned}$$

Q.E.D. ■

Remark 6.6

- *Continuity is not sufficient to guarantee differentiability. For instance, $f(x) = |x|$.*

However, differentiability \implies continuity.

- *Derivative of a differentiable function may not be continuous. For example, consider the function*

$$f(x) = \begin{cases} x^2 \sin \frac{1}{x}, & x \neq 0 \\ 0, & x = 0. \end{cases}; f : \mathbb{R}^1 \rightarrow \mathbb{R}^1. \text{ Then, we have}$$

$$f'(0) = \lim_{x \rightarrow 0} \frac{f(x) - f(0)}{x} = \lim_{x \rightarrow 0} \frac{x^2 \sin \frac{1}{x}}{x} = \lim_{x \rightarrow 0} x \sin \frac{1}{x} = 0.$$

When $x \neq 0$,

$$f'(x) = 2x \sin \frac{1}{x} + x^2 \cos \frac{1}{x} \left(-\frac{1}{x^2}\right) = 1x \sin \frac{1}{x} - \cos \frac{1}{x}.$$

Conclusion: f is differentiable in \mathbb{R}^1 . However,

$$f'(x) = \begin{cases} 2x \sin \frac{1}{x} - \cos \frac{1}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$$

is not continuous at $x = 0$.

Theorem 6.2.3 Necessary Condition for Differentiability II

Suppose $f : A \subset \mathbb{R}^n \rightarrow \mathbb{R}^m$ is differentiable. Then, the partial derivatives, $\frac{\partial f_j}{\partial x_i}$, exists $\forall i, j$.

Example 6.2.4 The Converse is not True

The converse of Theorem 6.2.3 is, in general, not true. Here we will consider a counterexample.

Consider function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ given by

$$f(x, y) = \begin{cases} \frac{xy}{\sqrt{x^2 + y^2}}, & (x, y) \neq (0, 0) \\ 0, & (x, y) = (0, 0) \end{cases}$$

Claim f is continuous at $(0, 0)$.

In fact, we have $(a - b)^2 \geq 0 \implies a^2 - 2ab + b^2 \geq 0$. So,

$$ab \leq \frac{a^2 + b^2}{2} \quad a, b \in \mathbb{R}.$$

Then,

$$|xy| \leq \frac{1}{2}(a^2 + b^2) \implies \frac{xy}{\sqrt{x^2 + y^2}} \rightarrow 0 \quad \text{as } (x, y) \rightarrow (0, 0).$$

Claim $\frac{\partial f(0, 0)}{\partial x} = 0$ and $\frac{\partial f(0, 0)}{\partial y} = 0$.

$$\frac{\partial f(0, 0)}{\partial x} = \lim_{x \rightarrow 0} \frac{f(x, 0) - f(0, 0)}{x - 0} = \lim_{x \rightarrow 0} \frac{0 - 0}{x} = 0.$$

Claim f is not differentiable at $(0, 0)$.

If f were differentiable, the matrix of $\mathbb{D}f(0, 0)$ is given by

$$J_f(0, 0) = \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \right) = (0, 0).$$

However, note that

$$\frac{\|f(x, y) - f(0, 0) - \mathbb{D}f(x, y)\|}{\|(x, y) - (0, 0)\|} = \frac{\frac{|xy|}{\sqrt{x^2 + y^2}}}{\sqrt{x^2 + y^2}} = \frac{|xy|}{x^2 + y^2}.$$

Since $\frac{|xy|}{x^2 + y^2}$ does not $\rightarrow 0$ as $(x, y) \rightarrow (0, 0)$, f is not differentiable at $(0, 0)$.

Conclusion: Continuity + Existence of Partial Derivative $\frac{\partial f_j}{\partial x_i} \not\Rightarrow$ Differentiability.

Theorem 6.2.5 Sufficient Condition for Differentiability

Let $A \subset \mathbb{R}^n$ be open and $f = (f_1, \dots, f_m) : A \rightarrow \mathbb{R}^m$. If all the partials $\frac{\partial f_j}{\partial x_i}$ exist and continuous on A , then f is differentiable on A .

Proof 2. WTS: $\forall x \in A$,

$$\lim_{y \rightarrow x} \frac{\|f(y) - f(x) - J_f(x)(y - x)\|}{\|y - x\|} = 0.$$

It is sufficient to show that this is true for each component f_j of $f = (f_1, f_2, \dots, f_m)$. Thus, we may assume $m = 1$: $f : A \subset \mathbb{R}^n \rightarrow \mathbb{R}^1$. Then,

$$J_f(x) = \left(\frac{\partial f}{\partial x_1} \quad \frac{\partial f}{\partial x_2} \quad \cdots \quad \frac{\partial f}{\partial x_n} \right).$$

So,

$$J_f(y - x) = \sum_{i=1}^n \frac{\partial f}{\partial x_i} (y_i - x_i),$$

and

$$\begin{aligned} f(y) - f(x) &= f(y_1, y_2, \dots, y_n) - f(x_1, x_2, \dots, x_n) \\ &= f(y_1, y_2, \dots, y_n) - f(x_1, y_2, \dots, y_n) \\ &\quad + f(x_1, y_2, \dots, y_n) - f(x_1, x_2, \dots, y_n) \\ &\quad + f(x_1, x_2, \dots, y_n) - \cdots \quad \text{each time, we change one component} \\ &\quad + f(x_1, x_2, \dots, y_n) - f(x_1, x_2, \dots, x_n) \end{aligned}$$

By MVT,

$$f(y_1, y_2, \dots, y_n) - f(x_1, y_2, \dots, y_n) = \frac{\partial f}{\partial x_1}(u_1, y_2, \dots, y_n) \cdot (y_1 - x_1).$$

Applying MVT to other terms, we obtain

$$f(y) - f(x) = \frac{\partial f(z^{(1)})}{\partial x_1}(y_1 - x_1) + \frac{\partial f(z^{(2)})}{\partial x_2}(y_2 - x_2) + \cdots + \frac{\partial f(z^{(n)})}{\partial x_n}(y_n - x_n).$$

Thus,

$$\begin{aligned} \|f(y) - f(x) - J_f(x)(y - x)\| &= \left\| \sum_{i=1}^n \frac{\partial f(z^{(i)})}{\partial x_i} (y_i - x_i) - \sum_{i=1}^n \frac{\partial f(x)}{\partial x_i} (y_i - x_i) \right\| \\ &\leq \sum_{i=1}^n \left| \frac{\partial f(z^{(i)})}{\partial x_i} - \frac{\partial f(x)}{\partial x_i} \right| \cdot \|y - x\| \end{aligned}$$

Triangle Inequality:
 $|y_i - x_i| \leq \|y - x\|$

By continuity of partial derivative, $\forall \varepsilon > 0, \exists \delta > 0$ s.t.

$$\|y - x\| < \delta \implies \sum_{i=1}^n \left| \frac{\partial f(z^{(i)})}{\partial x_i} - \frac{\partial f(x)}{\partial x_i} \right| < \varepsilon$$

Hence,

$$\|f(y) - f(x) - J_f(x)(y - x)\| < \varepsilon\|y - x\|.$$

Q.E.D. ■

Definition 6.2.6 (Directional Derivative). Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ and $e \in \mathbb{R}^n$ be a unit vector. The directional derivative of f at x_0 in the direction e is given by

$$D_e f(x_0) = \left. \frac{d}{dt} f(x_0 + te) \right|_{t=0} = \lim_{t \rightarrow 0} \frac{f(x_0 + te) - f(x_0)}{t}.$$

Claim 6.2.7 If f is differentiable at x_0 , then $D_e f(x_0) = \mathbb{D}f(x_0) \cdot e$

Proof 3.

$$\begin{aligned} \lim_{t \rightarrow 0} \frac{\|f(x_0 + te) - f(x_0) - \mathbb{D}f(x_0)(te)\|}{\|te\|} &= 0 \\ \lim_{t \rightarrow 0} \frac{f(x_0 + te) - f(x_0)}{t} &= \mathbb{D}f(x_0)(e) \\ D_e f(x_0) &= \mathbb{D}f(x_0)(e). \end{aligned}$$

Q.E.D. ■

Remark 6.7 Existence of directional derivatives $\not\Rightarrow$ differentiability

Example 6.2.8

Continuity of f + Existence of directional derivative $\not\Rightarrow$ differentiability.

Consider function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ given by

$$f(x, y) = \begin{cases} \frac{xy}{\sqrt{x^2 + y^2}}, & (x, y) \neq (0, 0) \\ 0, & (x, y) = (0, 0). \end{cases}$$

Claim $D_e f(0, 0)$ exists for any direction $e \in \mathbb{R}^2$.

$$\lim_{t \rightarrow 0} \frac{f((0, 0) + te) - f(0, 0)}{t} \text{ exists } \forall e \in \mathbb{R}^2.$$

Definition 6.2.9 (Tangent Line/Plane).

- The *tangent line* to the curve $y = f(x)$ at x_0 is given by

$$y = f(x_0) + f'(x_0)(x - x_0).$$

- The *tangent plane* to the surface $z = f(x)$ at x_0 is given by

$$z = f(x_0) + \mathbb{D}f(x_0)(x - x_0).$$

Example 6.2.10

Find the tangent plane at $(1, 2)$ to the surface $z = x^2 + y^2$.

Solution 4.

$$J_f(x) = \left(\frac{\partial f}{\partial x} \quad \frac{\partial f}{\partial y} \right) = (2x \quad 2y).$$

The tangent plane is given by

$$\begin{aligned} z &= f(1, 2) + \mathbb{D}f(1, 2)((x, y) - (1, 2)) \\ &= 1^2 + 2^2 + \begin{bmatrix} 2x & 2y \end{bmatrix} \Big|_{(x,y)=(1,2)} \begin{bmatrix} x-1 \\ y-2 \end{bmatrix} \\ &= 5 + \begin{bmatrix} 2 & 4 \end{bmatrix} \begin{bmatrix} x-1 \\ y-2 \end{bmatrix} \\ z &= 5 + 2(x-1) + 4(y-2). \end{aligned}$$

□

Summary III: Relations among Properties of $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$

Differentiability

\implies Existence of Directional Derivative

\implies Existence of Partial Derivative (moving in direction of the basis)

+ **Theorem 6.2.5** \implies Differentiability

\nRightarrow Continuity

\implies Continuity

\nRightarrow Existence of Partial Derivative

6.3 Differentiation Rules

6.3.1 Chain Rule

Recall the one variable case: $h = g(u)$, $u = f(x)$. Then,

$$h = f \circ g(x) = g(f(x)),$$

and

$$\frac{dh}{dx} = \frac{dh}{du} \cdot \frac{du}{dx} = g'(f(x)) \cdot f'(x).$$

Theorem 6.3.1 General Case Chain Rule

Let $f : A \subset \mathbb{R}^n \rightarrow \mathbb{R}^m$ and $g : B \rightarrow \mathbb{R}^p$ be differentiable with $f(A) \subset B$. Then, the composite $g \circ f : A \rightarrow \mathbb{R}^p$ is differentiable, and

$$\mathbb{D}(g \circ f)(x) = \mathbb{D}g(f(x)) \circ \mathbb{D}f(x),$$

a composition of linear mappings.

In matrix notation, define $h = g(u)$ and $u = f(x)$. Then, $h = g \circ f(x) = g(f(x))$, and

$$J_h(x) = J_g(f(x)) \cdot J_f(x) \quad \text{product of matrices}$$

$$= \begin{bmatrix} \frac{\partial g_1}{\partial u_1} & \cdots & \frac{\partial g_1}{\partial u_m} \\ \vdots & \ddots & \vdots \\ \frac{\partial g_p}{\partial u_1} & \cdots & \frac{\partial g_p}{\partial u_m} \end{bmatrix} \cdot \begin{bmatrix} \frac{\partial f_1}{\partial x_1} & \cdots & \frac{\partial f_1}{\partial x_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_m}{\partial x_1} & \cdots & \frac{\partial f_m}{\partial x_n} \end{bmatrix}$$

Proof 1. (Sketch). We need to show: for fixed $x \in A \subset \mathbb{R}^n$,

$$\lim_{y \rightarrow x} \frac{\|h(y) - h(x) - \mathbb{D}h(x)(y - x)\|}{\|y - x\|} = 0,$$

or

$$\lim_{y \rightarrow x} \frac{\|g(f(y)) - g(f(x)) - \mathbb{D}g(f(x))[\mathbb{D}f(x)(y - x)]\|}{\|y - x\|} = 0.$$

Work with the numerator:

$$\begin{aligned} \text{numerator} &= \|g(f(y)) - g(f(x)) - \mathbb{D}g(f(x))(f(y) - f(x)) \\ &\quad + \mathbb{D}g(f(x))(f(y) - f(x)) - \mathbb{D}g(f(x))[\mathbb{D}f(x)(y - x)]\| \\ &\leq \|g(f(y)) - g(f(x)) - \mathbb{D}g(f(x))[\mathbb{D}f(x)(y - x)]\| \quad \text{triangle inequality} \\ &\quad + \|\mathbb{D}g(f(x))(f(y) - f(x)) - \mathbb{D}g(f(x))[\mathbb{D}f(x)(y - x)]\| \\ &\leq \varepsilon_1 \|f(y) - f(x)\| + \|\mathbb{D}g(f(x))\| \cdot \|f(y) - f(x) - \mathbb{D}f(x)(y - x)\| \\ &\quad (\varepsilon_1 : g \text{ is differentiable; } \mathbb{D}g(f(x)) : \text{common factor}) \\ &\leq \varepsilon_1 \cdot L \|y - x\| + M \cdot \varepsilon_2 \|y - x\| \\ &\quad (L : \text{local Lipschitz; } M : \text{differential bounded; } \varepsilon_2 : f \text{ is differentiable}) \\ &= (L\varepsilon_1 + M\varepsilon_2) \cdot \|y - x\|. \end{aligned}$$

Therefore,

$$\lim_{y \rightarrow x} \frac{\text{numerator}}{\|y - x\|} = \lim_{y \rightarrow x} \frac{(L\varepsilon_1 + M\varepsilon_2)\|y - x\|}{\|y - x\|} = \lim_{y \rightarrow x} L\varepsilon_1 + M\varepsilon_2 = 0.$$

Q.E.D. ■

Example 6.3.2

- Change of Variable

$$(x, y, z) \longleftrightarrow (r, \theta, z) : \begin{cases} x = r \cos \theta \\ y = r \sin \theta \\ z = z \end{cases} \quad (\text{cylindrical coordinate})$$

Let $h(r, \theta, z) = f(x, y, z) = f(x(r, \theta, z), y(r, \theta, z), z(r, \theta, z))$. Then,

$$\mathbb{D}h = \frac{\partial h}{\partial(r, \theta, z)} = \frac{\partial f}{\partial(x, y, z)} \cdot \frac{\partial(x, y, z)}{\partial(r, \theta, z)} = J_f \cdot \begin{bmatrix} \cos \theta & -r \sin \theta & 0 \\ \sin \theta & r \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

- Consider composition of the maps $[0, 1] \xrightarrow{\gamma} \mathbb{R}^n \xrightarrow{f} \mathbb{R}$. Then, $h(t) = f(\gamma(t))$. By chain rule,

$$\begin{aligned} h'(t) &= \mathbb{D}f \circ \mathbb{D}\gamma = \begin{pmatrix} \frac{\partial f}{\partial x_1} & \frac{\partial f}{\partial x_2} & \cdots & \frac{\partial f}{\partial x_n} \end{pmatrix} \begin{pmatrix} x'_1(t) \\ x'_2(t) \\ \vdots \\ x'_n(t) \end{pmatrix} \\ &= \sum_{i=1}^n \frac{\partial f}{\partial x_i} x'_i(t) = \langle \nabla f, \gamma'(t) \rangle. \end{aligned}$$

6.3.2 Other Differentiation Rules**Theorem 6.3.3 Product Rule**

Let $f : A \subset \mathbb{R}^n \rightarrow \mathbb{R}^m$ and $g : A \rightarrow \mathbb{R}$ be differentiable. Then, the product $gf : A \rightarrow \mathbb{R}^m$ is differentiable, and

$$\mathbb{D}(gf) = g(\mathbb{D}f) + (\mathbb{D}g)f.$$

More precisely, for each $x \in A$ and $h \in \mathbb{R}^n$,

$$\mathbb{D}(gf) \cdot h = \underbrace{g(x)}_{\text{scalar}} \cdot \underbrace{\mathbb{D}f(x)(h)}_{\in \mathbb{R}^m} + \underbrace{\mathbb{D}g(x)(h)}_{\text{scalar}} \cdot \underbrace{f(x)}_{\in \mathbb{R}^m}.$$

In particular,

$$\frac{\partial g(f_j)}{\partial x_i} = g \cdot \frac{\partial f_j}{\partial x_i} + \frac{\partial g}{\partial x_i} \cdot f_j.$$

Theorem 6.3.4 Other Differentiation Rules

$$\mathbb{D}(f + g) = \mathbb{D}f + \mathbb{D}g$$

$$\mathbb{D}(\lambda f) = \lambda \mathbb{D}f$$

$$\mathbb{D}\left(\frac{f}{g}\right) = \frac{g\mathbb{D}f - (\mathbb{D}g)f}{g^2} \quad \left(\text{derived from product rule: } \frac{f}{g} = f \cdot \frac{1}{g}\right)$$

6.4 Geometric Interpretation of Gradient

Let $f : A \subset \mathbb{R}^n \rightarrow \mathbb{R}$ be differentiable.

Definition 6.4.1 $(\mathbb{D}f(x), \nabla f(x), D_e f(x))$.

- Differential of f : a matrix/linear transformation

$$\mathbb{D}f(x) = \begin{bmatrix} \frac{\partial f}{\partial x_1} & \frac{\partial f}{\partial x_2} & \cdots & \frac{\partial f}{\partial x_n} \end{bmatrix}$$

- Gradient of f : a vector

$$\nabla f(x) = \left(\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_n} \right).$$


- Directional derivative of f in the direction e :

$$D_e f(x) = \mathbb{D}f(x)e = \langle \nabla f(x), e \rangle.$$

Geometric meaning of $D_e f(x)$: Rate of change in the direction of e .

6.4.2 Geometric Meaning of Gradient.

Claim 6.4.3 ∇f is perpendicular to the level surface S defined by $f(x) = \text{constant}$.



Proof 1. Fix any curve $\gamma(t)$ on S : $\gamma : [a, b] \rightarrow S$. Then, $f(\gamma(t)) = c$. By chain rule,

$$\mathbb{D}f(\gamma(t)) \cdot \gamma'(t) = 0 \implies \langle \nabla f(x_0), \gamma'(x_0) \rangle = 0.$$

So, $\nabla f(x_0) \perp \gamma'(x_0)$. That is, $\nabla f \perp \text{curve } \gamma \text{ on } S \implies \nabla f \perp S$.

Q.E.D. ■

Corollary 6.4.4 Tangent Plane: The tangent plane at x_0 of the level surface is given by

$$\langle \nabla f(x_0), x - x_0 \rangle = 0.$$

Example 6.4.5

Find the tangent plane at $(1, 0, 1)$ to the surface $x^2 - y^2 + xyz = 1$.

Claim 6.4.6 The direction of ∇f is the direction in which f has the greatest rate of change, which is given by $\|\nabla f\|$.

Proof2. Fix a direction $e \in \mathbb{R}^n$. Then, the rate of change in direction e is given by

$$D_e f(x_0) = \langle \nabla f, e \rangle = \|\nabla f\| \|e\| \cos \theta,$$

where θ is the angle between $\nabla f(x_0)$ and e . Then, the rate of change is maximized when $\cos \theta = 1$. So, $\theta = 0$. That is, e is in the direction of ∇f .

Q.E.D. ■

6.5 Mean Value Theorem (MVT)

Theorem 6.5.1 MVT in 1-D


Let $f : [a, b] \rightarrow \mathbb{R}^1$ be continuous and differentiable on (a, b) . Then, $\exists c \in (a, b)$ s.t.

$$f'(c) = \frac{f(b) - f(a)}{b - a} \quad \text{or} \quad f(b) - f(a) = f'(c)(b - a).$$

Theorem 6.5.2 MVT in Higher Dimension

Let $f : A \subset \mathbb{R}^n \rightarrow \mathbb{R}$ be differentiable on an open set A . Then, for any pair of points $x, y \in A$ s.t. the line segment $[x, y]$ joining x and y is contained in A , \exists a point $c \in [x, y]$ s.t.

$$f(y) - f(x) = \mathbb{D}f(c)(y - x).$$



Proof 1. Let $g(t) = (1 - t)x + ty$ for $0 \leq t \leq 1$ and

$$h(t) = f \circ g(t) = f((1 - t)x + ty) : [0, 1] \rightarrow \mathbb{R}.$$

Apply Theorem 6.5.1 to h , we know $\exists t_0 \in (0, 1)$ s.t.

$$\begin{aligned} h(1) - h(0) &= h'(t_0)(1 - 0) \\ f(y) - f(x) &= \mathbb{D}f(g(t_0)) \cdot g'(t_0) && [\text{Chain Rule}] \\ &= \mathbb{D}f(g(t_0)) \cdot (y - x). \end{aligned}$$

Denote $g(t_0) = c \in [x, y]$. Then,

$$f(y) - f(x) = \mathbb{D}f(c)(y - x).$$

Q.E.D. ■

Definition 6.5.3 (Convex Set). A set $A \subset \mathbb{R}^n$ is *convex* if $\forall x, y \in A$, $[x, y] \subset A$.

Corollary 6.5.4 : Let $A \subset \mathbb{R}^n$ be open and convex, and $f : A \rightarrow \mathbb{R}^m$ differentiable. If $\mathbb{D}f \equiv 0$, then f is constant in A .

Proof 2. (Sketch)

Apply MVT to each component of $f = (f_1, f_2, \dots, f_m)$.

Q.E.D. ■

6.6 Taylor's Theorem & Higher Order Differentials

6.6.1 One Dimensional Case

Theorem 6.6.1 Taylor's Formula

Let $f : (a, b) \rightarrow \mathbb{R}$ be one of class C^r (i.e., $f'(x), f''(x), \dots, f^{(r)}(x)$ are continuous). Then, for any $x_0, x \in (a, b)$, we have

$$f(x) = \underbrace{f(x_0) + f'(x_0)(x - x_0) + \frac{f''(x_0)}{2!}(x - x_0)^2 + \dots + \frac{f^{(r-1)}(x_0)}{(r-1)!}(x - x_0)^{r-1}}_{\text{Taylor's polynomial of degree } r-1} + \overbrace{R_{r-1}(x_0)}^{\text{Remainder}},$$

where R_{r-1} is the remainder at x_0 and can be written as

$$R_{r-1}(x_0) = \frac{f^{(r)}(c)}{r!}(x - x_0)^r \quad \text{for some } c \text{ between } x \text{ and } x_0.$$

Remark 6.8 (Key Idea to Prove) Use integration by parts in a reversed way multiple times.

Proof 1. Write $h = x - x_0$. Then, by Fundamental Theorem of Calculus,

$$f(x) - f(x_0) = f(x_0 + h) - f(x_0) = \int_0^1 \frac{d}{dt} f(x_0 + th) dt.$$

Now, apply integration by parts. Taking $u = \frac{d}{dt}f(x_0 + th) = f'(x_0 + th)h$ and $dv = dt \implies v = t - 1$, we have

$$\begin{aligned} f(x) - f(x_0) &= \int_0^1 u \, dv \\ &= uv \Big|_0^1 - \int_0^1 v \, du \\ &= -(-1)f'(x_0)h - \int_0^1 (t-1)f''(x_0 + th)h^2 \, dt \\ &= f'(x_0)h - \int_0^1 f''(x_0 + th)h^2(t-1) \, dt. \end{aligned}$$

Apply integration by parts again with

$$u = f''(x_0 + th)h^2 \quad \text{and} \quad dv = (t-1) \, dt \implies v = \frac{1}{2}(t-1)^2.$$

Then, we obtain

$$\begin{aligned} \int_0^1 f''(x_0 + th)h^2(t-1) \, dt &= f''(x_0 + th)h^2 \frac{1}{2}(t-1)^2 \Big|_0^1 - \int_0^1 \frac{1}{2}(t-1)^2 f'''(x_0 + th)h^3 \, dt \\ &= \frac{f''(x_0)}{2}h^2 + \int_0^1 f^{(3)}(x_0 + th)h^3 \cdot \frac{1}{2}(t-1)^2 \, dt. \end{aligned}$$

So,

$$f(x) - f(x_0) = f'(x_0)h + \frac{f''(x_0)}{2}h^2 + \int_0^1 f^{(3)}(x_0 + th)h^3 \cdot \frac{1}{2}(t-1)^2 \, dt.$$

By induction, we obtain that

$$\begin{aligned} f(x) - f(x_0) &= \overbrace{f'(x_0)h + \frac{f''(x_0)}{2}h^2 + \frac{f'''(x_0)}{3!}h^3 + \cdots + \frac{f^{(r-1)}(x_0)}{(r-1)!}h^{r-1}}^{\text{Taylor's polynomial}} \\ &\quad + \underbrace{(-1)^{r-1} \int_0^1 f^{(r)}(x_0 + th)h^r \frac{(t-1)^{r-1}}{(r-1)!} \, dt}_{\text{Remainder}} \end{aligned}$$

Lemma 6.6.2 2nd MVT for Integral: If $g \geq 0$, then $\int_a^b f(x)g(x) \, dx = f(\lambda) \int_a^b g(x) \, dx$.

Apply 2nd MVT to the remainder, we have

$$\begin{aligned} R_{r-1} &= (-1)^{r-1} f^{(r)}(x_0 + t_0 h)h^r \int_0^1 \frac{(t-1)^{r-1}}{(r-1)!} \, dt \\ &= f^{(r)}(x_0 + t_0 h)h^r \cdot \frac{1}{r} && \left[(-1)^{r-1} \text{ is absorbed when evaluating the integral} \right] \\ &= \frac{f^{(r)}(c)}{r!}h^r && \left[\text{Denote } c=x_0+t_0h, \text{ a point between } x_0 \text{ and } x \right] \end{aligned}$$

Combining everything, we get exactly what we have claimed.

Q.E.D. ■

Summary IV: Taylor's Formula & Taylor's Approximation

- Taylor's Formula:

$$f(x) = P_{r-1}(x) + R_{r-1}.$$

- Taylor's Approximation:

$$f(x) \approx P_{r-1}(x).$$

6.6.2 Taylor Series

Definition 6.6.3 (Taylor Series). Let $f \in C^\infty$. Then, the *Taylor series* is defined as

$$\sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n = f(x_0) + f'(x_0)(x - x_0) + \frac{f''(x_0)}{2}(x - x_0)^2 + \cdots.$$

Definition 6.6.4 (Real Analytic). f is (real) *analytic* at x_0 if its Taylor series converges to $f(x)$ in a neighborhood of x_0 . i.e.,

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n, \quad |x - x_0| < R.$$

Corollary 6.6.5 : If $f \in C^\infty(\mathbb{R})$ and for each interval $[a, b]$, \exists constant M s.t.

$$|f^{(n)}(x)| \leq M^n \quad \forall n \text{ and } x \in [a, b],$$

then, f is real analytic at each point x_0 and it has Taylor series representation. Namely,

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n, \quad |x - x_0| < \infty.$$

Proof 2. Fix $x_0 \in \mathbb{R}$. For any $x \in \mathbb{R}$, choose $b > 0$ s.t. $x_0, x \in [-b, b]$. By Taylor's Formula,

$$f(x) = \underbrace{P_{n-1}(x)}_{\text{partial sum of the series}} + R_{n-1}.$$

Recall:

$$R_{n-1} = \frac{f^{(n)}(c)}{n!} (x - x_0)^n \quad \text{for some } c.$$

Then,

$$|R_{n-1}| \leq \frac{M^n}{n!} |x - x_0|^n \quad \forall x \in [-b, b].$$

Since the series $\sum_{n=0}^{\infty} \frac{M^n}{n!} |x - x_0|^n$ converges by ratio test, its n -th term,

$$\frac{M^n}{n!} |x - x_0|^n \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Hence, $R_{n-1} \rightarrow 0$ as $n \rightarrow \infty$. Then, $P_{n-1}(x) \rightarrow f(x)$ as $n \rightarrow \infty$.

Q.E.D. ■

Example 6.6.6

- e^x and $\sin x$ are real analytic in \mathbb{R} . Find Taylor series at $x_0 = 0$:

Solution 3.

$$e^x = \sum_{n=0}^{\infty} \frac{1}{n!} x^n, \quad |x - x_0| < \infty.$$

□

- Is every \mathcal{C}^∞ real analytic? No.

Counterexample 6.7. Consider the function $f(x)$:

$$f(x) = \begin{cases} 0, & x = 0 \\ e^{-1/x^2}, & x \neq 0. \end{cases}$$

Claim $f(x) \in \mathcal{C}^\infty$.

Proof. At $x = 0$,

$$f'(0) = \lim_{x \rightarrow 0} \frac{f(x) - f(0)}{x - 0} = \lim_{x \rightarrow 0} \frac{e^{-1/x^2}}{x} = 0 \quad (\text{by L.H.})$$

At $x \neq 0$,

$$f'(x) = e^{-1/x^2} \left(\frac{2}{x^3} \right) = \frac{2/x^3}{e^{1/x^2}} \rightarrow 0 \quad \text{as } x \rightarrow 0 \quad (\text{by L.H.})$$

So, $f'(x)$ is continuous at $x = 0$, and

$$f'(x) = \begin{cases} 0, & x = 0 \\ e^{-1/x^2} \left(\frac{2}{x^3} \right) & x \neq 0. \end{cases}$$

By induction, one can show that

- $f^{(n)}(0) = 0 \quad \forall n$
- $f^{(n)}(x) \rightarrow 0$ as $x \rightarrow 0$.

So, $f^{(n)}(x)$ is continuous at $x = 0$. So, $f \in \mathcal{C}^\infty$. \square

Claim $f(x)$ is not real analytic at $x = 0$.

Proof. Taylor series:

$$\sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} (x)^n = 0.$$

So, the Taylor series does not converge to $f(x)$ on any neighborhood of $x = 0$. \square

6.6.3 Higher Dimensional Case

Observation: Let $f : A \subset \mathbb{R}^n \rightarrow \mathbb{R}$.

- Differential: $\mathbb{D}f(x)$ is a linear transformation $\mathbb{R}^n \rightarrow \mathbb{R}$.
- Let $g(x) = \mathbb{D}f(x)$. Then, $g : A \subset \mathbb{R}^n \rightarrow \mathbf{L}(\mathbb{R}^n, \mathbb{R}) \approx \mathbb{R}^n$, where $\mathbf{L}(M, N)$ is the space of linear transformation from M to N .
- $\mathbb{D}g(x)$ is a linear transformation $\mathbb{R}^n \rightarrow \mathbb{R}^n$ or $\mathbf{L}(\mathbb{R}^n, \mathbb{R})$.

Notation 6.8. Higher Order Differential The second order differential of f at x is denoted as

$$\mathbb{D}^2 f(x) = \mathbb{D}g(x) = \mathbb{D}(\mathbb{D}f(x)).$$

Definition 6.6.9 (Bilinear Maps). Given f and $x \in A$. Define a *bilinear map*, $\mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ by

$$\mathbb{D}^2 f(x)(u, v) = \left[\mathbb{D}^2 f(x)(u) \right] (v),$$

where $u, v \in \mathbb{R}^n$ and $\mathbb{D}^2 f(x)(u) \in \mathbf{L}(\mathbb{R}^n, \mathbb{R})$. In matrix notation,

$$uBv^\top,$$

where u is $1 \times n$, B is $n \times n$, and v^\top is $n \times 1$.

Definition 6.6.10 (Matrix Representation of the Bilinear Map). $\mathbb{D}^2 f(x) : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ is given by

$$\begin{bmatrix} \frac{\partial^2 f}{\partial x_1 \partial x_1} & \frac{\partial^2 f}{\partial x_2 \partial x_1} & \cdots & \frac{\partial^2 f}{\partial x_n \partial x_1} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 f}{\partial x_1 \partial x_n} & \frac{\partial^2 f}{\partial x_2 \partial x_n} & \cdots & \frac{\partial^2 f}{\partial x_n \partial x_n} \end{bmatrix}_{n \times n}$$

This matrix is denoted as $H_x(f)$, the Hessian matrix of f at x . Then, in matrix form, we have that for

$u = (u_1, u_2, \dots, u_n) \in \mathbb{R}^n$ and $v = (v_1, v_2, \dots, v_n) \in \mathbb{R}^n$, and

$$\mathbb{D}^2 f(x)(u, v) = u \cdot H_x(f) \cdot v^\top \in \mathbb{R}.$$

Proof 4. Note that

$$g(x) = \left(\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_n} \right) : \mathbb{R}^n \rightarrow \mathbb{R}^n.$$

Then,

$$\begin{aligned} \mathbb{D}^2 f(x) &= \mathbb{D}g(x) \\ &= \begin{bmatrix} \frac{\partial}{\partial x_1} \frac{\partial f}{\partial x_1} & \frac{\partial}{\partial x_2} \frac{\partial f}{\partial x_1} & \cdots & \frac{\partial}{\partial x_n} \frac{\partial f}{\partial x_1} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial}{\partial x_1} \frac{\partial f}{\partial x_n} & \frac{\partial}{\partial x_2} \frac{\partial f}{\partial x_n} & \cdots & \frac{\partial}{\partial x_n} \frac{\partial f}{\partial x_n} \end{bmatrix}. \end{aligned}$$

Q.E.D. ■

Lemma 6.6.11 Symmetry of the Partial and Differentials: Let $f(x, y) : A \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ be of class \mathcal{C}^2 . Then,

$$\frac{\partial^2 f}{\partial x \partial y} = \frac{\partial^2 f}{\partial y \partial x}.$$

In general, for $f : A \subset \mathbb{R}^n \rightarrow \mathbb{R}$ in class \mathcal{C}^2 ,

$$\frac{\partial^2 f}{\partial x_i \partial x_j} = \frac{\partial^2 f}{\partial x_j \partial x_i} \quad \forall i, j.$$

Extension 6.1 If $f \in \mathcal{C}^{(n)}$, the order of taking n -th derivative does not matter.

Corollary 6.6.12 : If f is of class \mathcal{C}^2 , then $\mathbb{D}^2 f(x) : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ is symmetric. That is,

$$\mathbb{D}^2 f(x)(u, v) = \mathbb{D}^2 f(x)(v, u).$$

Proof 5.

$$\mathbb{D}^2 f(x)(u, v) = u \cdot H_x(f) \cdot v^\top$$

Since $\mathbb{D}^2 f(x)(u, v) \in \mathbb{R}$, we have

$$\begin{aligned} \mathbb{D}^2 f(x)(u, v) &= [\mathbb{D}^2 f(x)(u, v)]^\top = (u \cdot H_x(f) \cdot v^\top)^\top \\ &= v \cdot H_x(f)^\top \cdot u^\top \\ &= v \cdot H_x(f) \cdot u^\top && \text{[by symmetry of } H_x(f)] \\ &= \mathbb{D}^2 f(x)(v, u). \end{aligned}$$

Q.E.D. ■

Example 6.6.13 Symmetry of Partial

Let $f(x, y, z) = e^{xy} + xyz : \mathbb{R}^3 \rightarrow \mathbb{R}$. Verify the symmetry of the partials.

Solution 6.

$$\frac{\partial f}{\partial x} = ye^{xy} + yz; \quad \frac{\partial f}{\partial y} = xe^{xy} + yz; \quad \frac{\partial f}{\partial z} = xy.$$

$$\frac{\partial^2 f}{\partial y \partial x} = \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial x} \right) = e^{xy} + xye^{xy} + z;$$

$$\frac{\partial^2 f}{\partial x \partial y} = \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial y} \right) = e^{xy} + xye^{xy} + z.$$

□

Summary V: Higher Order Differentials

- 1-st Order Differential: $\mathbb{D}f(x_0) : \mathbb{R}^n \rightarrow \mathbb{R}$: 1-linear form

$$\mathbb{D}f(x_0)(v) = J_f(x_0) \cdot v = \sum_{i=1}^n \frac{\partial f}{\partial x_i} v_i.$$

- 2-nd Order Differential: $\mathbb{D}^2 f(x_0) : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$: bilinear form

$$\mathbb{D}^2 f(x_0)(v, w) = v \cdot H_f(x_0) \cdot w^\top = \sum_{i,j=1}^n \frac{\partial^2 f}{\partial x_i \partial x_j} v_i \cdot w_j.$$

- k -th Order Differential: $\mathbb{D}^k f(x_0) : \mathbb{R}^n \times \mathbb{R}^n \times \dots \times \mathbb{R}^n \rightarrow \mathbb{R}$: k -linear form

$$\mathbb{D}^k f(x_0)(v^{(1)}, v^{(2)}, \dots, v^{(k)}) = \sum_{i_1, i_2, \dots, i_k=1}^n \frac{\partial^k f}{\partial x_{i_1} \partial x_{i_2} \dots \partial x_{i_k}} v_{i_1}^{(1)} v_{i_2}^{(2)} \dots v_{i_k}^{(k)}$$

In particular, denote $h = x - x_0 \in \mathbb{R}^n$, then

$$\mathbb{D}^k f(x_0)(h, h, \dots, h) = \sum \frac{\partial^k f}{\partial x_{i_1} \partial x_{i_2} \dots \partial x_{i_k}} h_{i_1} h_{i_2} \dots h_{i_k}.$$

- Speical case: $n = 2$: Write $\mathbb{D}^k f(x_0)(h, h) = \left(\frac{\partial}{\partial x_1} + \frac{\partial}{\partial x_2} \right)^k f(x_0) \cdot (h, h)$. Then,

$$\begin{aligned} \mathbb{D}^1 f &= \left(\frac{\partial}{\partial x_1} + \frac{\partial}{\partial x_2} \right)^1 f = \frac{\partial f}{\partial x_1} + \frac{\partial f}{\partial x_2}; \quad \mathbb{D}^2 f = \left(\frac{\partial}{\partial x_1} + \frac{\partial}{\partial x_2} \right)^2 f = \frac{\partial^2 f}{\partial x_1^2} + 2 \frac{\partial^2 f}{\partial x_1 \partial x_2} + \frac{\partial^2 f}{\partial x_2^2}, \\ \mathbb{D}^3 f(h, h, h) &= \frac{\partial^3 f}{\partial x_1^3} h_1^3 + 3 \frac{\partial^3 f}{\partial x_1^2 \partial x_2} h_1^2 h_2 + 3 \frac{\partial^3 f}{\partial x_1 \partial x_2^2} h_1 h_2^2 + \frac{\partial^3 f}{\partial x_2^3} h_2^3 \end{aligned}$$

Theorem 6.6.14 Taylor's Theorem

Let $f : A \subset \mathbb{R}^n \rightarrow \mathbb{R}$ be of class \mathcal{C}^r . Suppose $x, x_0 \in A$ s.t. the line segment joining x and x_0 , $[x, x_0] \subset A$. Then, $\exists c \in [x, x_0]$ s.t.

$$f(x) = f(x_0) + \mathbb{D}f(x_0)(x - x_0) + \frac{1}{2!}\mathbb{D}^2f(x_0)(x - x_0, x - x_0) + \cdots \\ + \frac{1}{(r-1)!}\mathbb{D}^{r-1}f(x_0)(x - x_0, x - x_0, \dots, x - x_0) + R_{r-1},$$

where R_{r-1} is the remainder given by

$$R_{r-1} = \frac{1}{r!}\mathbb{D}^r f(c)(x - x_0, \dots, x - x_0)$$

and satisfies

$$\frac{R_{r-1}(x_0)}{\|x - x_0\|^{r-1}} \rightarrow 0 \quad \text{as } x \rightarrow x_0.$$

Proof 7. Consider 1-variable form, $\varphi(t) = x_0 + t(x - x_0)$. Define

$$g(t) = f(x_0 + t(x - x_0))$$

for $t \in (a, b)$ with $[0, 1] \subset (a, b)$.

Apply Taylor's Theorem in 1-D to $g(t)$, we get

$$g(1) = g(0) + g'(0)(1 - 0) + \frac{g''(0)}{2!}(1 - 0)^2 + \cdots + \frac{g^{(r-1)}(0)}{(r-1)!}(1 - 0)^{r-1} + R_{r-1}$$

$$f(x) = f(x_0) + \sum_{k=1}^{r-1} \frac{g^{(k)}(0)}{k!} + \frac{1}{r!}g^{(r)}(\tilde{c}), \quad \tilde{c} \in [0, 1].$$

By chain rule, one can get

$$g'(t) = \mathbb{D}f(\varphi(t))\varphi'(t)$$

$$g'(0) = \mathbb{D}f(x_0)(x - x_0)$$

$$g''(t) = \mathbb{D}^2f(\varphi(t))\varphi'(t) \cdot \varphi'(t)$$

$$g''(0) = \mathbb{D}^2f(x_0)(x - x_0)^2 = \mathbb{D}^2f(x_0)(x - x_0, x - x_0).$$

So,

$$g^{(k)}(0) = \mathbb{D}^k f(x_0)(x - x_0, x - x_0, \dots, x - x_0).$$

Q.E.D. ■

Example 6.6.15 Polynomial Approximation using Taylor's Theorem

Determine the 2-nd order Taylor's formula for $f(x, y) = e^{(x-1)^2} \cos y$ at $(1, 0)$.

Solution 8.

- Compute partials:

$$\frac{\partial f}{\partial x} = 2(x-1)e^{(x-1)^2} \cos y; \quad \frac{\partial f}{\partial y} = -e^{(x-1)^2} \sin y.$$

$$\frac{\partial^2 f}{\partial x^2} = 2e^{(x-1)^2} \cos y + 4(x-1)^2 e^{(x-1)^2} \cos y; \quad \frac{\partial^2 f}{\partial y^2} = -e^{(x-1)^2} \cos y.$$

$$\frac{\partial^2 f}{\partial x \partial y} = -2(x-1)e^{(x-1)^2} \sin y$$

- Evaluate at base point $(1, 0)$:

$$\left. \frac{\partial f}{\partial x} \right|_{(1,0)} = 0, \quad \left. \frac{\partial f}{\partial y} \right|_{(1,0)} = 0, \quad \left. \frac{\partial^2 f}{\partial x^2} \right|_{(1,0)} = 2, \quad \left. \frac{\partial^2 f}{\partial x \partial y} \right|_{(1,0)} = 0, \quad \left. \frac{\partial^2 f}{\partial y^2} \right|_{(1,0)} = -1.$$

- Taylor's Formula: $h = x - x_0 = (x, y) - (1, 0)$.

$$f(x, y) = f(1, 0) + \mathbb{D}f(1, 0)(h) + \frac{1}{2} \mathbb{D}^2 f(1, 0)(h, h) + R_2,$$

where $f(1, 0) = 1$, $\mathbb{D}f(1, 0) = \begin{bmatrix} 0 & 0 \end{bmatrix}$, and $\mathbb{D}^2 f(1, 0) = \begin{bmatrix} 2 & 0 \\ 0 & -1 \end{bmatrix}$. So,

$$\mathbb{D}f(1, 0)(h) = 0$$

$$\mathbb{D}^2 f(1, 0)(h, h) = \begin{pmatrix} x-1 & y \end{pmatrix} \begin{bmatrix} 2 & 0 \\ 0 & -1 \end{bmatrix} \begin{pmatrix} x-1 \\ y \end{pmatrix} = 2(x-1)^2 - y^2.$$

Then,

$$f(x, y) = 1 + \frac{1}{2} [2(x-1)^2 - y^2] + R_2,$$

where

$$\frac{R_2}{\|(x-1, y)\|^2} \rightarrow 0 \quad \text{as} \quad (x-1, y) \rightarrow (1, 0).$$

□

6.7 Minima & Maxima in \mathbb{R}^n

Question: Given function $f : A \subset \mathbb{R}^n \rightarrow \mathbb{R}$, how do we find (local) maximum or minimum points for f in A ?

6.7.1 Optimization in 1-D. Suppose $f : (a, b) \rightarrow \mathbb{R}$

- A local max/min point (or extreme point) x_0 must be a critical point:

$$f'(x_0) = 0 \quad \text{or} \quad f'(x_0) \text{ D.N.E.}$$

- 2-nd Order Derivative Test (for critical points):

$$f''(x_0) > 0 : \text{local min}; \quad f''(x_0) < 0 : \text{local max.}$$

Definition 6.7.2 (Extrema). Suppose $f : A \subset \mathbb{R}^n \rightarrow \mathbb{R}$.

- Then, $x_0 \in A$ is a *local minimum* if $\exists \delta > 0$ s.t. $x \in A$ and

$$|x - x_0| < \delta \implies f(x) \geq f(x_0).$$


- Similarly, $x_0 \in A$ is a *local maximum* if $\exists \delta > 0$ s.t. $x \in A$ and

$$|x - x_0| < \delta \implies f(x) \leq f(x_0).$$

Theorem 6.7.3 Necessary Condition for Extreme Points

If $f : A \subset \mathbb{R}^n \rightarrow \mathbb{R}$ is differentiable and $x_0 \in A$ is an extreme point for f , then x_0 is a *critical point*, i.e., $\mathbb{D}f(x_0) = 0$.

Remark 6.9 This is only a necessary condition but not sufficient. For example, in \mathbb{R}^1 , $f(x) = x^2$ at $(0, 0)$ or in \mathbb{R}^2 , $f(x, y) = x^2 - y^2$ at $(0, 0)$.



For a critical point that is not an extreme point, we call it a saddle point.

Proof 1. (Sketch).

Assume $\mathbb{D}f(x_0) \neq 0$. Then, WLOG, $\exists v \in \mathbb{R}^n$ s.t. $\mathbb{D}f(x_0)(v) = c > 0$. By definition of differential, choose $\delta > 0$ s.t.

$$\|f(x_0 + h) - f(x_0) - \underbrace{\mathbb{D}f(x_0)(h)}_{=\varepsilon}\| < \frac{c}{2\|v\|} \cdot \|h\| \quad \forall \|h\| < \delta.$$

Choose $h = \lambda v$ with $\lambda > 0$ and $\|h\| < \delta$. Then, by triangle inequality,

$$f(x_0 + \lambda v) - f(x_0) > 0 \quad \text{but} \quad f(x_0 - \lambda v) - f(x_0) < 0.$$

Contradiction!

Q.E.D. ■

Definition 6.7.4 (Positive/Negative (Semi)definite). A bilinear form $B : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ is called *positive definite* (or *negative definite*) if $B(x, x) > 0$ (or < 0) $\forall x \in \mathbb{R}^n, x \neq 0$. We say B is *positive* (or *negative*) *semidefinite* if $B(x, x) \geq 0$ (or ≤ 0) $\forall x \in \mathbb{R}^n$.

Definition 6.7.5 (Major Diagonal Factors). Recall B is determined by a matrix H as follows:

$$B(x, x) = xHx^\top, \quad \text{where } H = \begin{bmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & \ddots & \vdots \\ a_{n1} & \cdots & a_{nn} \end{bmatrix}_{n \times n}$$

The *major diagonal factors* are given by

$$\begin{aligned} \Delta_1 &= \det \begin{pmatrix} a_{11} \end{pmatrix} = a_{11} \\ \Delta_2 &= \det \begin{pmatrix} a_{11} & a_{21} \\ a_{12} & a_{22} \end{pmatrix} \\ &\vdots \\ \Delta_n &= \det(H). \end{aligned}$$

Lemma 6.7.6 :

- H is positive definite $\iff \Delta_k > 0 \quad \forall k = 1, \dots, n$
- H is positive semi-definite $\implies \Delta_k \geq 0 \quad \forall k = 1, \dots, n$.
- H is negative definite $\iff (-H)$ is positive definite.

Example 6.7.7

$$H = \begin{bmatrix} 2 & -1 \\ -1 & 3 \end{bmatrix} \implies \Delta_1 = 2, \Delta_2 = 5 \implies H \text{ is positive definite.}$$

Theorem 6.7.8 Second Order Sufficient Condition

Suppose $f : A \subset \mathbb{R}^n \rightarrow \mathbb{R}$ is of class \mathcal{C}^2 and $x_0 \in A$ is a critical point (i.e., $\mathbb{D}f(x) = 0$).

- If $H_f(x_0)$ is negative (or positive) definite, then x_0 is a local maximum (or minimum).
- If x_0 is a local maximum (or minimum), then $H_f(x_0)$ is negative (or positive) semidefinite.

Remark 6.10

- $\text{Max of } f \iff \text{Min of } (-f)$
- *About minimum point:*
 - $\Delta_k > 0 \quad \forall k, H_f(x_0) \text{ is positive definite} \implies x_0 \text{ is local minimum.}$

- x_0 is a local minimum $\implies H_f(x_0)$ is positive semidefinite $\implies \Delta_k \geq 0 \quad \forall k$.
- $\Delta_k < 0$ for some $k \implies x_0$ is not a local minimum.
- About maximum point:
 - $\Delta_k < 0$ for odd k and $\Delta_k > 0$ for even $k \implies (-H_f(x_0))$ is negative definite $\implies H_f(x_0)$ is negative definite $\implies x_0$ is local maximum.
 - x_0 is local maximum $\implies H_f(x_0)$ is negative semidefinite $\implies \Delta_k \leq 0$ for odd k and $\Delta_k \geq 0$ for even k .
 - $\Delta_k < 0$ for some even $k \implies x_0$ is not a local maximum $\implies x_0$ is a saddle point.

Proof2. (of ①)

- Set-up: Suppose H_f is negative definite. Need to show:

$$\exists \delta > 0 \text{ s.t. } \|y - x\| < \delta \implies f(y) \leq f(x_0). \quad (\star)$$

Scartch:

By Taylor's Theorem

$$\begin{aligned} f(y) &= f(x_0) + \underbrace{\mathbb{D}f(x_0)}_{=0, \text{critical point}} (y - x_0) + \frac{1}{2} \mathbb{D}^2 f(c)(y - x_0, y - x_0) \\ f(y) - f(x_0) &= \frac{1}{2} \mathbb{D}^2 f(c)(y - x_0, y - x_0). \end{aligned}$$

If $\mathbb{D}^2 f(c)$ is negative semidefinite, we are done with the proof. However, we only know definiteness at x_0 . Let's add and subtract $\mathbb{D}^2 f(x_0)$:

$$f(y) - f(x_0) = \frac{1}{2} \underbrace{\mathbb{D}^2 f(x_0)(y - x_0, y - x_0)}_{\text{negative}} + \frac{1}{2} \underbrace{[\mathbb{D}^2 f(c) - \mathbb{D}^2 f(x_0)](y - x_0, y - x_0)}_{\text{make it small}}$$

- Consider the function

$$g(x) = \mathbb{D}^2 f(x_0)(x, x) : \mathbb{R}^n \rightarrow \mathbb{R}.$$

Denote $\mathbb{D}^2 f(x_0) = H$, then $g(x) = H(x, x)$. g is continuous. Then, $\exists \bar{x} \in S = \{x \in \mathbb{R}^n \mid \|x\| = 1\}$ s.t.

$$H(x, x) \leq H(\bar{x}, \bar{x}).$$

Extreme Value Theorem: Continuous function on closed and bounded set attains its maximum and minimum. Since H is negative definite, $H(\bar{x}, \bar{x}) < 0$. Let $\varepsilon = -H(\bar{x}, \bar{x}) > 0$. Then, for any

$h \in \mathbb{R}^n$ with $h \neq 0$, we have

$$H(h, h) = \|h^2\| \cdot H\left(\frac{h}{\|h\|}, \frac{h}{\|h\|}\right) \leq \|h^2\| \cdot H(\delta x, \bar{x}).$$

So,

$$H(h, h) \leq -\varepsilon \|h^2\| \quad (\text{I})$$

- Prove (\star) is true in a neighborhood.

By continuity of $\mathbb{D}^2 f$ at x_0 , $\exists \delta > 0$ s.t.

$$\|y - x_0\| < \delta \implies y \in A, \underbrace{\|\mathbb{D}^2 f(y) - \mathbb{D}^2 f(x_0)\|}_{\text{operator norm}} < \frac{\varepsilon}{2} \quad (\text{II})$$

Operator norm satisfies: $\|T(x, y)\| \leq \|T\| \cdot \|x\| \cdot \|y\|$.

By Taylor's Formula, because $\mathbb{D}f(x_0) = 0$, we have

$$f(y) - f(x) = \frac{1}{2} \mathbb{D}^2 f(c)(h, h),$$

where $y \in B(x_0, \delta)$, $h = y - x_0$, and $c \in [x_0, y]$. Note that

$$\begin{aligned} \mathbb{D}^2 f(c)(h, h) &= [\mathbb{D}^2 f(c) - \mathbb{D}^2 f(x_0)](h, h) + \mathbb{D}^2 f(x_0)(h, h) \\ &\leq \|\mathbb{D}^2 f(c) - \mathbb{D}^2 f(x_0)\| \cdot \|h\|^2 + (-\varepsilon) \|h\|^2 && \text{By (I)} \\ &\leq \frac{1}{2} \varepsilon \|h\|^2 + (-\varepsilon) \|h\|^2 && \text{By (II)} \\ &= -\frac{\varepsilon}{2} \|h\|^2 \leq 0. \end{aligned}$$

Then, $f(y) \leq f(x) \quad \forall y \in B(x_0, \delta)$. So, x_0 is the local maximum.

Q.E.D. ■

Example 6.7.9

Find and classify the critical points for $f(x, yz) = \cos 2x \sin y + z^2$.

Solution 3.

- Find the critical point:

$$\frac{\partial f}{\partial x} = -2 \sin 2x \sin y; \quad \frac{\partial f}{\partial y} = \cos 2x \cos y; \quad \frac{\partial f}{\partial z} = 2z.$$

Set

$$\frac{\partial f}{\partial x} = \frac{\partial f}{\partial y} = \frac{\partial f}{\partial z} = 0.$$

Then,

$$\begin{cases} -2 \sin 2x \sin y = 0 \\ \cos 2x \cos y = 0 \\ 2z = 0 \end{cases} \implies \begin{cases} x = \frac{k\pi}{2} \\ y = \frac{2j+1}{2}\pi \\ z = 0 \end{cases} \quad \text{or} \quad \begin{cases} x = \frac{2k+1}{4}\pi \\ y = j\pi \\ z = 0. \end{cases}$$

- Classify critical points:

Compute the Hessian

$$\frac{\partial^2 f}{\partial x^2} = -4 \cos 2x \sin y; \quad \frac{\partial^2 f}{\partial y \partial x} = -2 \sin 2x \cos y; \quad \frac{\partial^2 f}{\partial z \partial x} = 0$$

$$\frac{\partial^2 f}{\partial y^2} = -\cos 2x \sin y; \quad \frac{\partial^2 f}{\partial z \partial y} = 0; \quad \frac{\partial^2 f}{\partial z^2} = 2.$$

So,

$$H_f(x) = \begin{bmatrix} -4 \cos 2x \sin y & -2 \sin 2x \cos y & 0 \\ -2 \sin 2x \cos y & -\cos 2x \sin y & 0 \\ 0 & 0 & 2 \end{bmatrix}.$$

Case I $x = \frac{k\pi}{2}, y = \frac{2j+1}{2}\pi, z = 0$. Then,

$$H_f\left(\frac{k\pi}{2}, \frac{2j+1}{2}\pi, 0\right) = \begin{bmatrix} -4(-1)^k(-1)^j & 0 & 0 \\ 0 & -1(-1)^k(-1)^j & 0 \\ 0 & 0 & 2 \end{bmatrix}.$$

Then, $\Delta_1 = -4(-1)^{j+k}, \Delta_2 = 4(-1)^{2k}(-1)^{2j} = 4 > 0$, and $\Delta_3 = 2 \cdot \Delta_2 = 8 > 0$.

- If $j + k$ is odd, then $\Delta_1 > 0$. Then, H_f is positive definite, and the critical point is a local minimum.
- If $j + k$ is even, then $\Delta_1 < 0$. Then, the critical point is not a local minimum. But $\Delta_3 = 0 > 0$, so it cannot be a local maximum. Hence, it must be a saddle point.

Case II $x = \frac{2k+1}{4}\pi, y = j\pi, z = 0$. Then,

$$H_f\left(\frac{2k+1}{4}\pi, j\pi, 0\right) = \begin{bmatrix} 0 & (-2)(-1)^k(-1)^j & 0 \\ (-2)(-1)^k(-1)^j & 0 & 0 \\ 0 & 0 & 2 \end{bmatrix}.$$

Then, $\Delta_1 = 0, \Delta_2 = -(-2)(-1)^{k+j} \cdot (-2)(-1)^{k+j} = -4(-1)^{2(k+j)} = -4 < 0$, and $\Delta_3 = 0$. As $\Delta_2 < 0$, they are saddle points.

- Conclusion:

$$\left(\frac{k\pi}{2}, \frac{2j+1}{2}\pi, 0\right) \begin{cases} \text{local minimum when } k+j \text{ is odd} \\ \text{saddle point when } k+j \text{ is even.} \end{cases}$$

$$\left(\frac{2k+1}{4}\pi, j\pi, 0\right) : \text{saddle point.}$$

□

7 Inverse and Implicit Function Theorem

7.1 Inverse Function Theorem

7.1.1 Linear Case.

- Consider a linear map: $y = f(x) : \mathbb{R}^n \rightarrow \mathbb{R}^n$ given by

$$\begin{cases} y_1 &= a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n \\ &\vdots \\ y_n &= a_{n1}x_1 + a_{n2}x_2 + \cdots + a_{nn}x_n. \end{cases}$$

Or, in matrix notation:

$$Ax = y \quad (\star)$$

- Given $y \in \mathbb{R}^n$, (\star) is a linear system of equations.
- Fact:** (\star) has unique solution $x \iff A$ is invertible. i.e., $\det(A) \neq 0$. In this case, the solution is given by $x = A^{-1}y$.
- $x = A^{-1}y$ is the inverse function of $y = f(x)$.

7.1.2 When can we solve a nonlinear system?.

- Nonlinear System:

$$\begin{cases} f_1(x_1, x_2, \dots, x_n) &= y_1 \\ &\vdots \\ f_n(x_1, x_2, \dots, x_n) &= y_n \end{cases}, \quad \text{or } f(x) = y.$$

In order to have inverse, dimension must match.

• Notation 7.3.

- $y = f(x) : A \subset \mathbb{R}^n \rightarrow \mathbb{R}^n$, where A is open and f is differentiable on A . Suppose $y = (y_1, y_2, \dots, y_n)$, $x = (x_1, x_2, \dots, x_n)$, and $f = (f_1, f_2, \dots, f_n)$.
- $\mathbb{D}f(x) = \left(\frac{\partial f_j}{\partial x_i} \right)_{ij}$ and $J_f(x) = \det(\mathbb{D}f(x))$ is the *Jacobian determinant* of f at x .

Theorem 7.1.4 Inverse Function Theorem

Let $y = f(x) : A \subset \mathbb{R}^n \rightarrow \mathbb{R}^n$ be of class C^1 . Suppose $x_0 \in A$ and $J_f(x_0) \neq 0$. Then, \exists neighborhoods U of x_0 and W of $y_0 = f(x_0)$ s.t.

- $f(U) = W$ and $f : U \rightarrow W$ has an inverse $f^{-1} : W \rightarrow U$
- $f^{-1} : W \rightarrow U$ is of class C^1 . Additionally, if $f \in C^r$, then $f^{-1} \in C^r$.
- $\mathbb{D}f^{-1}(y) = [\mathbb{D}f(x)]^{-1} \quad \forall y \in W \text{ and } y = f(x)$.

► **Proof 1 of Inverse Function Theorem**

Theorem (Contraction Mapping Principle / CMP) Let \mathcal{X} be a complete metric space and $\varphi : \mathcal{X} \rightarrow \mathcal{X}$. Suppose $\exists 0 < k < 1$ s.t.

$$d(\varphi(x), \varphi(y)) \leq k \cdot d(x, y) \quad \forall x, y \in \mathcal{X}.$$

Then, \exists unique fixed point x^* s.t. $\varphi(x^*) = x^*$.

Step 1 Reductions

- We may assume that $\mathbb{D}f(x_0) = I$.

In fact, let $T = \mathbb{D}f(x_0)$. Then, $J_f(x_0) \neq 0 \implies T^{-1}$ exists. Consider a new map: $T^{-1} \circ f : A \rightarrow \mathbb{R}$. Then,

$$\begin{aligned} \mathbb{D}(T^{-1} \circ f) &= \mathbb{D}T^{-1}(f(x_0)) \circ \mathbb{D}f(x_0) \\ &= T^{-1} \circ T \\ &= I. \end{aligned}$$

If the inverse of $T^{-1} \circ f$ exists, then the inverse of f also exists. So, once the identity case is true, we just multiply T^{-1} to f and we can get the general case is true.

- We may assume that $x_0 = 0$ and $f(x_0) = 0$.

To see this, let $h(x) = f(x + x_0) - f(x_0)$. Then, $h(0) = 0$ and $\mathbb{D}h(0) = \mathbb{D}f(x_0)$. If the inverse of $h(x)$ exists, then the equation $f(x) = y$ can be solved:

$$\begin{aligned} f(x) &= h(x - x_0) + f(x_0) = y \\ h(x - x_0) &= y - f(x_0) \\ x - x_0 &= h^{-1}(y - f(x_0)) \\ x &= h^{-1}(y - f(x_0)) + x_0. \end{aligned}$$

Step 2 Existence of Inverse

- By reduction above, we have $x_0 = 0$, $y_0 = f(x_0) = 0$, $\mathbb{D}f(x_0) = \mathbb{D}f(0) = I$.

WTS: \exists neighborhoods U, W of 0 s.t. the map $y = f(x) : U \rightarrow W$ has an inverse in W . i.e., $\forall y \in W, \exists$ unique $x \in U$ s.t. $y = f(x)$.

For a fixed $y \in \mathbb{R}^n$, define $g_y(x) := y + x - f(x) : A \rightarrow \mathbb{R}^n$.

If $g_y(x)$ has a fixed point: $g_y(x^*) = x^* = y + x^* - f(x^*) \implies y - f(x^*) = 0$. So, we want to show $g_y(x)$ has a unique fixed point.

- Construction of neighborhoods U and W .

Let $g(x) = x - f(x)$. Then,

$$\mathbb{D}g(0) = I - \mathbb{D}f(0) = I - I = 0.$$

Since $f \in \mathcal{C}^1$, $g \in \mathcal{C}^1$. Then, $\mathbb{D}g(x)$ is continuous at 0. Then, $\forall \varepsilon = \frac{1}{2n}$, $\exists \delta > 0$ s.t.

$$\|x - 0\| < \delta \implies \|\mathbb{D}g_i(x) - \mathbb{D}g_i(0)\| = \|\mathbb{D}g_i(x) - 0\| = \|\mathbb{D}g_i(x)\| < \frac{1}{2n},$$

where $g = (g_1, g_2, \dots, g_n)$.

Apply MVT to each of g_i , we obtain $\forall x \in \overline{B}(x_0, \delta)$, $\exists c_i \in [0, x]$ s.t.

$$g_i(x) = g_i(x) - g_i(0) = \mathbb{D}g_i(c_i)(x - 0).$$

So,

$$\begin{aligned} \|g(x)\| &\leq \sum_{i=1}^n \|g_i(x)\| = \sum_{i=1}^n |\mathbb{D}g_i(c_i) \cdot x| \\ &\leq \sum_{i=1}^n \|\mathbb{D}g_i(c_i)\| \cdot \|x\| && \text{[operator norm]} \\ &\leq \sum_{i=1}^n \frac{1}{2n} \|x\| && \text{[continuity of } \mathbb{D}g\text{]} \\ &= \frac{1}{2} \|x\|. \end{aligned}$$

i.e., $\|g(x)\| \leq \frac{1}{2} \|x\|$. Thus, $g : \overline{B}(0, \delta) \rightarrow \overline{B}(0, \frac{1}{2}\delta) \subset \overline{B}(0, \delta)$ is a contraction map. Let $W = B(0, \frac{\delta}{2})$ and $U = \{x \in B(0, \delta) : f(x) \in W\}$. **WTS: U and W are the desired neighborhoods.**

- Show existence of $f^{-1} : W \rightarrow U$.

Fix $y \in W$. Then, $\forall x \in \overline{B}(0, \delta)$,

$$\begin{aligned} \|g_y(x)\| &= \|y + g(x)\| \leq \|y\| + \|g(x)\| \\ &< \frac{\delta}{2} + \frac{1}{2}\delta = \delta \quad \left[y \in W = B\left(0, \frac{\delta}{2}\right), \|g(x)\| \leq \frac{1}{2}\|x\|, x \in U = B(0, \delta) \right] \end{aligned}$$

Then, $g_y(x) : \overline{B}(0, \delta) \rightarrow \overline{B}(0, \delta)$ and g_y is also a contraction map with $k = \frac{1}{2}$. Then, by CMP, \exists unique x s.t. $g_y(x) = x$. Then,

$$\begin{aligned} g_y(x) &= y + x - f(x) = x \\ y - f(x) &= 0 \implies y = f(x). \end{aligned}$$

So, for fixed y , \exists unique x s.t. $y = f(x)$. Then, f is a bijection, and thus the inverse exists.

Step 3 Continuity of f^{-1} .

WTS: f^{-1} is Lipschitz continuous.

Fix $y_1, y_2 \in W$. Let $x_i = f^{-1}(y_i)$ for $i = 1, 2$. Then,

$$\begin{aligned} \|f^{-1}(y_1) - f^{-1}(y_2)\| &= \|x_1 - x_2\| = \|g(x_1) + f(x_1) - g(x_2) - f(x_2)\| \\ &\leq \|g(x_1) - g(x_2)\| + \|f(x_1) - f(x_2)\| \\ &= \|g(x_1) - g(x_2)\| + \|y_1 - y_2\|. \end{aligned}$$

Since $\|\mathbb{D}g(x)\| \leq \frac{1}{2}$ for $x \in \overline{B}(0, \delta)$, by Mean Value Inequality,

$$\|g(x_1) - g(x_2)\| \leq \frac{1}{2}\|x_1 - x_2\|.$$

Then,

$$\|x_1 - x_2\| \leq \frac{1}{2}\|x_1 - x_2\| + \|y_1 - y_2\|.$$

So,

$$\frac{1}{2}\|x_1 - x_2\| \leq \|y_1 - y_2\| \implies \|x_1 - x_2\| \leq 2\|y_1 - y_2\|.$$

That is,

$$\|f^{-1}(y_1) - f^{-1}(y_2)\| \leq 2\|y_1 - y_2\| \quad (\star)$$

Thus, f^{-1} is Lipschitz and thus continuous.

Step 4 Differentiability of f^{-1}

- **Proposition** $[\mathbb{D}f(0)]^{-1}$ exists and $\mathbb{D}f(x)$ is continuous at 0 $\implies \exists \delta' > 0$ s.t. $[\mathbb{D}f(x)]^{-1}$ exists and bounded by M :

$$\underbrace{\|\mathbb{D}f(x) \cdot (v)\|}_{\text{operator norm}} \leq \|M\| \cdot \|v\| \quad \forall \|x\| < \delta' \text{ and } v \in \mathbb{R}^n.$$

- **WTS:** $f^{-1}(y)$ is differentiable at any fixed point $y_0 \in W$ and

$$\mathbb{D}f^{-1}(y_0) = [\mathbb{D}f(x_0)]^{-1} \quad \text{with } y_0 = f(x_0).$$

Fix $y_0 \in W$. Then,

$$\begin{aligned} & \frac{\|f^{-1}(y) - f^{-1}(y_0) - \mathbb{D}f^{-1}(y_0) \cdot (y - y_0)\|}{\|y - y_0\|} \\ &= \frac{\|[\mathbb{D}f(x_0)]^{-1} \cdot [\mathbb{D}f(x_0) \cdot f^{-1}(y) - \mathbb{D}f(x_0) \cdot f^{-1}(y_0) - (y - y_0)]\|}{\|y - y_0\|} \quad [\text{factor out } \mathbb{D}f^{-1}(y_0) = [\mathbb{D}f(x_0)]^{-1}] \\ &= \frac{\|[\mathbb{D}f(x_0)]^{-1} \cdot [\mathbb{D}f(x_0)(x - x_0) - (f(x) - f(x_0))]\|}{\|f(x) - f(x_0)\|} \quad [y = f(x)] \\ &= \frac{\|[\mathbb{D}f(x_0)]^{-1} \cdot [\mathbb{D}f(x_0)(x - x_0) - (f(x) - f(x_0))]\| \cdot \|x - x_0\|}{\|f(x) - f(x_0)\| \cdot \|x - x_0\|} \quad [\text{Multiply by magic 1}] \\ &\leq \frac{2\|[\mathbb{D}f(x_0)]^{-1}[\mathbb{D}f(x_0)(x - x_0) - (f(x) - f(x_0))]\|}{\|x - x_0\|} \quad [\text{Lipschitz continuity, Eq } (\star)] \\ &\rightarrow 0 \quad \text{as } x \rightarrow x_0. \end{aligned}$$

So, f^{-1} is differentiable, and

$$[\mathbb{D}f^{-1}(y)] = [\mathbb{D}f(x)]^{-1}.$$

Q.E.D. ■

Example 7.1.5

Investigate the invertibility (both local and global) for the map $W = (u, v) = f(x, y) : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by $u = e^x \cos y$ and $v = e^x \sin y$.

Solution 2.

Firstly, we know $f \in C^\infty$. Compute the Jacobian determinant:

$$\begin{aligned} J_f(x, y) &= \det(\mathbb{D}f(x)) = \det \begin{pmatrix} \partial u / \partial x & \partial u / \partial y \\ \partial v / \partial x & \partial v / \partial y \end{pmatrix} \\ &= \det \begin{pmatrix} e^x \cos y & -e^x \sin y \\ e^x \sin y & e^x \cos y \end{pmatrix} \\ &= e^{2x} \cos^2 y + e^{2x} \sin^2 y \\ &= e^{2x} > 1. \end{aligned}$$

So, by the Inverse Function Theorem, f is invertible locally at any point, and the differentiable of the inverse is given by

$$\mathbb{D}f^{-1}(u, v) = [\mathbb{D}f(x, y)]^{-1} = \begin{bmatrix} e^x \cos y & -e^x \sin y \\ e^x \sin y & e^x \cos y \end{bmatrix}^{-1}.$$

Now, let's examine if f is globally invertible (i.e., if f is a one-to-one function on \mathbb{R}^2). Note that

$$f(x_0, y_0) = e^{x_0} \cos y_0$$

and

$$f(x_0, y_0 + 2\pi) = e^{x_0} \cos(y_0 + 2\pi) = e^{x_0} \cos(y_0) \text{ and } f(x_0, y_0 - 2\pi) = e^{x_0} \cos(y_0 - 2\pi) = e^{x_0} \cos(y_0).$$

So, f is not globally invertible since f is not an injection. □

Remark 7.1 f can be written in complex notation: $f(z) = e^z$, where $z = x + iy \in \mathbb{C}$. Then,

$$f(z) = e^z = e^{x+iy} = e^x (\cos x + i \sin y).$$

Meanwhile, $f^{-1}(z) = \ln(z)$.

7.2 Implicit Function Thm and Applications

Motivation

- Given a function $f : \mathbb{R} \rightarrow \mathbb{R}$. Consider an equation $f(y) = x$. If it can be solved for y (uniquely in terms of x), then the solution $y = g(x)$ is the inverse of f . That is, $(f \circ g)(x) = x$.
- Reinterpretation of Inverse:
Rewrite $f(y) = x$ as $x - f(y) = 0$ ①.
Then, f is invertible \iff Equation ① is solvable for y .
- **Question:** When can we solve a general equation for y , $F(x, y) = 0$ ($F : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^m$)?
The solution of $F(x, y) = 0$, denoted by $y = g(x)$, is called the *implicit function* determined by $F(x, y) = 0$.

Example 7.2.1

Consider equation $x^2 + y^2 - 1 = 0$ to be $F(x, y) : \mathbb{R}^1 \times \mathbb{R}^1 \rightarrow \mathbb{R}^1$.

Given (x_0, y_0) s.t. $F(x_0, y_0) = 0$ with $y_0 \neq 0$. Then, \exists a unique solution

$$y = \begin{cases} \sqrt{1 - x^2} & \text{if } y_0 > 0 \\ -\sqrt{1 - x^2} & \text{if } y_0 < 0. \end{cases}$$

in the neighborhood of x_0 .

Note that $\left. \frac{\partial F}{\partial y} \right|_{y=y_0} = 2y_0 \neq 0$ when $y_0 \neq 0$.

Theorem 7.2.2 Implicit Function Theorem

Let $A \subset \mathbb{R}^n \times \mathbb{R}^m$ and $F(x, y) : A \rightarrow \mathbb{R}^m$ be of class \mathcal{C}^1 . Suppose $(x_0, y_0) \in A$ with $F(x_0, y_0) = 0$. If

$$\begin{aligned} \Delta = \det \left(\frac{\partial F}{\partial y} \right) &= \det \left(\frac{\partial(F_1, \dots, F_m)}{\partial(y_1, \dots, y_m)} \right) \\ &= \det \begin{bmatrix} \frac{\partial F_1}{\partial y_1} & \dots & \frac{\partial F_1}{\partial y_m} \\ \vdots & \ddots & \vdots \\ \frac{\partial F_m}{\partial y_1} & \dots & \frac{\partial F_m}{\partial y_m} \end{bmatrix} \neq 0 \quad \text{at } (x_0, y_0), \end{aligned}$$

then \exists neighborhoods U of x_0 , V of y_0 , and a unique function $y = f(x) : U \rightarrow V$ such that $F(x, f(x)) = 0 \quad \forall x \in U$. i.e., $y = f(x)$ is the solution of $F(x, y) = 0$.

Furthermore, if $F \in \mathcal{C}^r$, then $f \in \mathcal{C}^r$.

Remark 7.2

- $y = f(x)$ is called the *implicit function determined by the equation* $F(x, y) = 0$ based at the point (x_0, y_0) .

- *Differential of implicit function:*

Suppose $n = m = 1$ and $F(x, y) = 0$. Then, by chain rule,

$$\begin{aligned} \frac{\partial F}{\partial x} \cdot \frac{dx}{dx} + \frac{\partial F}{\partial y} \cdot \frac{dy}{dx} &= 0 \\ \frac{dy}{dx} &= -\frac{\partial F / \partial x}{\partial F / \partial y}. \end{aligned}$$

In the general case, let $y = f(x) = (f_1, \dots, f_m) : \mathbb{R}^n \rightarrow \mathbb{R}^m$. Let f be the implicit function determined by $F(x, y) = 0$. Then,

$$\mathbb{D}f = -\left(\frac{\partial F}{\partial y}\right)^{-1} \cdot \left(\frac{\partial F}{\partial x}\right).$$

► Proof 1 of Implicit Function Theorem

Given $F(x, y) = 0$, $A \subset \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^m$. Consider the map $G : A \subset \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^n \times \mathbb{R}^m$ given by

$$G(x, y) = (x, F(x, y)).$$

We want to use Inverse Function Theorem. So, we need a map that maps to the same dimension.

Suppose G^{-1} exists in a neighborhood of (x_0, y_0) . Write

$$G^{-1}(x, 0) = (x, f(x)).$$

Then, $y = f(x)$ is the solution of $F(x, y) = 0$ because

$$\begin{aligned} G(x, f(x)) &= (x, 0) \\ &= (x, F(x, f(x))). \end{aligned}$$

So, $F(x, f(x)) = 0$.

It remains to show that G is invertible. This follows from the inverse function theorem. Consider

$$\mathbb{D}G \Big|_{(x,y)=(x_0,y_0)} = \begin{bmatrix} \begin{array}{cccc|cccc} 1 & 0 & \cdots & 0 & 0 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 & 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 & 0 & 0 & \cdots & 0 \end{array} \\ \hline \begin{array}{cccc|cccc} \frac{\partial F_1}{\partial x_1} & \cdots & \frac{\partial F_1}{\partial x_n} & \frac{\partial F_1}{\partial y_1} & \cdots & \frac{\partial F_1}{\partial y_m} \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ \frac{\partial F_m}{\partial x_1} & \cdots & \frac{\partial F_m}{\partial x_n} & \frac{\partial F_m}{\partial y_1} & \cdots & \frac{\partial F_m}{\partial y_m} \end{array} \end{bmatrix}.$$

So,

$$J_G(x_0, y_0) = \det \begin{bmatrix} \partial F_1 / \partial y_1 & \cdots & \partial F_1 / \partial y_m \\ \vdots & \ddots & \vdots \\ \partial F_m / \partial y_1 & \cdots & \partial F_m / \partial y_m \end{bmatrix} = \Delta \neq 0,$$

as assumed in implicit function theorem. Therefore, by the inverse function theorem, G is invertible.

Q.E.D. ■

Example 7.2.3

Discuss the solvability of $\begin{cases} y + x + uv = 0 \\ uxy + v = 0 \end{cases}$ for u, v in terms of x, y near the point $(0, 0, 0, 0)$ and the point $(1, 1, \sqrt{2}, -\sqrt{2})$. If impossible, compute $\frac{\partial u}{\partial x}$ and $\frac{\partial v}{\partial x}$ if exists.

Solution 2.

$$F(x, y, u, v) = 0 \text{ and } \begin{cases} F_1 = y + x + uv \\ F_2 = uxy + v. \end{cases} \quad \text{Let's compute } \Delta:$$

$$\begin{aligned} \Delta &= \det \left(\frac{\partial(F_1, F_2)}{\partial(u, v)} \right) = \det \begin{bmatrix} \partial F_1 / \partial u & \partial F_1 / \partial v \\ \partial F_2 / \partial u & \partial F_2 / \partial v \end{bmatrix} \\ &= \det \begin{bmatrix} v & u \\ xy & 1 \end{bmatrix} \\ &= v - uxy. \end{aligned}$$

Then, $\Delta(0, 0, 0, 0) = 0$. So, Implicit Function Theorem does not apply. On the other hand,

$$\Delta(1, 1, \sqrt{2}, -\sqrt{2}) = -\sqrt{2} - \sqrt{2} = -2\sqrt{2} \neq 0.$$

So, by Implicit Function Theorem, \exists unique solution $u = u(x, y)$ and $v = v(x, y)$ in a neighborhood.

Furthermore, the differentiable is given by

$$\begin{aligned} \frac{\partial(u, v)}{\partial(x, y)} &= - \left(\frac{\partial F}{\partial(u, v)} \right)^{-1} \left(\frac{\partial F}{\partial(x, y)} \right) \\ &= - \begin{bmatrix} v & u \\ xy & 1 \end{bmatrix}^{-1} \begin{bmatrix} 1 & 1 \\ uy & ux \end{bmatrix}. \end{aligned}$$

□

Theorem 7.2.4 Application: Domain-Straightening Theorem

Let $f : A \subset \mathbb{R}^n \rightarrow \mathbb{R}$. Suppose $\mathbb{D}f(x_0) \neq 0$ and $f(x_0) = 0$. Then, \exists open sets U and V (with $x_0 \in V$) and invertible map $h : U \rightarrow V$ s.t. $f(h(x_1, \dots, x_n)) = x_0$.

Remark 7.3 Under change of variables h , one can flatten the level curves of function $f(x)$.

Theorem 7.2.5 Application: Range-Straightening Theorem

Suppose $f : A \subset \mathbb{R}^p \rightarrow \mathbb{R}^n$ with $p < n$ and $\text{rank of } \mathbb{D}f(x_0) = p$. Then, \exists neighborhoods U, V , and invertible map $g : U \rightarrow V$ s.t. $g \circ f(x_1, \dots, x_p) = (x_1, \dots, x_p, 0, \dots, 0)$.

7.3 Constrained Extrema**7.3.1 Morse Theory: Local Behavior Near a Critical Point**

Let $f(x) : A \subset \mathbb{R}^n \rightarrow \mathbb{R}$ be of class \mathcal{C}^2 and x_0 is a critical point. Then, one can use $H_f(x_0)$ to classify critical point x_0 .

- Morse Theory makes this classification more precise.
- **Lemma 7.3.1 Morse Lemma:** Let $f(x) : A \subset \mathbb{R}^n \rightarrow \mathbb{R}$ be of class \mathcal{C}^2 with critical point $x_0 \in A$. If $H_f(x_0)$ is nondegenerate (i.e., $\det(H_f(x_0)) \neq 0$), then \exists neighborhoods U for x_0 and V for 0, and invertible map $g : V \rightarrow U$ s.t. the function $h = f \circ g$ has the form

$$h(y) = f(x_0) - [y_1^2 + y_2^2 + \dots + y_\lambda^2] + [y_{\lambda+1}^2 + \dots + y_n^2],$$

where λ is an integer called the *index* of f at x_0 .

- Interpretation/Application:
 1. $\lambda = 0$: x_0 is a local minimum. Paraboloid open up.
 2. $\lambda = n$: x_0 is a local maximum. Paraboloid open down.
 3. $0 < \lambda < n$: x_0 is a saddle point. Hyperboloid.
- What is λ ?

λ (the index of f at x_0) is the number of negative eigenvalues of $H_f(x_0)$.

Example 7.3.2

Determine the shape of the surface given by $z = x^2 + 3xy - y^2$ near critical point $(0, 0)$.

Solution 1.

$\mathbb{D}f = \begin{pmatrix} 2x + 3y & 3x - 2y \end{pmatrix}$. Therefore,

$$H_f(x, y) = \begin{bmatrix} 2 & 3 \\ 3 & -2 \end{bmatrix}$$

The eigenvalues are $t = \pm\sqrt{13}$. So, index $\lambda = 1$. As $0 < \lambda < n$, $(0, 0)$ is a saddle point. The shape is thus a hyperboloid.

□

7.3.2 Constrained Extremal Problem

Goal: To maximum (or minimize) a function $f(x) : \mathbb{R}^n \rightarrow \mathbb{R}$ under the constraint $g(x) = c$.

Tool: Lagrange Multiplier Method.

Theorem 7.3.3 Necessary Condition

Let $f, g : U \subset \mathbb{R}^n \rightarrow \mathbb{R}$ be of class C^1 . Assume $g(x_0) = c_0$ with $\nabla g(x_0) \neq 0$. If f restricted to the surface $S : g(x) = c_0$ has maximum or minimum at x_0 , then $\exists \lambda \in \mathbb{R}$ s.t.

$$\nabla f(x_0) = \lambda \nabla g(x_0).$$

Remark 7.4 (Geometric Meaning) $\nabla f(x_0)$ is parallel to $\nabla g(x_0)$.

Proof2.

- Geometric proof: **WTS:** $\nabla f(x_0) \perp S$.

Fix curve $c(t)$ at t_0 . So, $c(t_0) = x_0$. **WTS:** $\nabla f(x) \perp c'(t)$.

Since f restricted to S has a maximum at x_0 , $h(t) = f(c(t))$ has a maximum at t_0 . Then,

$$0 = h'(t_0) = \nabla f(x_0) \cdot c'(t_0) = \langle \nabla f(x_0), c'(t_0) \rangle.$$

So, $\nabla f(x_0) \perp c'(t_0)$, and thus $\nabla f(x_0) \perp S$.

- Analytical proof: **Substitute the condition** $g(x) = c_0$ **into** $f(x)$

Since

$$\nabla g(x_0) = \left(\frac{\partial g}{\partial x_1}, \dots, \frac{\partial g}{\partial x_n} \right) \neq \vec{0},$$

then $\exists \frac{\partial g}{\partial x_i} \neq 0$ for some $i = 1, \dots, n$. WLOG, assume $\frac{\partial g}{\partial x_n}(x_0) \neq 0$. By Implicit Function Theorem, the equation

$$g(x_1, \dots, x_n) = c_0$$

can be uniquely solve for x_n :

$$x_n = h(x_1, \dots, x_{n-1}).$$

Let $k(x_1, \dots, x_{n-1}) = f(x_1, \dots, x_{n-1}, h(x_1, \dots, x_{n-1}))$. Then, the maximum of f correspond to maximum of k . Then,

$$0 = \frac{\partial k}{\partial x_i} = \frac{\partial f}{\partial x_i} + \frac{\partial f}{\partial x_n} \cdot \frac{\partial h}{\partial x_i} \quad \text{for } i = 1, \dots, n-1. \quad (1)$$

Furthermore, $g(x) = c_0$. So, $g(x_1, \dots, x_{n-1}, h(x_1, \dots, x_{n-1})) = c_0$. Then,

$$\frac{\partial g}{\partial x_i} + \frac{\partial g}{\partial x_n} \cdot \frac{\partial h}{\partial x_i} = 0 \quad \text{for } i = 1, \dots, n-1.$$

Then,

$$\frac{\partial h}{\partial x_i} = -\frac{\partial g/\partial x_i}{\partial g/\partial x_n} \quad (2)$$

Substitute (2) into (1):

$$\begin{aligned} \frac{\partial f}{\partial x_i} &= -\frac{\partial f}{\partial x_n} \cdot \frac{\partial h}{\partial x_i} = -\frac{\partial f}{\partial x_n} \cdot \frac{-\partial g/\partial x_i}{\partial g/\partial x_n} \\ &= \underbrace{\frac{\partial f/\partial x_n}{\partial g/\partial x_n}}_{\lambda} \cdot \frac{\partial g}{\partial x_i} \\ &= \lambda \frac{\partial g}{\partial x_i}. \end{aligned}$$

So,

$$\frac{\partial f}{\partial x_i} = \lambda \frac{\partial g}{\partial x_i} \quad \forall i = 1, \dots, n.$$

That is,

$$\nabla f(x) = \lambda \nabla g(x).$$

Q.E.D. ■

Theorem 7.3.4 General Procedure to Solve an Extremal Problem

- Solve the equations for $x \in \mathbb{R}^n$ and $\lambda \in \mathbb{R}$:

$$\begin{cases} g(x) = c_0 \\ \nabla f(x) = \lambda \nabla g(x) \end{cases}$$

- Compare values of f at these points.

Example 7.3.5

Find extrema for the function $f(x, y) = x^2 - y^2$ subject to the constraint $x^2 + y^2 = 1$.

Solution 3.

Solve the equations:

$$\begin{cases} g(x) = c_0 \\ \nabla f(x) = \lambda \nabla g(x) \end{cases} \implies \begin{cases} x^2 + y^2 = 1 \\ \begin{bmatrix} 2x \\ -2y \end{bmatrix} = \lambda \begin{bmatrix} 2x \\ 2y \end{bmatrix} \end{cases} \implies \begin{cases} x^2 + y^2 = 1 \\ 2x = \lambda 2x \\ -2y = \lambda 2y. \end{cases}$$

- If $x = 0$, $y = \pm 1$, and $\lambda = -1$.
- If $y = 0$, $x = \pm 1$, and $\lambda = 1$.

Possible candidates: $(0, 1)$, $(0, -1)$, $(1, 0)$, and $(-1, 0)$.

- At $(0, 1)$, $f(0, 1) = 0^2 - 1^2 = -1$.
- At $(0, -1)$, $f(0, -1) = 0^2 - (-1)^2 = -1$.
- At $(1, 0)$, $f(1, 0) = 1^2 - 0^2 = 1$.
- At $(-1, 0)$, $f(-1, 0) = (-1)^2 - 0 = 1$.

Then, $(0, 1)$ and $(0, -1)$ are local minimum, and $(1, 0)$ and $(-1, 0)$ are local maximum.

□

Theorem 7.3.6 Extremal Problem with Multiple Constraints

Maximize/Minimize $f(x)$ with constraints $g_1(x) = c_1, \dots, g_m(x) = c_m$. Then, we solve

$$\begin{cases} g_1(x) = c_1 \\ \vdots \\ g_m(x) = c_m \\ \nabla f(x) = \lambda_1 \nabla g_1(x) + \dots + \lambda_m \nabla g_m(x). \end{cases}$$

8 Integration

8.1 Definition of Integration

8.1.1 Geometric Motivation. To compute the area of region under the curve $y = f(x)$.

- Form the upper and lower approximation:

$$U(f, \mathcal{P}) = \sum_{i=1}^n \sup_{I_i} f(x) \ell(I_i)$$

$$L(f, \mathcal{P}) = \sum_{i=1}^n \inf_{I_i} f(x) \ell(I_i).$$


- Form the upper and lower integral:

$$\bar{\int}_A f = \inf_{\mathcal{P}} U(f, \mathcal{P})$$

$$\underline{\int}_A f = \sup_{\mathcal{P}} L(f, \mathcal{P}).$$

8.1.2 General Formulation of Integral.

- Set-up: Let $f : A \subset \mathbb{R}^n \rightarrow \mathbb{R}$ be a bounded function on a bounded set A .
- Goal: define the volume of the region under the surface $y = f(x)$ (or the integral $\int_A f \, dx$).
- Step 1: choose a rectangle $B = [a_1, b_1] \times [a_2, b_2] \times \cdots \times [a_n, b_n]$ that contains A . Extend f s.t. $f(x) = 0$ when $x \notin A$.



Then, the volume over A is the same as the volume over B . That is,

$$\int_A f(x) \, dx = \int_B f(x) \, dx.$$

- Step 2: partition B : divide slides of B into sub-intervals to obtain a partition P , collection of smaller rectangles.

- Step 3: Form upper and lower sums:

$$U(f, \mathcal{P}) = \sum_{R \in \mathcal{P}} \underbrace{\sup_R f(x)}_{\text{height}} \cdot \underbrace{v(R)}_{\text{base}} \quad (\text{Upper Sum of } f \text{ w.r.t. } \mathcal{P})$$

$$L(f, \mathcal{P}) = \sum_{R \in \mathcal{P}} \inf_R f(x) \cdot v(R) \quad (\text{Lower Sum of } f \text{ w.r.t. } \mathcal{P})$$

- Step 4: Form upper and lower integrals:

$$\int_A^{\bar{}} f = \inf_{\mathcal{P}} (U(f, \mathcal{P})) \quad \text{and} \quad \int_A^{\underline{}} f = \sup_{\mathcal{P}} (L(f, \mathcal{P})).$$

- Observation:

$$L(f, \mathcal{P}) \leq \text{real volume} \leq U(f, \mathcal{P}) \implies \int_A^{\underline{}} f \leq \text{real volume} \leq \int_A^{\bar{}} f.$$

- **Definition 8.1.3 (Integrable).** We say f is *Riemann integrable* if

$$\int_A^{\underline{}} f = \int_A^{\bar{}} f.$$

The integral of f on the set A is defined as $\int_A f(x) dx = \int_A^{\underline{}} f = \int_A^{\bar{}} f$. Sometimes, the integral is also written as $\int_A f$ or $\int_A f(x) dx_1 dx_2 \cdots dx_n$.

Theorem 8.1.4 Equivalent Conditions for Integrability

Suppose $f : A \subset \mathbb{R}^n \rightarrow \mathbb{R}$ is bounded and A and B are bounded. Let B be a rectangle in \mathbb{R}^n . Assume $f(x) = 0$ for $x \notin A$. Then, the following are equivalent conditions for f to be integrable:

- (Riemann's Condition): $\forall \varepsilon > 0, \exists$ partition \mathcal{P}_ε (of B) s.t.

$$0 \leq U(f, \mathcal{P}_\varepsilon) - L(f, \mathcal{P}_\varepsilon) < \varepsilon.$$

- (Darboux's Condition): \exists a number I s.t. $\forall \varepsilon > 0, \exists \delta > 0$ s.t.

1. \mathcal{P} is any partition of B into rectangles B_1, B_2, \dots, B_N with side length less than δ , and
2. If $x_1 \in B_1, x_2 \in B_2, \dots, x_N \in B_N$, then we have

$$\left| \sum_{i=1}^N f(x_i) v(B_i) - I \right| < \varepsilon.$$

Remark 8.1 • The number I is the value of the integral

- $\sum_{i=1}^N f(x_i)v(B_i)$ is called the Riemann sum of f w.r.t. \mathcal{P} .
- Interpretation: Darboux's condition says that when the partition is fine enough (side length $< \delta$), then the Riemann sum is a good approximation of the integral.

► **Proof 1 of Equivalent Conditions for Integrability**

Step 1 f integrable \implies Riemann's Condition

Given $\varepsilon > 0$, need to find a partition \mathcal{P}_ε s.t. $U(f, \mathcal{P}_\varepsilon) - L(f, \mathcal{P}_\varepsilon) < \varepsilon$.

Since

$$\int_A f = \inf_{\mathcal{P}} U(f, \mathcal{P}),$$

by definition of infimum,

$$\exists \mathcal{P}_1 \text{ s.t. } U(f, \mathcal{P}_1) < \int_A f + \frac{\varepsilon}{2}.$$

Similarly,

$$\exists \mathcal{P}_2 \text{ s.t. } L(f, \mathcal{P}_2) > \int_A f - \frac{\varepsilon}{2}.$$

Let $\mathcal{P}_\varepsilon = \mathcal{P}_1 \cup \mathcal{P}_2$ (partition refinement). Then, \mathcal{P}_ε is a refinement of \mathcal{P}_1 and \mathcal{P}_2 . Therefore,

$$U(f, \mathcal{P}_\varepsilon) \leq U(f, \mathcal{P}_1) < \int_A f + \frac{\varepsilon}{2}, \quad \text{and} \quad L(f, \mathcal{P}_\varepsilon) \geq L(f, \mathcal{P}_2) > \int_A f - \frac{\varepsilon}{2}.$$

Hence,

$$\begin{aligned} U(f, \mathcal{P}_\varepsilon) - L(f, \mathcal{P}_\varepsilon) &\leq U(f, \mathcal{P}_1) - L(f, \mathcal{P}_2) \\ &< \int_A f + \frac{\varepsilon}{2} - \int_A f + \frac{\varepsilon}{2} \\ &= \int_A f - \int_A f + \varepsilon \\ &= 0 + \varepsilon \\ &= \varepsilon. \quad \square \end{aligned} \quad [f \text{ integrable}]$$

Step 2 Riemann's Condition $\implies f$ integrable

By Assumption, $\forall \varepsilon > 0$, \exists partition \mathcal{P}_ε s.t.

$$U(f, \mathcal{P}_\varepsilon) - L(f, \mathcal{P}_\varepsilon) < \varepsilon.$$

Since $\int_A f = \inf_{\mathcal{P}} U(f, \mathcal{P})$, we have

$$\int_A f \leq U(f, \mathcal{P}_\varepsilon).$$

Similarly, we have $\int_A f \geq L(f, \mathcal{P}_\varepsilon)$. Then,

$$0 \leq \int_A f - \int_A f \leq U(f, \mathcal{P}_\varepsilon) - L(f, \mathcal{P}_\varepsilon) < \varepsilon.$$

Thus,

$$\int_A f = \int_A f \implies f \text{ is integrable. } \square$$

Step 3 **Darboux's Condition \implies Integrability**

Let I be the number in Darboux's condition.

$$\text{WTS: } \int_A f = I = \int_A f.$$

Claim 8.1.5 $\forall \varepsilon > 0, \exists$ partition \mathcal{P} s.t.

$$|L(f, \mathcal{P}) - I| < \varepsilon \quad (\star)$$

Scratch:

$$\begin{aligned} |L(f, \mathcal{P}) - I| &< \underbrace{\left| L(f, \mathcal{P}) - \sum_{i=1}^N f(x_i)v(B_i) \right|}_{=\sum_{i=1}^N \left| \inf_{B_i} f(x_i) - f(x_i) \right| v(B_i)} + \underbrace{\left| \sum_{i=1}^N f(x_i)v(B_i) - I \right|}_{< \frac{\varepsilon}{2}, \text{ by Darboux}} \\ &< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon. \end{aligned}$$

So, we will make

$$\left| \inf_{B_i} f(x_i) - f(x_i) \right| < \frac{\varepsilon}{2v(B_i)N}$$

since we want $\frac{\varepsilon}{2}$ eventually. Then,

$$|L(f, \mathcal{P}) - I| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon.$$

Given $\varepsilon > 0$. By Darboux's condition, $\exists \delta > 0$ s.t. $\forall \mathcal{P} = \{B_1, B_2, \dots, B_N\}$ with sides $< \delta$, we have

$$\left| \sum_{i=1}^N f(x_i)v(B_i) - I \right| < \frac{\varepsilon}{2}.$$

for any $x_i \in B_i$, where $i = 1, \dots, N$.

To prove (\star) , we can choose $x_i \in B_i$ s.t.

$$0 \leq f(x_i) - \inf_{B_i} f(x_i) < \frac{\varepsilon}{2v(B_i)N}.$$

Then, it follows that

$$\begin{aligned}
 |L(f, \mathcal{P}) - I| &< \left| L(f, \mathcal{P}) - \sum_i f(x_i)v(B_i) \right| + \left| \sum_i f(x_i)v(B_i) - I \right| \\
 &< \sum_{i=1}^N \left| \inf_{B_i} f(x_i) - f(x_i) \right| v(B_i) + \frac{\varepsilon}{2} \\
 &< \sum_{i=1}^N \frac{\varepsilon}{2N \cdot \cancel{v(B_i)}} \cdot \cancel{v(B_i)} + \frac{\varepsilon}{2} \\
 &= \cancel{N} \cdot \frac{\varepsilon}{2\cancel{N}} + \frac{\varepsilon}{2} \\
 &= \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon \implies (*)
 \end{aligned}$$

Furthermore, $(*) \implies L(f, \mathcal{P}) > I - \varepsilon \quad \forall \varepsilon > 0$. So,

$$\int_A f = \sup_{\mathcal{P}} L(f, \mathcal{P}) \geq I.$$

Similarly, $\forall \varepsilon > 0, \exists \mathcal{P}$ s.t. $|U(f, \mathcal{P}) - I| < \varepsilon \implies U(f, \mathcal{P}) < I + \varepsilon$. Then,

$$\int_A f = \inf_{\mathcal{P}} U(f, \mathcal{P}) \leq I.$$

So, it must be


$$\int_A f = \int_A f = I.$$

Step 4 Integrability \implies Darboux's Condition (Scratch)

- Given $\varepsilon > 0, \exists \mathcal{P}$ s.t.

$$I - \frac{\varepsilon}{2} < L(f, \mathcal{P}) \leq \sum_i f(x_i)v(B_i) \leq U(f, \mathcal{P}) < I + \frac{\varepsilon}{2}.$$

- Given partition $\mathcal{P}, \exists \delta > 0$ s.t. for any partition \mathcal{P}' with side length $< \delta$, the sum of volumes of sub-rectangles in \mathcal{P}' that are not completely/entirely contained in a sub-rectangle in \mathcal{P} is less than ε .



Q.E.D. ■

Example 8.1.6 An Exercise

Compute the upper and lower sums for $\int_0^1 x \, dx$ over special partition \mathcal{P} :

$$\mathcal{P} = \left\{ 0, \frac{1}{n}, \frac{2}{n}, \dots, \frac{n-1}{n} \right\}.$$

8.2 Criterion for Integrability

Question: When is f integrable? How can we tell from other properties?

Short Answer: f is integrable when the set of discontinuity is “small.”

8.2.1 Measure Zero: How to Measure the Size of a Set

Definition 8.2.1 (Volume of A). Given a bounded set $A \subset \mathbb{R}^n$, define *characteristic function* of A by

$$\mathbb{1}_A(x) = \begin{cases} 1 & \text{if } x \in A \\ 0 & \text{if } x \notin A \end{cases}.$$

We say that A has *volume* (or *Jordan measurable*) if $\mathbb{1}_A(x)$ is integrable on A . We write

$$v(A) = \int_A \mathbb{1}_A(x) \, dx.$$

Remark 8.2 When $n = 1$, $v(A)$ is the length of A . When $n = 2$, $v(A)$ is the area of A .

Fact: A set has volume 0 (i.e., $v(A) = 0$) $\iff \forall \varepsilon > 0, \exists$ finite cover of A by rectangles S_1, S_2, \dots, S_N s.t.

$$\sum_{i=1}^N v(S_i) < \varepsilon.$$

Proof 1. Suppose $v(A) = \int_A \mathbb{1}_A(x) \, dx = 0$. Then, $\forall \varepsilon > 0, \exists$ partition $\mathcal{P} = \{\mathcal{P}_1, \dots, \mathcal{P}_N\}$ of B s.t.

$$U(\mathbb{1}_A(x), \mathcal{P}) < I + \varepsilon = \varepsilon.$$

$$\implies \sum_{\mathcal{P}_j \cap A \neq \emptyset} \underbrace{\sup_{\mathcal{P}} \mathbb{1}_A(x)}_{=1} \cdot v(\mathcal{P}_j) = \sum_{\mathcal{P}_j \cap A \neq \emptyset} v(\mathcal{P}_j) < \varepsilon.$$

Note that $\{\mathcal{P}_j \mid \mathcal{P}_j \cap A \neq \emptyset\}$ is a finite cover of A .

Q.E.D. ■

Definition 8.2.2 (Measure Zero Set). A set $A \subset \mathbb{R}^n$ (not necessarily bounded) is said to have measure zero, $m(A) = 0$, if $\forall \varepsilon > 0, \exists$ countable cover of A by rectangles $\{S_i\}$ s.t.

$$\sum_{i=1}^{\infty} v(S_i) < \varepsilon.$$

Remark 8.3

- $v(A) = 0 \implies m(A) = 0$
- Any finite set has volume zero.
- Any countable set has measure zero. (use geometric sum: first point covered by $\frac{\varepsilon}{2}$, second point covered by $\frac{\varepsilon}{4}$, ..., N -th point covered by $\frac{\varepsilon}{2^N}$)

Example 8.2.3

Let A be the x -axis (real line).

- If A is considered as a subset of \mathbb{R}^2 , then $m(A) = 0$.

Proof 2. To cover the x -axis, we can cover it interval by interval. But the volumes of the rectangles need to get smaller and smaller:

$$S_n = [n, n+1] \times \left[-\frac{\varepsilon}{2^{|n|+2}}, \frac{\varepsilon}{2^{|n|+2}} \right]$$

for $n = 0, \pm 1, \pm 2, \dots$

Q.E.D. ■

- However, if A is considered as a subset of \mathbb{R}^1 , then $m(A) \neq 0$.

Theorem 8.2.4

Suppose $A_i \subset \mathbb{R}^n$ with $m(A_i) = 0 \quad \forall i = 1, 2, \dots$. Then,

$$A = \bigcup_{i=1}^{\infty} A_i$$

has measure zero.

Proof 3. Given $\varepsilon > 0$ for each $i = 1, 2, \dots, m(A_i) = 0$. So, \exists rectangles $\{S_j^{(i)}\}_{j=1}^{\infty}$ s.t. $A_i \subset \bigcup_{j=1}^{\infty} S_j^{(i)}$

with $\sum_{j=1}^{\infty} v(S_j^{(i)}) < \frac{\varepsilon}{2^i}$. Then, $\{S_j^{(i)}\}_{i,j=1}^{\infty}$ is a countable collection of rectangles with

- $A = \bigcup_{i=1}^{\infty} A_i \subset \bigcup_{i=1}^{\infty} \bigcup_{j=1}^{\infty} S_j^{(i)}$

$$\bullet \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} v(S_j^{(i)}) < \sum_{i=1}^{\infty} \frac{\varepsilon}{2^i} = \varepsilon.$$

So, by definition, $m(A) = 0$.

Q.E.D. ■

Remark 8.4

- The above theorem is not true for volume zero sets. A counterexample is the rationals in $[0, 1]$. Each rational is volume zero, but their union is not volume zero as 1_A is not integrable.
- In Definition 8.2.2, we can replace “closed rectangles S_i ” by “open rectangles S_i .”

8.2.2 Lebesgue's Theorem

Theorem 8.2.5 Lebesgue's Theorem

Let A be a bounded set in \mathbb{R}^n and f be a bounded function on A . Extend f to \mathbb{R}^n by letting $f(x) = 0 \quad \forall x \notin A$. Then, f is integrable on $A \iff$ the points on which the *extended function* f is discontinuous form a set of measure zero. That is, extended f has support on A , and if D denotes the set of discontinuity of extended f , then $m(D) = 0$.

Example 8.2.6

- $A = [0, 1]$ and

$$f(x) = \begin{cases} 1 & x \text{ rational} \\ 0 & \text{o/w.} \end{cases}$$

Then, the set of discontinuity is $D = [0, 1]$, and $m(D) \neq 0$. By Lebesgue's Theorem, f is not integrable.

- $A = \{\text{rationals} \in [0, 1]\}$ and $f(x) : A \rightarrow \mathbb{R}$ by $f(x) \equiv 1$. Then, f is continuous on A , but it is not integrable on A . The extended f has $D = [0, 1]$, not measure zero. So, by Lebesgue's Theorem, f is not integrable.
- $A = \{(x, y) \mid x^2 + y^2 < 1\} \subset \mathbb{R}^2$ and $f(x) : A \rightarrow \mathbb{R}$ by

$$f(x, y) = \begin{cases} x^2 + \sin\left(\frac{1}{y}\right) & y \neq 0 \\ x^2 & y = 0. \end{cases}$$

Then, the set of discontinuity is $D = [-1, 0] \times [1, 0] + \partial A$. Then, $m(D) = 0$ in \mathbb{R}^2 . So, by Lebesgue's Theorem, f is integrable on A .

Corollary 8.2.7 of Lebesgue's Theorem:

- A bounded set $A \subset \mathbb{R}^n$ has volume $\iff \partial A$ has measure 0.

Proof 4. Assume $v(A)$ exists. Then, $\mathbb{1}_A(x)$ is integrable. So, the set of discontinuity of extended $\mathbb{1}_A(x)$ is $D = \partial A$. By Lebesgue's Theorem, $f = \mathbb{1}_A(x)$ is integrable $\iff m(\partial A) = 0$.

Q.E.D. ■

- Let $A \subset \mathbb{R}^n$ be a bounded set with volume. If $f : A \rightarrow \mathbb{R}$ is bounded and has only a (finite or) countable number of discontinuity, then f is integrable.

Proof 5. Denote the set of discontinuity of f on A as M . The set of discontinuity of the extended f will be $D \subset \partial A \cup M$. Since A has volume, by the previous Corollary, we know $m(\partial A) = 0$. Since M is countable, $m(M) = 0$. Then, $m(\partial A \cup M) = 0 \implies D \subset \partial A \cup M$ has measure zero. By Lebesgue's Theorem, f is integrable.

Q.E.D. ■

► Proof 6 of Lebesgue's Theorem**Step 1 Preparation and Reduction**

- The set-up: Fix a rectangle $B \supset A$ (so $\text{cl}(A) \subset \int(B)$) and define $g : B \rightarrow \mathbb{R}$ by

$$g(x) = \begin{cases} f(x), & x \in A \\ 0, & x \notin A. \end{cases}$$

Let D denote the set of discontinuity of $g(x)$. That is,

$$D = \{x \in B \mid g(x) \text{ is not continuous at } x\}.$$

Need to show: f integrable on $A \iff m(D) = 0$.

- How to quantify discontinuity?

1. **Definition 8.2.8 (Oscillation).** The *oscillation* of a function $h(x)$ at a point x_0 is

$$\mathcal{O}(h, x_0) = \inf \left\{ \sup \{ |h(x_2) - h(x_1)| : x_1, x_2 \in U \} : U \text{ is a neighborhood of } x_0 \right\},$$

where $\mathcal{O}(f, U) = \sup \{ |h(x_2) - h(x_1)| : x_1, x_2 \in U \}$ is the oscillation in a neighborhood U , and \inf takes over all possible neighborhoods of x_0 .

2. **Claim 8.2.9** h is continuous at $x_0 \implies \mathcal{O}(h, x_0) = 0$.


Proof. h is continuous at $x_0 \implies \forall \varepsilon > 0, \exists \delta > 0$ s.t.

$$|x - x_0| < \delta \implies |h(x) - h(x_0)| < \frac{\varepsilon}{2}.$$

For $U = \{|x - x_0| < \delta\} \cap A$,

$$\begin{aligned} x_1, x_2 \in U &\implies |h(x_2) - h(x_1)| \leq |h(x_2) - h(x_0)| + |h(x_0) - h(x_1)| \\ &< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon. \end{aligned}$$

Then, $\mathcal{O}(h, U) < \varepsilon \implies \mathcal{O}(h, x_0) = 0$. \square



Step 2 (\Leftarrow) Assume $m(D) = 0$. Prove g is integrable.

We will show: g satisfies Riemann condition.

- Set up:

Fix $\varepsilon > 0$. Let $D_\varepsilon = \{x \in B \mid \mathcal{O}(g, x) > \varepsilon\}$. Then, $D_\varepsilon \subset D$. So, $m(D_\varepsilon) = 0$.

By Definition, \exists collection of open rectangles $|B_i|$ s.t.

$$D_\varepsilon \subset \bigcup_i B_i \quad \text{and} \quad \sum_i v(B_i) < \varepsilon.$$

Claim 8.2.10 D_ε is closed (and hence compact).

Proof. (Sketch) D_ε contains all its limits points. That is,

$$x_n \in D_\varepsilon, \{x_n\} \rightarrow x \implies x \in D_\varepsilon.$$

Assume, for the sake of contradiction,

$$x \notin D_\varepsilon \implies \mathcal{O}(g, x) < \varepsilon.$$

But $\mathcal{O}(g, x_n) \geq \varepsilon$, we can derive a contradiction from there. \square


Since D_ε is compact, it has a finite subcover:

$$\{B_1, B_2, \dots, B_N\} \text{ s.t. } \sum_{i=1}^N v(B_i) < \varepsilon.$$

- Initial Partition of B :

Construct a partition \mathcal{P} from $\{B_i\}_{i=1}^N$ s.t. each rectangle $S \in \mathcal{P}$ is either:

1. disjoint from D_ε , or
2. its interior is contained in one of the B_i 's.



The way to construct \mathcal{P} is to extend the sides of B_i to form a partition on B .

Let $C_1 = \{S \in \mathcal{P} : \text{int}(S) \text{ is contained in one of } B_i\}$ and $C_2 = \{S \in \mathcal{P} : S \cap D_\varepsilon = \emptyset\}$.

- Refinement of \mathcal{P}

Fix $S \in C_2$, $S \cap D_\varepsilon = \emptyset \implies \mathcal{O}(g, x) < \varepsilon \quad \forall x \in S$. Then, $\forall x \in S, \exists$ neighborhood U_x s.t.

$$\sup \{|g(x_1) - g(x_2)| : x_1, x_2 \in U_x\} < \mathcal{O}(g, x) + \delta,$$

where $\delta = \frac{1}{2}(\varepsilon - \mathcal{O}(g, x))$. Then,

$$\sup_{U_x} g - \inf_{U_x} g < \mathcal{O}(g, x) + 2\delta = \varepsilon.$$

Denote $M_{U_x}(g) = \sup_{U_x} g$ and $m_{U_x}(g) = \inf_{U_x} g$. Then,

$$\boxed{M_{U_x}(g) - m_{U_x}(g) < \varepsilon} \quad (\star)$$

Since S is compact and $S \subset \bigcup_{x \in S} U_x$.

$\implies \exists$ finite collection of neighborhoods $\{U_{x_i}\}$ that covers S . Partition S so that each rectangle is contained in some U_{x_i} . Do this partition for each $S \in C_2$, and we obtain a refinement of \mathcal{P} , denoted by \mathcal{P}' .

- Verify Riemann's condition for \mathcal{P}' :

Note that

$$\begin{aligned}
 U(g, \mathcal{P}') - L(g, \mathcal{P}') &= \sum_{S' \in \mathcal{P}'} (M_{S'}(g) - m_{S'}(g))v(S') \\
 &= \sum_{S' \subset S \in C_1} (M_{S'}(g) - m_{S'}(g))v(S') + \sum_{S' \subset S \in C_2} (M_{S'}(g) - m_{S'}(g))v(S') \\
 &\leq \sum_{S' \subset S \in C_1} 2Mv(S') + \sum_{S' \subset S \in C_2} \varepsilon v(S') \quad [|g(x)| \leq M \text{ and } (\star)] \\
 &\leq 2M \subset \sum_i v(B_i) + \varepsilon v(B) \quad [C_1 \text{ is covered by } B'_i s] < 2M\varepsilon + \varepsilon v(B) \quad [L] \\
 &= \varepsilon(2M + v(B)).
 \end{aligned}$$

In summary, given $\varepsilon > 0$, \exists partition \mathcal{P}' s.t.

$$U(g, \mathcal{P}') - L(g, \mathcal{P}') < \varepsilon(2M + v(B)).$$

So, we satisfy Riemann condition. \square

Step 3 $(\Rightarrow) f \text{ is integrable} \Rightarrow m(D) = 0$.

For $n = 1, 2, \dots$, let

$$D_{1/n} = \left\{ x \in D \mid \mathcal{O}(g, x) \geq \frac{1}{n} \right\}.$$

Then,

$$D = \bigcup_{i=1}^{\infty} D_{1/n}.$$

Need to show: $m(D_{1/n}) = 0 \quad \forall n$.

Fix $n \geq 1$. For any partition \mathcal{P} , write

$$D_{1/n} = S_1 \cup S_2,$$

where

$$S_1 = \{x \in D_{1/n} \mid x \text{ is on the boundary of some } S \in \mathcal{P}\}$$

and

$$S_2 = \{x \in D_{1/n} \mid x \in \text{int}(S) \text{ for some } S \in \mathcal{P}\}.$$

Then, $m(S_1) = 0$. We need to show $m(S_2) = 0$.

Given $\varepsilon > 0$. By Riemann's condition, \exists partition \mathcal{P} s.t.

$$\sum_{S \in \mathcal{P}} (M_S(g) - m_S(g))v(S) < \frac{\varepsilon}{n}.$$

Let C denote the collection of all $S \in \mathcal{P}$ s.t. $D_{1/n} \cap \text{int}(S) \neq \emptyset$. Then, C covers S_2 and for any $S \in C$,

$$M_S(g) - m_S(g) \geq \mathcal{O}(g, x) \geq \frac{1}{n}.$$

Thus,

$$\sum_{S \in C} (M_S(g) - m_S(g))v(S) \leq \sum_{S \in \mathcal{P}} (M_S(g) - m_S(g))v(S) < \frac{\varepsilon}{n}.$$

Since

$$\sum_{S \in C} (M_S(g) - m_S(g))v(S) \geq \sum_{S \in C} \frac{1}{n}v(S) = \frac{1}{n} \sum_{S \in C} v(S),$$

we have

$$\frac{1}{n} \sum_{S \in C} v(S) \leq \sum_{S \in C} (M_S(g) - m_S(g))v(S) < \frac{\varepsilon}{n}.$$

That is,

$$\frac{1}{n} \sum_{S \in C} v(S) < \frac{\varepsilon}{n} \implies \sum_{S \in C} v(S) < \varepsilon.$$

Therefore, $m(S_2) = 0$ as well.

Since $m(S_1) = m(S_2) = 0$ and $D_{1/n} = S_1 \cup S_2$, $m(D_{1/n}) = 0 \quad \forall n$. Then,

$$m(D) = m\left(\bigcup_{i=1}^{\infty} D_{1/n}\right) = 0.$$

Q.E.D. ■

Theorem 8.2.11 Properties of Integration

Let $A, B \subset \mathbb{R}^n$ be bounded, $c \in \mathbb{R}$, and $f, g : A \rightarrow \mathbb{R}$ be integrable. Then,

- $f + g$ is integrable and $\int_a (f + g) = \int_A f + \int_A g$.
- cf is integrable and $\int_A (cf) = c \int_A f$.
- $|f|$ is integrable and $\left| \int_A f \right| \leq \int_A |f|$.
- If $f \leq g$, then $\int_A f \leq \int_A g$.
- If A has volume and $|f| \leq M$, then $\left| \int_A f \right| \leq Mv(A)$.
- **(Mean Value Theorem for Integrals):** If $f : A \rightarrow \mathbb{R}$ is continuous and A has volume and is compact and connected, then $\exists x_0 \in A$ s.t. $\int_A f(x) dx = f(x_0)v(A)$. The quantitative $\frac{1}{v(A)} \cdot \int_A f$ is called the *average* of f over A .
- Let $f : A \cup B \rightarrow \mathbb{R}$. If the sets A and B are such that $A \cap B$ has measure zero and $f|_{(A \cap B)}$, $f|_A$, and $f|_B$ are all integrable, then f is integrable on $A \cup B$ and $\int_{A \cup B} = \int_A f + \int_B f$.

8.3 Improper Integrals

Goal: Study integral of the form $\int_A f(x)$, where $f : A \subset \mathbb{R}^n \rightarrow \mathbb{R}$ is an arbitrary function and $A \subset \mathbb{R}^n$ is an arbitrary set.


Definition 8.3.1 (Integral).

- If $A \subset \mathbb{R}^n$ is bounded and f is bounded, then

$$\int_A f(x) = \bar{\int}_A f(x) = \underline{\int}_A f(x) \quad (\text{Riemann Condition})$$

- $f(x) \geq 0$ bounded and A is arbitrary, then


$$\int_A f(x) = \lim_{a \rightarrow \infty} \int_{A_a} f(x)$$



- $f(x) \geq 0$ unbounded and A is arbitrary.

For $M > 0$, define

$$f_M(x) = \begin{cases} f(x) & \text{for } f(x) \leq M \\ 0 & \text{o/w.} \end{cases}$$



Then,

$$\int_A f(x) = \lim_{M \rightarrow \infty} \int_A f_M(x).$$

- f is arbitrary and A is arbitrary.

Let

$$f^+(x) = \begin{cases} f(x) & f(x) \geq 0 \\ 0 & f(x) < 0, \end{cases} \quad \text{and} \quad f^-(x) = \begin{cases} 0 & f(x) \geq 0 \\ -f(x) & f(x) < 0. \end{cases}$$

Remark 8.5 1. $f^+(x)$ is the positive part of f and $f^-(x)$ is the negative part of f .

2. $f^+, f^- \geq 0$.

3. $f(x) = f^+(x) - f^-(x)$. We can write any function as the difference of two non-negative functions.

$$4. |f(x)| = f^+(x) + f^-(x).$$

So, f is integrable on A if both f^+ and f^- are integrable on A . We write

$$\int_A f(x) = \int_A f^+(x) - \int_A f^-(x).$$

Remark 8.6 1. One can show this definition preserves linearity of integral from bounded case.
 2. **Observation:** f integrable $\implies f^+$ and f^- integrable $\implies |f| = f^+ + f^-$ is also integrable.
 However, $|f|$ integrable $\not\implies f$ integrable. For counterexample,

$$f(x) = \begin{cases} 1 & x \text{ rational} \\ -1 & x \text{ irrational} \end{cases} \quad \text{on } [0, 1].$$

$|f(x)| \equiv 1 \implies$ integrable. But f^+ , f^- , or f are not integrable.

Theorem 8.3.2 Comparison Principle

Suppose

- $0 \leq g \leq f$ on A and $\int_A f$ converges (i.e., f is integrable on A)
- g is integrable on each finite rectangle $[-a, a]^n$.

Then, g is also integrable on A , and $\int_A g \leq \int_A f$.

Remark 8.7 The second condition is crucial and cannot be removed.

Proof 1. Since $g \geq 0$ and is integrable on $[-a, a]^n$, define

$$G(a) = \int_{[-a, a]^n} g(x).$$

Then, $G(a)$ is an increasing function of a . Furthermore,

$$g \leq f \implies G(a) = \int_{[-a, a]^n} g(x) \leq \int_{[-a, a]^n} f(x) \leq \int_A f(x).$$

So,

$$\int_A g(x) = \lim_{a \rightarrow \infty} G(a) \leq \int_A f(x).$$

Q.E.D. ■

Question: When does an integrable $\int_a^b f(x)$ (one-variable function) converge? If it converges, how to compute?

Theorem 8.3.3 Integral of Functions of One-Variable

- Suppose $f : [a, \infty) \rightarrow \mathbb{R}$ is continuous with $f(x) \geq 0$. Let $F'(x) = f(x)$. Then,

$$\int_a^\infty f(x) \, dx = \lim_{b \rightarrow \infty} \int_a^b f(x) \, dx = \lim_{b \rightarrow \infty} [F(b) - F(a)].$$

- Suppose $f : (a, b] \rightarrow \mathbb{R}$ is continuous with $f(x) \geq 0$. Then,

$$\int_a^b f(x) \, dx = \lim_{\varepsilon \rightarrow 0^+} \int_{a+\varepsilon}^b f(x) \, dx.$$

Example 8.3.4

- Consider $\int_1^\infty x^p \, dx$.

Solution 2.For $b \geq 1$,

$$\int_1^b x^p \, dx = \begin{cases} \ln b & p = -1 \\ \frac{1}{p+1} (b^{p+1} - 1) & p \neq -1. \end{cases}$$

When $b \rightarrow \infty$, $\int_1^b x^p \, dx$ diverges when $p \geq -1$ and converges when $p < -1$. So,

$$\int_1^\infty x^p \, dx \text{ is divergent when } p \geq -1$$

and

$$\int_1^\infty x^p \, dx = -\frac{1}{p+1} \text{ is convergent when } p < -1.$$

□

- Consider $\int_1^\infty e^{-x^2} x^3 \, dx$.

Solution 3.

Converges by comparison.

□

Definition 8.3.5 (Conditional Convergence).

$$\int_a^\infty f(x) \, dx \text{ (conditional)} = \lim_{b \rightarrow \infty} \int_a^b f(x) \, dx.$$

Remark 8.8 (Types of Convergence) For an improper integral $\int_a^\infty f(x) \, dx$, there are three types of convergence:

- **Absolute Convergence:** $\int_a^\infty |f(x)| \, dx$ exists.
- **Conditional Convergence:** $\lim_{b \rightarrow \infty} \int_a^b f(x) \, dx$ exists.
- **Divergence.**

For general function, absolute convergence $\not\Rightarrow$ conditional convergence. For continuous function, absolute convergence is stronger, and \Rightarrow conditional convergence.

Example 8.3.6

Determine whether the integral $\int_1^\infty \frac{\cos x}{x} \, dx$ is absolute convergence, conditional convergence, or neither (divergence).

Solution 4.

- First, consider absolute convergence.

Observe that

$$\begin{aligned} \int_0^\infty \left| \frac{\cos x}{x} \right| \, dx &= \int_1^\infty \frac{|\cos x|}{x} \, dx \geq \int_{\pi/2}^{n\pi/2} \frac{|\cos x|}{x} \, dx \\ &= \sum_{k=1}^{n-1} \int_{k\pi/2}^{(k+1)\pi/2} \frac{|\cos x|}{x} \, dx \\ &\geq \sum_{k=1}^{n-1} \underbrace{\frac{1}{(k+1)\frac{\pi}{2}}}_{\text{harmonic}} \int_{k\pi/2}^{(k+1)\pi/2} |\cos x| \, dx \\ &\rightarrow \infty \quad \text{as } n \rightarrow \infty. \end{aligned}$$

So, $\int_1^\infty \left| \frac{\cos x}{x} \right| \, dx$ diverges, and thus $\int_1^\infty \frac{\cos x}{x} \, dx$ is not absolutely convergent.

- Conditional convergence:

$$\int_1^b \frac{\cos x}{x} \, dx = \left. \frac{\sin x}{x} \right|_1^b + \int_1^b \frac{\sin x}{x^2} \, dx \quad \text{[Integration by Parts]}$$

When $b \rightarrow \infty$,

$$\lim_{b \rightarrow \infty} \left. \frac{\sin x}{x} \right|_1^b = \frac{\sin 1}{1} \quad \text{converges.}$$

Further,

$$\left| \frac{\sin x}{x^2} \right| \leq \left| \frac{1}{x^2} \right| = \frac{1}{x^2} \Rightarrow \int_1^\infty \left| \frac{\sin x}{x^2} \right| \, dx \leq \int_1^\infty \frac{1}{x^2} \, dx.$$

So, $\int_1^\infty \frac{\sin x}{x^2} dx$ absolutely converges by comparison.

Then, $\int_1^b \frac{\cos x}{x} dx$ is conditional convergence.

□

8.4 Lebesgue Convergence Theorem

Goal: When do we have

$$\lim_{n \rightarrow \infty} \int_A f(x) dx = \int_A \left(\lim_{n \rightarrow \infty} f(x) \right) dx? \quad (\star)$$

Theorem 8.4.1 Lebesgue Monotone Convergence Theorem (LMCT)

Let $g_n : [0, 1] \rightarrow \mathbb{R}$ be a sequence of non-negative integrable function such that

- $g_{n+1}(x) \leq g_n(x) \quad \forall x \in [0, 1]$ (decreasing sequence)
- $\lim_{n \rightarrow \infty} g_n(x) = 0 \quad \forall x \in [0, 1]$.

Then,

$$\lim_{n \rightarrow \infty} \int_0^1 g_n(x) dx = \int_0^1 0 dx = 0.$$

Corollary 8.4.2 : Suppose $f_n(x), f(x) : [0, 1] \rightarrow \mathbb{R}$ with

- $f_n \leq f_{n+1}(x) \leq f(x) \quad \forall x \in [0, 1]$
- $f_n(x) \rightarrow f(x) \quad \forall x$.

Then,

$$\lim_{n \rightarrow \infty} \int_0^1 f_n(x) dx = \int_0^1 f(x) dx.$$

Proof 1. Apply LMCT to the sequence $g_n(x) = f(x) - f_n(x) \geq 0$.

Q.E.D. ■

Remark 8.9

- For (\star) to hold, we only need $f_n(x) \uparrow f(x)$ ($f_n(x)$ is monotone increasing and the limit of $f_n(x)$ is $f(x)$)
- The assumption that $A = [0, 1] \subset \mathbb{R}$ is not essential. Result is true for any set $A \subset \mathbb{R}^n$.
- The monotonicity assumption cannot be removed. For example:

$$g_n(x) = \begin{cases} n, & 0 < x < \frac{1}{n} \\ 0, & \text{o/w} \end{cases}$$

Then, we have $g_n(x) \rightarrow 0 \quad \forall x \in [0, 1]$. However,

$$\int_0^1 g_n(x) dx = 1 \quad \forall n \quad \text{and} \quad \int_0^1 0 dx = 0.$$

So,

$$\int_0^1 g_n dx \neq \int_0^1 0 dx,$$


and LMCT does not hold anymore.

► Proof 2 of Lebesgue Monotone Convergence Theorem

Lemma 8.4.3 : Suppose $f : [0, 1] \rightarrow \mathbb{R}$ be integrable with $|f| \leq M$ and $\int_0^1 f \geq \alpha > 0$. Then, the set

$$E = \left\{ x \in [0, 1] \mid f(x) \geq \frac{\alpha}{2} \right\}$$

contains a finite union of disjoint open intervals of total length $\geq \frac{\alpha}{4M}$.



Proof. By definition of integral, \exists partition \mathcal{P} s.t.

$$0 \leq \int_0^1 f - L(f, \mathcal{P}) < \frac{\alpha}{4}.$$

Then,

$$L(f, \mathcal{P}) > \int_0^1 f - \frac{\alpha}{4} \geq \alpha - \frac{\alpha}{4} = \frac{3\alpha}{4}.$$

Let ℓ denote the total length of the intervals I in \mathcal{P} with $I \subset E$. Then,

$$\begin{aligned} \frac{3\alpha}{4} &< L(f, \mathcal{P}) = \sum_{I \in \mathcal{P}} \left(\inf_I f(x) \right) \ell(I) \\ &= \sum_{I \in \mathcal{P} \cap E} \left(\inf_I f(x) \right) \ell(I) + \sum_{I \in \mathcal{P} \setminus E} \left(\inf_I f(x) \right) \ell(I) \\ &\leq \sum_{I \in \mathcal{P} \cap E} M \cdot \ell(I) + \sum_{I \in \mathcal{P} \setminus E} \frac{\alpha}{2} \ell(I) \\ &\leq \ell M + \frac{\alpha}{2} \cdot 1 \end{aligned} \quad \left[\text{If } I \notin E, f(x) \leq \frac{\alpha}{2} \right]$$

So, $\ell \cdot M \geq \frac{\alpha}{4} \implies \ell \geq \frac{\alpha}{4M}$. Remove endpoints from I , we get open intervals. \square

- **Step 1** Set up and Reduction:

$$0 \leq g_{n+1} \leq g_n \implies \int_0^1 g_{n+1}(x) dx \leq \int_0^1 g_n(x) dx.$$

Then, the limit exists:

$$\lim_{n \rightarrow \infty} \int_0^1 g_n(x) dx =: \lambda \geq 0.$$

Need to show: $\lambda = 0$.

Assume $\lambda > 0$, and we will derive a contradiction (with the assumption $g_n(x) \rightarrow 0 \quad \forall x \in [0, 1]$).

- **Step 2** Apply the above Lemma 8.4.3 to the cut-off function $(g_n)_M$, where $M > 0$.

$$(g_n(x))_M := \begin{cases} g_n(x), & g_n(x) \leq M \\ M, & g_n(x) > M. \end{cases}$$

Then,

$$\int_0^1 g_n(x) dx = \lim_{M \rightarrow \infty} \int_0^1 (g_n)_M.$$

Choose $M = \frac{2\lambda}{5}$ s.t.

$$0 \leq \int_0^1 (g_n - (g_n)_M) \leq \int_0^1 (g_1 - (g_1)_M) \leq \frac{\lambda}{5}.$$

Let $E_n = \left\{ x \in [0, 1] \mid g_n(x) \geq \frac{2\lambda}{5} \right\}$. Then,

1. $E_{n+1} \subset E_n$ **by monotonicity**
2. $\left\{ x \in [0, 1] \mid (g_n)_M(x) \geq \frac{\alpha}{2} \right\} \subset E_n$. **Choose α s.t. $\frac{2\lambda}{5} = \frac{\alpha}{2}$ to apply the Lemma. $\implies \alpha = \frac{4\lambda}{5}$.**

Apply Lemma 8.4.3 to $(g_n)_M$ and $\alpha = \frac{4\lambda}{5}$. Then, E_n contains a finite union of disjoint open intervals of total length

$$\ell \geq \frac{\alpha}{4M} = \frac{4\lambda}{5} \cdot \frac{1}{4M} = \frac{\lambda}{5M}$$

- **Step 3** Show that $\bigcap_{n=1}^{\infty} E_n \neq \emptyset$.

Let

$$D = \bigcup_{n=1}^{\infty} \{x \in [0, 1] \mid g_n \text{ not continuous at } x\} = \bigcup_{n=1}^{\infty} D_n.$$

Since g_n is integrable, we have $m(D_n) = 0$. So,

$$m(D) = m\left(\bigcup_{n=1}^{\infty} D_n\right) = 0.$$

That is, D is covered by U , a countable union of open intervals of total length $< \varepsilon = \frac{\lambda}{5M}$.

By Step 2, $E_n \not\subset U$.

Claim 8.4.4 $\text{cl}(E_n) \subset E_n \cup U$.

Proof. In fact, if $x_0 \in \text{cl}(E_n) \setminus E_n$, then [WTS: $x_0 \in U$]

$$g_n(x_0) < \frac{2\lambda}{5} \implies g_n \text{ is not continuous at } x_0.$$

Suppose $x_0 \in \text{cl}(E_n) \implies \exists x_k \in E_n$ s.t. $x_k \rightarrow x_0$ as $k \rightarrow \infty$. Also, $g_n(x_k) \geq \frac{2\lambda}{5}$, but $g_n(x_0) < \frac{2\lambda}{5} \implies g_n(x_k) \neq g_n(x_0) \implies \text{discontinuous}$

So, $x_0 \in D_n$, and thus $x_0 \in U$. So, this Claim 8.4.4 is true. \square

Note, let $F_n = \text{cl}(E_n) \setminus U$. Then,

1. F_n is compact
2. $F_n \subset E_n$ (by Claim 8.4.4)

So, by the nested set property: $\bigcap_{n=1}^{\infty} F_n \neq \emptyset$. As $F_n \subset E_n$, we further have $\bigcap_{n=1}^{\infty} E_n \neq \emptyset$.

Let $x_0 \in \bigcap_{n=1}^{\infty} E_n$, then $g_n(x_0) \geq \frac{2\lambda}{5}$. Then, $\lim_{n \rightarrow \infty} g_n(x_0) \neq 0$. *This derives a contradiction with the second assumption in LMCT (i.e., $g_n(x) \rightarrow 0$). So, $\lambda > 0$ is impossible, and it must be that $\lambda = 0$.

Q.E.D. \blacksquare

Corollary 8.4.5: Let $g_n : A \rightarrow \mathbb{R}$ be integrable and non-negative. Assume

$$g(x) = \sum_{n=1}^{\infty} g_n(x)$$

is also integrable. Then,

$$\int_A g(x) = \int_A \sum_{n=1}^{\infty} g_n(x) = \sum_{n=1}^{\infty} \int_A g_n(x).$$

Proof3. Let $f_n(x) = \sum_{k=1}^n g_k(x)$, the partial sum.

Then,

$$\int_A f_n(x) = \int_A \sum_{k=1}^n g_k(x) = \sum_{k=1}^n \int_A g_k(x) \quad [\text{property of integral}]$$

As $n \rightarrow \infty$, $f_n \rightarrow g(x)$, and $f_{n+1} \geq f_n$ (g_n is non-negative). Then, apply Corollary 8.4.2, we have

$$\int_A g(x) = \sum_{n=1}^{\infty} \int_A g_n(x).$$

Q.E.D. \blacksquare

9 Computing Integrals

Question: In practice, how do we compute the integral $\int_A f(x) dx$?

- In \mathbb{R}^1 : Fundamental Theorem of Calculus.

$$\int_a^b f(x) dx = F(x) \Big|_a^b = F(b) - F(a).$$

- In \mathbb{R}^n : Reduce to \mathbb{R}^1 case by *Fubini's Theorem*. Or, use *change of variable* (substitution first), and then use Fubini's Theorem.

9.1 Fubini's Theorem

Theorem 9.1.1 Fubini's Theorem

Let $A = \{(x, y) \mid a \leq x \leq b, c \leq y \leq d\}$ be a rectangle in \mathbb{R}^2 and $f : A \rightarrow \mathbb{R}$ be integrable. Suppose for each fixed $x \in [a, b]$, the following integral exists:

$$g(x) = \int_c^d f(x, y) dy.$$

Then, $g(x)$ is integrable on $[a, b]$, and

$$\int_A f(x, y) = \int_a^b g(x) dx = \int_a^b \left(\int_c^d f(x, y) dy \right) dx.$$

Corollary 9.1.2: If $f : A \rightarrow \mathbb{R}$ is continuous, then

$$\int_A f(x, y) = \int_a^b \left(\int_c^d f(x, y) dy \right) dx \stackrel{\text{symmetry}}{=} \int_c^d \left(\int_a^b f(x, y) dx \right) dy.$$

Corollary 9.1.3 Generalization: Let A be a region given by $A = \{(x, y) \mid a \leq x \leq b, \varphi(x) \leq y \leq \psi(x)\}$, where φ and ψ are continuous. If $f : A \rightarrow \mathbb{R}$ is continuous, then

$$\int_A f(x, y) = \int_a^b \left(\int_{\varphi(x)}^{\psi(x)} f(x, y) dy \right) dx.$$


Remark 9.1

- The roles of x and y can be interchanged.
- Results are true in higher dimensions. For example, let $C = A \times B \subset \mathbb{R}^{n+m}$, where $A \subset \mathbb{R}^n$ and $B \subset \mathbb{R}^m$. Fix $x \in A$ and $y \in B$. Then,

$$\int_{A \times B} f = \int_A \left(\int_B f(x, y) dy \right) dx.$$

Example 9.1.4 Computing Integral

Compute $\int_A (x + y) \, dx \, dy$, where A is the following region:

**Solution 1.**

$$\begin{aligned}
 \int_A (x + y) \, dx \, dy &= \int_0^1 \left(\int_0^x (x + y) \, dy \right) dx \\
 &= \int_0^1 \left(xy + \frac{1}{2}y^2 \right) \Big|_0^x dx \\
 &= \int_0^1 x^2 + \frac{1}{2}x^2 \, dx \\
 &= \frac{3}{2} \cdot \frac{1}{3} x^3 \Big|_0^1 \\
 &= \frac{1}{2}.
 \end{aligned}$$

□

► Proof 2 of Fubini's Theorem

- Let $g(x) = \int_c^d f(x, y) \, dy$. **WTS:** (1) g is integrable on $[a, b]$, and (2) $\int_a^b g \, dx = \int_A f$. We will compute the upper and lower sums of f and g .
- Fix any partition \mathcal{P}_A of A , where $\mathcal{P}_A = \{S_{i,j}\}_{i,j}$, where $S_{i,j} = v_i \times w_j$. Then, \mathcal{P}_A induces a partition of $[a, b]$, where $\mathcal{P}_{[a,b]} = \{v_i\}_i$ and a partition of $[c, d]$, $\mathcal{P}_{[c,d]} = \{w_j\}_j$.
- Next, estimate the lower sum $L(f, \mathcal{P}_A)$:

$$\begin{aligned}
 L(f, \mathcal{P}_A) &= \sum_{i,j} \underbrace{\inf_{x \in S_{i,j}} f(x)}_{\text{denote as } m_{i,j}(f)} v(S_{i,j}) \\
 &= \sum_{i,j} m_{i,j}(f) v(v_i \times w_j) \\
 &= \sum_{i,j} m_{i,j}(f) v(v_i) \cdot v(w_j).
 \end{aligned}$$

Key Observation:

$$\inf \{f(x, y) \mid (x, y) \in v_i \times w_j\} \leq \underbrace{\inf \{f(x, y) : y \in w_j\}}_{\text{fix } x, \text{ allow } y \text{ to vary}} \quad \forall x \in v_i.$$

Denote $\inf \{f(x, y) \mid y \in w_j\} = m_j(f, x)$. Then, for any fixed $x \in [a, b]$,

$$\begin{aligned} m_{i,j}(f) &\leq m_j(f, x) \\ m_{i,j}(f)v(w_j) &\leq m_j(f, x)v(w_j) \\ \sum_j m_{i,j}(f)v(w_j) &\leq \underbrace{\sum_j m_j(f, x) \cdot v(w_j)}_{\text{lower sum of } f(x,y) \text{ in the} \\ &\quad \text{variable } y \text{ w.r.t. partition } \mathcal{P}_{[c,d]}} \\ &= L(f(x, y), \mathcal{P}_{[c,d]}) \\ &\leq \int_c^d f(x, y) dy = g(x) \quad \forall x. \end{aligned}$$

Thus,

$$\begin{aligned} \sum_j m_{i,j}(f)v(w_j) &\leq \inf_{v_i} g(x) \\ \sum_j m_{i,j}(f)v(w_j)v(v_i) &\leq \inf_{v_i} g(x)v(v_i) \\ \sum_i \sum_j m_{i,j}(f)v(w_j)v(v_i) &\leq \sum_i \inf_{v_i} g(x)v(v_i) \\ \underbrace{\sum_{i,j} m_{i,j}(f)v(w_j)v(v_i)}_{L(f, \mathcal{P}_A)} &\leq \underbrace{\sum_i \inf_{v_i} g(x)v(v_i)}_{L(g, \mathcal{P}_{[a,b]})} \end{aligned}$$

So,

$$L(f, \mathcal{P}_A) \leq L(g, \mathcal{P}_{[a,b]}).$$

- Similarly, we have

$$U(f, \mathcal{P}_A) \geq U(g, \mathcal{P}_{[a,b]}).$$

- Therefore, we have

$$L(f, \mathcal{P}_A) \leq L(g, \mathcal{P}_{[a,b]}) \leq U(g, \mathcal{P}_{[a,b]}) \leq U(f, \mathcal{P}_A).$$

Since f is integrable, by Riemann's condition,

$$0 \leq U(f, \mathcal{P}_A) - L(f, \mathcal{P}_A) < \varepsilon.$$

Then,

$$0 \leq U(g, \mathcal{P}_{[a,b]}) - L(g, \mathcal{P}_{[a,b]}) < \varepsilon.$$

So, g is integrable as well. Moreover,

$$\int_a^b g(x) \, dx = \int_A f.$$

Q.E.D. ■

Example 9.1.5

Compute the volume of the region

$$A = \{(x, y, z) \mid x \geq 0, y \geq 0, z \geq 0, x + y + z \leq 1\}$$

by integration.


Solution 3.

$$v(A) \int_A \mathbf{1}_A = \int_0^1 \int_0^{1-x} \int_0^{1-x-y} 1 \, dz \, dy \, dx.$$

□

9.2 Change of Variable

General Setting: $f : B \rightarrow \mathbb{R}$ bounded is an integrable function



Goal: Transform integral $\int_B f(y)$ to an integral on A .

Example 9.2.1 1D Case

$$\int f(y) \, dy = \int f(g(x)) \underbrace{g'(x) \, dx}_{dy}.$$

Theorem 9.2.2 Change of Variable Formula in Higher Dimension

Assume $J_g(x) \neq 0 \quad \forall x \in A$. If $f : B \rightarrow \mathbb{R}$ is bounded and integrable on $B = g(A)$, then $f \circ g(x) \cdot (J_g(x))$ is integrable on A , and

$$\int_B f(y) \, dy = \int_A f(g(x)) \cdot \underbrace{|J_g(x)| \, dx}_{dy}.$$

Proof 1. (Sketch)

- Change of volume under linear map:

Let $\mathbf{L} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear map given by

$$\mathbf{L} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}.$$

Denote $y = \mathbf{L}x$. Then,

$$v(L(s)) = \left| \det \begin{bmatrix} a & b \\ c & d \end{bmatrix} \right| \cdot v(S).$$

- Linear approximation of $g : A \rightarrow B$:

Fix $x_0 \in A$. Then, in a neighborhood of x_0 , g can be approximated by a linear map:

$$g(x) = g(x_0) + \mathbb{D}g(x_0)(x - x_0) + \text{error}.$$

- Conversion into integral formula:

Fix small rectangles S in A . Then, $g(S)$ is “nearly” parallelogram. So,

$$v(g(S)) \approx |J_g(x_0)|v(S).$$

Do this for each rectangle S_{ij} in a partition:

$$v(g(S_{ij})) \approx |J_g(x_{ij})|v(S_{ij}).$$

Then,

$$\begin{aligned} f(y_{ij})v(g(S_{ij})) &\approx f(g(x_{ij}))|J_g(x_{ij})|v(S_{ij}) \\ \sum f(y_{ij})v(g(S_{ij})) &\approx \sum f(g(x_{ij}))|J_g(x_{ij})|v(S_{ij}) \end{aligned}$$

Through the summation and limit process:

$$\int_B f(y) dy = \int_A f(g(x))|J_g(x)| dx.$$


Q.E.D. ■

Example 9.2.3

Evaluate the integral using the change of variables $u = x^2 - y^2$ and $v = 2xy$.

$$\int_0^1 \int_0^1 (x^2 + y^2) \sin(x^2 - y^2) dx dy.$$

1. Sketch the regions in xy -plane and uv -plane:



$$\begin{cases} u = x^2 - y^2 \\ v = 2xy \end{cases} \quad \text{and} \quad g : (u, v) \rightarrow (x, y) \implies u = x^2 - \frac{v^2}{4x^2}.$$

2. Compute the determinant: $g^{-1} : (u, v) \rightarrow (x, y)$.

$$J_{g^{-1}}(x, y) = \begin{vmatrix} \partial u / \partial x & \partial u / \partial y \\ \partial v / \partial x & \partial v / \partial y \end{vmatrix} = \begin{vmatrix} 2x & -2y \\ 2y & 2x \end{vmatrix} = 4x^2 + 4y^2.$$

So,

$$J_g = \frac{1}{J_{g^{-1}}(x, y)} = \frac{1}{4(x^2 + y^2)}$$

3. Apply the change of variable formula:

$$\begin{aligned} \int_0^1 \int_0^1 (x^2 + y^2) \sin(x^2 - y^2) \, dx \, dy &= \int_0^2 \int_{(1/4) \cdot v^2 - 1}^{1 - (1/4) \cdot v^2} (x^2 + y^2) \sin(x^2 - y^2) |J_g(x)| \, du \, dv \\ &= \int_0^2 \int_{(1/4) \cdot v^2 - 1}^{1 - (1/4) \cdot v^2} \cancel{(x^2 + y^2)} \sin(u) \frac{1}{4 \cancel{(x^2 + y^2)}} \, du \, dv \\ &= \frac{1}{4} \int_0^2 \int_{(1/4) \cdot v^2 - 1}^{1 - (1/4) \cdot v^2} \sin(u) \, du \, dv. \end{aligned}$$

Remark 9.2 (Special Coordinate Systems)

- *Polar Coordinate in \mathbb{R}^2 :*


$$\begin{cases} x = r \cos \theta \\ y = r \sin \theta \end{cases} \quad J_g(r, \theta) = r,$$

$$\implies \int_B f(x, y) \, dx \, dy = \int_A f(r \cos \theta, r \sin \theta) r \, dr \, d\theta.$$

- *Spherical Coordinate in \mathbb{R}^3 :*

$$\begin{cases} x = r \sin \varphi \cos \theta \\ y = r \sin \varphi \sin \theta \\ z = r \cos \varphi \end{cases} \quad J_g(r, \theta, \varphi) = r^2 \sin \varphi$$

$$\Rightarrow \int_B f(x, y, z) \, dx \, dy \, dz = \int_A f(r \sin \varphi \cos \theta, r \sin \varphi \sin \theta, r \cos \varphi) r^2 \sin \varphi \, dr \, d\theta \, d\varphi.$$



- *Cylindrical Coordinate in \mathbb{R}^3 :*

$$\begin{cases} x = r \cos \theta \\ y = r \sin \theta \\ z = z \end{cases} \quad J_g(r, \theta, z) = r$$

$$\Rightarrow \int_B f(x, y, z) \, dx \, dy \, dz = \int_A f(r \cos \theta, r \sin \theta, z) r \, dr \, d\theta \, dz.$$

Example 9.2.4

- Evaluate $\int_{-\infty}^{\infty} e^{-x^2} \, dx$

Solution 2.

Step 1 Evaluate integral

$$\int_{\mathbb{R}^2} e^{-x^2-y^2} \, dx \, dy$$

by polar coordinate ($x = r \cos \theta$ and $y = r \sin \theta$). Let D_R denote the circle centered at origin with radius R Then,

$$\begin{aligned} \int_{D_R} e^{-x^2-y^2} \, dx \, dy &= \int_0^{2\pi} \int_0^R e^{-r^2} r \, dr \, d\theta = \int_0^{2\pi} \left(-\frac{1}{2} e^{-r^2} \right) \Big|_0^R \, d\theta \\ &= 2\pi \left(-\frac{1}{2} e^{-R^2} + \frac{1}{2} \right) = -\pi e^{-R^2} + \pi. \end{aligned}$$

So,

$$\begin{aligned}\int_{\mathbb{R}^2} e^{-x^2-y^2} dx dy &= \lim_{R \rightarrow \infty} \int_{D_R} e^{-x^2-y^2} dx dy \\ &= \lim_{R \rightarrow \infty} \left(-\pi e^{-R^2} + \pi \right) \\ &= \pi.\end{aligned}$$

Step 2 Evaluate

$$\int_{\mathbb{R}^2} e^{-x^2-y^2} dx dy$$

by Fubini's Theorem.

Let $S_b = [-b, b] \times [-b, b] \subset \mathbb{R}^2$. Then,

$$\begin{aligned}\int_{\mathbb{R}^2} e^{-x^2-y^2} dx dy &= \lim_{b \rightarrow \infty} \int_{S_b} e^{-x^2-y^2} dx dy \\ &= \lim_{b \rightarrow \infty} \int_{-b}^b \int_{-b}^b e^{-x^2} \cdot e^{-y^2} dx dy \\ &= \lim_{b \rightarrow \infty} \left(\int_{-b}^b e^{-x^2} dx \right) \cdot \left(\int_{-b}^b e^{-y^2} dy \right) \\ &= \left(\int_{-\infty}^{\infty} e^{-x^2} dx \right)^2\end{aligned}$$

Step 3 Combine Steps 1 and 2:

$$\pi = \int_{\mathbb{R}^2} e^{-x^2-y^2} dx dy = \left(\int_{-\infty}^{\infty} e^{-x^2} dx \right)^2$$

So,

$$\int_{-\infty}^{\infty} e^{-x^2} dx = \sqrt{\pi}.$$

□

- Evaluate $\int_{\mathbb{R}^3} \frac{1}{x^2 + y^2 + z^2} dx dy dz$
- Evaluate $\int_R 2e^{x^2-y^2} dx dy dz$, where $R = \{(x, y, z) \mid x^2 + y^2 \leq 1, 1 \leq x \leq 2\}$.

10 Fourier Analysis

10.1 Introduction

General Idea: Try to decompose certain objects into simpler components.

- Algebraic Model: \mathbb{R}^n

$$x = \sum_{i=1}^n x_i e_i,$$

where e_i 's are the standard basis.

- Calculus Model: Taylor Series

$$f(x) = \sum_{n=0}^{\infty} c_n (x - a)^n.$$

- Fourier Analysis: Theory of infinite dimensional inner product space of functions.

Goal: Decompose a function $f(x)$ into a “linear combination of basis:”

$$f(x) = \sum_{n=-\infty}^{\infty} c_n \varphi_n(x).$$

Physics Motivation: Decompose complicated waves into harmonies.

10.2 Inner Product Space of Functions

10.2.1 Basic Concepts

Definition 10.2.1 (Inner Product). Let V be a complex vector space. Then, an *inner product* on V is a map $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{C}$ s.t. $\forall f, g, h \in V$ and $a, b \in \mathbb{C}$, we have

- Linearity:

$$\langle af + bg, h \rangle = a \langle f, h \rangle + b \langle g, h \rangle.$$

- Conjugate Symmetry:

$$\langle f, g \rangle = \overline{\langle g, f \rangle}.$$

- Positive Definiteness:

$$\langle f, f \rangle \geq 0 \quad \text{and} \quad \langle f, f \rangle = 0 \iff f = 0.$$

Example 10.2.2

\mathbb{C} is an inner product space under the inner product:

$$\langle z_1, z_2 \rangle = z_1 \overline{z_2}.$$

Corollary 10.2.3 Conjugate Linearity in the Second Component:

$$\langle h, af + bg \rangle = \bar{a} \langle h, f \rangle + \bar{b} \langle h, g \rangle.$$

Proof 1.

$$\begin{aligned} \langle h, af + bg \rangle &= \overline{\langle af + bg, h \rangle} && \text{[Conjugate symmetry]} \\ &= \overline{a \langle f, g \rangle + b \langle g, h \rangle} && \text{[Linearity]} \\ &= \bar{a} \langle h, f \rangle + \bar{b} \langle h, g \rangle. && \text{[Conjugate symmetry]} \end{aligned}$$

Q.E.D. ■

Definition 10.2.4 (Norm and Distance Induced by Inner Product).

- Norm:

$$\|f\| := \sqrt{\langle f, f \rangle}.$$

- Distance from f to g :


$$d(f, g) := \|f - g\|.$$

Corollary 10.2.5 Facts:

- $(V, \|\cdot\|)$ is a normed space.
- (V, d) is a metric space.

Lemma 10.2.6 Cauchy-Schwarz Inequality:

$$|\langle f, g \rangle| \leq \|f\| \cdot \|g\|$$

Proof 2.

The projection should have the smallest length:

$$\begin{aligned} 0 \leq \|f - \langle f, g \rangle g\|^2 &= \langle f, \langle f, g \rangle g, f - \langle f, g \rangle g \rangle \\ &= \langle f, f - \langle f, g \rangle g \rangle - \langle f, g \rangle \langle g, f - \langle f, g \rangle g \rangle \\ &= \langle f, f \rangle - \overline{\langle f, g \rangle} \langle f, g \rangle - \langle f, g \rangle \langle g, f \rangle + \langle f, g \rangle \overline{\langle f, g \rangle} \langle g, g \rangle \\ &= \|f\|^2 - |\langle f, g \rangle|^2 - |\langle f, g \rangle|^2 + |\langle f, g \rangle|^2 \|g\|^2. \end{aligned}$$

Normalize: let $\|g\| = 1$. Then,

$$\begin{aligned} 0 &\leq \|f\|^2 - |\langle f, g \rangle|^2 \\ |\langle f, g \rangle|^2 &\leq \|f\|^2 \\ |\langle f, g \rangle| &\leq \|f\| = \|f\| \cdot \|g\|. \end{aligned}$$

Q.E.D. ■

Definition 10.2.7 (Convergence). Suppose $f_n, f \in V$. Then, $f_n \rightarrow f$ in V if $\|f_n - f\| \rightarrow 0$ as $n \rightarrow \infty$. We call this *convergence in norm*.

10.2.2 The Space \mathcal{C} and L^2

Definition 10.2.8 (Integral of Complex Valued Functions). Suppose $f(x) = f_1(x) + \mathbf{i}f_2(x) : [a, b] \rightarrow \mathbb{C}$ be a complex-valued function, where $f_1, f_2 : [a, b] \rightarrow \mathbb{R}$. Then,

$$\int_a^b f(x) \, dx := \int_a^b f_1(x) \, dx + \mathbf{i} \int_a^b f_2(x) \, dx.$$

Definition 10.2.9 (The Space \mathcal{C} and L^2). Fix an interval $[a, b]$.

- $\mathcal{C} := \{f(x) \mid f : [a, b] \rightarrow \mathbb{C} \text{ is continuous}\}.$
- $L^2 := \left\{ f : [a, b] \rightarrow \mathbb{C} \mid \int_a^b |f(x)|^2 \, dx < \infty \right\}.$

The condition $\int_a^b |f(x)|^2 \, dx < \infty$ is called L^2 *integrable*.

Corollary 10.2.10 Facts:

- \mathcal{C} and L^2 are vector spaces. \mathcal{C} is a subspace of L^2 .
- Zero vector in \mathcal{C} : $f(x) \equiv 0$.
- Zero vector in L^2 : $f(x) = 0$ a. e. (almost everywhere).

That is, $m(\{x \in [a, b] \mid f(x) \neq 0\}) = 0$.

- $\underbrace{f_1 = f_2}_{\text{vectors}} \text{ in } L^2 \iff \underbrace{f_1(x) = f_2(x)}_{\text{function}} \text{ a. e.}$

- Inner Product:

$$\langle f, g \rangle = \int_a^b f(x) \overline{g(x)} \, dx.$$

Claim 10.2.11 With the above definition of inner product, \mathcal{C} and L^2 are inner product spaces.

10.3 Fourier Analysis on Inner Product Space

10.3.1 Geometry of an Inner Product Space

Definition 10.3.1 (Orthogonality). $f, g \in V$ are *orthogonal* (denoted as $f \perp g$) if $\langle f, g \rangle = 0$.

Definition 10.3.2 (Orthonormal Family). A family $\{\varphi_1, \varphi_2, \dots\} \subset V$ is called an *orthonormal family* if

- $\langle \varphi_i, \varphi_j \rangle = 0 \quad \forall i \neq j$
- $\text{norm} \varphi_i = 1 \quad \forall i$.

Or equivalently,

$$\langle \varphi_i, \varphi_j \rangle = \delta_{ij} = \begin{cases} 1, & \text{if } i = j \\ 0, & \text{if } i \neq j. \end{cases}$$

Example 10.3.3

In \mathbb{R}^n : $\{e_1, e_2, \dots, e_n\}$, the standard basis, is an orthonormal basis.

Theorem 10.3.4 Gram-Schmidt Process: Generate Orthonormal Family from Linear Independent Family

$$\underbrace{\{g_0, g_1, \dots\}}_{\text{linear independent}} \rightarrow \underbrace{\{\varphi_0, \dots, \varphi_1, \dots\}}_{\text{orthonormal}}$$

1. Orthogonal projection:

$$x = \sum_i c_i e_i,$$

where $c_i = \langle x, e_i \rangle$. Then, we have

$$\langle x - \langle x, e_i \rangle e_i, e_i \rangle = 0.$$

2. Inductive Process:

$$\varphi_0 = \frac{g_0}{\|g_0\|}$$

$$f_1 = g_1 - \langle g_1, \varphi_0 \rangle \varphi_0, \quad \implies \varphi_1 = \frac{f_1}{\|f_1\|}$$

\vdots

$$f_n = g_n - \sum_{i=0}^{n-1} \langle g_n, \varphi_i \rangle \varphi_i, \quad \implies \varphi_n = \frac{f_n}{\|f_n\|}.$$

10.3.2 Fourier Series and Complete Family

Definition 10.3.5 (Complete Orthonormal Family). An orthonormal family $\{\varphi_0, \varphi_1, \dots\}$ (countable) is called *complete* if each $f \in V$ can be written as

$$f = \sum_{k=0}^{\infty} c_k \varphi_k \quad (\star)$$

Remark 10.1

- The meaning of (\star) :

$$\left\| f - \sum_{k=0}^n c_k \varphi_k \right\| \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

- (\star) is called the Fourier series of f w.r.t. $\{\varphi_0, \varphi_1, \dots\}$.
- If $\{\varphi_0, \varphi_1, \dots\}$ is complete, then it is an orthonormal basis of V .

Objective: Find suitable complete orthonormal family and expand $f \in V$ into Fourier series.

Theorem 10.3.6

If f has Fourier series expansion:

$$f = \sum_{k=0}^{\infty} c_k \varphi_k,$$

then,

$$c_k = \langle f, \varphi_k \rangle \quad \text{for } k = 0, 1, \dots$$

c_k 's are called the *Fourier coefficients* of f .

Proof 1. Let

$$S_n = \sum_{k=0}^n c_k \varphi_k.$$

Then,

$$\|f - S_n\| \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Fix $m \geq 0$. Then, for any $n \geq m$,

$$\begin{aligned} c_m &\stackrel{\text{want}}{=} \langle f, \varphi_m \rangle = \langle f - S_n + S_n, \varphi_m \rangle \\ &= \langle f - S_n, \varphi_m \rangle + \langle S_n, \varphi_m \rangle \\ &= \langle f - S_n, \varphi_m \rangle + c_m \\ &= 0 + c_m \quad \text{as } n \rightarrow \infty \end{aligned}$$

[orthogonality]

[Cauchy-Schwarz]
 $\|f - S_n\| \rightarrow 0$

So, $\langle f, \varphi_m \rangle = c_m$.

Q.E.D. ■

Question: Given f and $\{\varphi_1, \varphi_2, \dots\}$, does the series

$$\sum_{k=0}^{\infty} \langle f, \varphi_k \rangle \varphi_k$$

converge to f ?

Theorem 10.3.7 Properties of Fourier Coefficients

Assume $\{\varphi_0, \varphi_1, \dots\}$ is an orthonormal family in V .

- Bessel's Inequality:

$$\sum_{k=0}^{\infty} |\langle f, \varphi_k \rangle|^2 \leq \|f\|^2.$$

- Parseval's Equality (One can View this as the Pythagorean Theorem):

If

$$f = \sum_{k=0}^{\infty} \langle f, \varphi_k \rangle \varphi_k,$$

then

$$\sum_{k=0}^{\infty} |\langle f, \varphi_j \rangle|^2 = \|f\|^2.$$

Proof 2. Let $S_n = \sum_{k=0}^n \langle f, \varphi_k \rangle \varphi_k$. Denote $c_k = \langle f, \varphi_k \rangle$.

$$\begin{aligned} \|f\|^2 &= \|f - S_n + S_n\|^2 \\ &= \langle f - S_n + S_n, f - S_n + S_n \rangle && \text{[definition]} \\ &= \|f - S_n\|^2 + \|S_n\|^2 && \text{[Linearity, } f - S_n \perp S_n] \\ \|S_n\| &= \langle S_n, S_n \rangle = \sum_{k=0}^n |c_k|^2. \end{aligned}$$

Then,

$$\|f\|^2 = \underbrace{\|f - S_n\|^2}_{\geq 0} + \sum_{k=0}^n |c_k|^2 \implies \|f\|^2 \geq \sum_{k=0}^n |c_k|^2 = \sum_{k=0}^n |\langle f, \varphi_k \rangle|^2$$

true for any n . So, we get ① by letting $n \rightarrow \infty$.

Under the assumption of ②, when $n \rightarrow \infty$, we have $\|f - S_n\|^2 \rightarrow 0$. So,

$$\|f\|^2 = \sum_{k=0}^{\infty} |c_k|^2 = \sum_{k=0}^{\infty} |\langle f, \varphi_k \rangle|^2.$$


Q.E.D. ■

Theorem 10.3.8 Best mean Approximation Theorem (BMAT)

Assume $\{\varphi_0, \varphi_1, \dots\}$ is an orthonormal family in V . For any scalars $t_0, t_1, \dots, t_n \in \mathbb{C}$, we have

$$\left\| f - \sum_{k=0}^n t_k \varphi_k \right\| \geq \left\| f - \sum_{k=0}^n \langle f, \varphi_k \rangle \varphi_k \right\|.$$

- The first sum is an arbitrary element in the plane formed by $\{\varphi_0, \dots, \varphi_n\}$.
- The second sum is the orthogonal projection of f onto the plane.

Remark 10.2 (Geometric Interpretation)

LHS \leq RHS: the shortest distance from a point f to the plane is achieved by the orthogonal projection (or, the perpendicular line).

Proof 3. Let $h_n = \sum_{k=0}^n t_k \varphi_k$. Then,

$$\begin{aligned} \|f - h_n\|^2 &= \langle f - h_n, f - h_n \rangle \\ &= \langle f, f \rangle - \langle h_n, f \rangle - \langle f, h_n \rangle + \langle h_n, h_n \rangle \\ &= \|f\|^2 - \sum_{k=0}^n t_k \overline{c_k} - \sum_{k=0}^n \overline{t_k} c_k + \sum_{k=0}^n |t_k|^2 \\ &\quad \vdots \text{ linearity} \\ &= \|f\|^2 - \sum_{k=0}^n |c_k|^2 + \sum_{k=0}^n |t_k - c_k|^2 \\ &= \|f - f_n\|^2 + \underbrace{\sum_{k=0}^n |t_k - c_k|^2}_{\geq 0}. \end{aligned}$$

So, BMAT is proven.

Q.E.D. ■

10.4 Completeness and Convergence in L^2

Theorem 10.4.1 Orthogonal Functions in L^2

Let $V = L^2([a, b])$, where $[a, b] = [0, 2\pi]$.

- Exponential family:

$$\varphi_n(x) = \frac{e^{inx}}{\sqrt{2\pi}}, \quad n = 0, \pm 1, \pm 2, \dots$$

- Trig. family:

$$\frac{1}{\sqrt{2\pi}}, \frac{\cos mx}{\sqrt{2\pi}}, \frac{\sin nx}{\sqrt{2\pi}}, \quad n, m = 1, 2, \dots$$

Claim 10.4.2 Both families are orthogonal.

Proof 1. (of exponential family)

WTS:

$$\langle \varphi_n, \varphi_m \rangle = \delta_{n,m} = \begin{cases} 1, & n = m \\ 0, & n \neq m \end{cases}$$

$$\begin{aligned} \langle \varphi_n, \varphi_m \rangle &= \int_0^{2\pi} \varphi_n(x) \overline{\varphi_m(x)} \, dx \\ &= \frac{1}{2\pi} \int_0^{2\pi} e^{inx} \cdot e^{-imx} \, dx \\ &= \frac{1}{2\pi} \int_0^{2\pi} e^{i(n-m)x} \, dx \\ &= \begin{cases} 1, & n = m \\ \frac{1}{2\pi} \cdot \frac{1}{i(n-m)} e^{i(n-m)x} \Big|_0^{2\pi} = 0, & n \neq m. \end{cases} \end{aligned}$$

Q.E.D. ■

Theorem 10.4.3 Mean Convergence Property/Completeness

The exponential family $\{\varphi_n\}_{n=-\infty}^{\infty}$ is complete in L^2

Remark 10.3 To prove this Theorem, we aim to show: any function $f(x) \in L^2$ can be represented by its Fourier series:

$$f(x) = \sum_{n=-\infty}^{\infty} \langle f, \varphi_n \rangle \varphi_n.$$

i.e.,

$$\left\| f(x) - \sum_{k=-n}^n \langle f, \varphi_k \rangle \varphi_k \right\|_{L^2} \xrightarrow{(n \rightarrow \infty)} 0.$$

Lemma 10.4.4 Stone-Weierstrass Theorem: Continuous functions can be approximated by polynomials of e^{ix} and e^{-ix} . More precisely, given $f : [0, 2\pi] \rightarrow \mathbb{C}$ continuous with $f(0) = f(2\pi)$. Then, $\forall \varepsilon > 0$,

$\exists n \geq 1$ and $c_k, k = 0, \pm 1, \dots$ s.t.

$$|f(x) - p_n(x)| < \varepsilon \quad \forall x \in [0, 2\pi],$$

where

$$p_n(x) = \sum_{k=-n}^n c_k e^{ikx},$$

a polynomial in e^{ix} and e^{-ix} .

Lemma 10.4.5: Integrable functions can be approximated by continuous functions. That is, let $f \in L^2$ and $\varepsilon > 0$ be given, \exists continuous function $g : [0, 2\pi] \rightarrow \mathbb{C}$ with $g(0) = g(2\pi)$ s.t.

$$\|f - g\| < \varepsilon.$$

► Proof 2 of Mean Convergence Property

- Step 1 Special Case:

Let f be continuous with $f(0) = f(2\pi)$. Write

$$S_n = \sum_{k=-n}^n \langle f, \varphi_k \rangle \varphi_k, \quad \text{where } \varphi_k(x) = \frac{e^{ikx}}{\sqrt{2\pi}}$$

WTS: $\|f - S_n\| \rightarrow 0$ as $n \rightarrow \infty$.

Fix $\varepsilon > 0$. By Lemma 10.4.4, we can choose $p_N(x)$ s.t.

$$|f(x) - p_N(x)| < \frac{\varepsilon}{\sqrt{2\pi}} \quad \forall x \in [0, 2\pi].$$

Then,

$$\|f - p_N\| \leq \left(\int_0^{2\pi} \left(\frac{\varepsilon}{\sqrt{2\pi}} \right)^2 \right)^{1/2} = \varepsilon.$$

Thus, $\forall n \geq N$, we have

$$\begin{aligned} \|f - S_n\| &\leq \|f - S_N\| && \text{[BMAT. } L_N \subset L_n \implies S_N \in L_n.] \\ &\leq \|f - p_N\| && \text{[BMAT. } p_N \in L_N] \\ &\leq \varepsilon. \end{aligned}$$

So, $\|f - S_n\| \rightarrow 0$ as $n \rightarrow \infty$.

- Step 2 General Case:

Fix $f \in L^2$. WTS: $f = \sum_{k=-\infty}^{\infty} \langle f, \varphi_k \rangle \varphi_k$.

By Lemma 10.4.5, \exists sequence of continuous functions $g_n : [0, 2\pi] \rightarrow \mathbb{C}$ with $g(0) = g(2\pi)$ s.t.

$$\|f - g_n\| \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

By Step 1, for each g_n , we have

$$g_n = \sum_{k=-\infty}^{\infty} \langle g_n, \varphi_k \rangle \varphi_k.$$

WTS: $\|f - S_n\| \rightarrow 0$.

Fix $\varepsilon > 0$. Choose N s.t.

$$\|f - g_N\| < \frac{\varepsilon}{3}.$$

Then, choose M s.t.

$$n \geq M \implies \|g_N - S_n(g_N)\| < \frac{\varepsilon}{3},$$

where $S_n(g_N)$ denotes the partial sum of Fourier series of g_N .

$$S_n(g_N) = \sum_{k=-n}^n \langle g_N, \varphi_k \rangle \varphi_k.$$

Thus, $\forall n \geq M$, we have

$$\begin{aligned} \|f - S_n\| &= \|S_n - S_n(g_N) + S_n(g_N) - g_N + g_N - f\| \\ &\leq \|S_n - S_n(g_N)\| + \|S_n(g_N) - g_N\| + \|g_N - f\| \\ \|S_n - S_n(g_N)\| &= \left\| \sum_{k=-n}^n \langle f, \varphi_k \rangle \varphi_k - \sum_{k=-n}^n \langle g_N, \varphi_k \rangle \varphi_k \right\| \\ &= \left\| \sum_{k=-n}^n \langle f - g_N, \varphi_k \rangle \varphi_k \right\| \\ &= \left\langle \sum_{k=-n}^n \langle f - g_N, \varphi_k \rangle \varphi_k, \sum_{k=-n}^n \langle f - g_N, \varphi_k \rangle \varphi_k \right\rangle^{1/2} \\ &= \left(\sum_{k=-n}^n |\langle f - g_N, \varphi_k \rangle|^2 \right)^{1/2} \quad \text{[Pythagorean Theorem]} \\ &\leq \|f - g_N\| < \frac{\varepsilon}{3}. \end{aligned}$$

So,

$$n \geq M \implies \|f - S_n\| < \frac{\varepsilon}{3} + \frac{\varepsilon}{3} + \frac{\varepsilon}{3} = \varepsilon.$$

Therefore,

$$\|f - S_n\| \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Q.E.D. ■

With that, these notes mark the end of a journey through the rigorous landscapes of Real Analysis. From the foundational structure of \mathbb{R} to the elegance of Fourier series in L^2 , this document reflects not only the theorems and proofs, but also the quiet persistence of curiosity.

I would like to express my sincere gratitude to:

Professor Shanshuang Yang

for his guidance, clarity, and intellectual generosity throughout MATH 411 & 412.

Jerrold E. Marsden and Michael J. Hoffman

whose text Elementary Classical Analysis served as a source of inspiration, challenge, and reflection.

My peers and friends

for the many shared ideas, problem-solving moments, and philosophical tangents.

Myself

for persistence, patience, and the willingness to wrestle with the abstract.

End of Notes

Jiuru Lyu

April 29, 2025