

Numerical differentiation and integration

JrPhy

Introduction

- Many problems are without the analytical solution, it can just be solved by some numerical methods. A few century ago, scientists calculated by hands. Now we use computers to do it, so the numerical method will be more important.

Definition of the differentiation


- Give a function $f(x)$, f is continuous in $[a,b]$, and $x \in [a,b]$
- Differential calculate a slope of a curve

$$\frac{d}{dx} f(x) = f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

- The grid is discrete in the computer, so the differential equation becomes to difference equation

$$\frac{d}{dx} f(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \rightarrow \frac{\Delta f(x)}{\Delta x} = \frac{f(x+h) - f(x)}{h}$$

Difference in computer



$$\frac{\Delta f(x)}{\Delta x} = \frac{f(x+h) - f(x)}{h}$$

- Here h is the distance of the difference grid. For example, $h = 2$, then

$$\frac{\Delta f(x)}{\Delta x} = \frac{f(x+2) - f(x)}{2} \rightarrow \frac{\Delta f[0]}{\Delta x[0]} = \frac{f[2] - f[0]}{2} \quad \frac{\Delta f[1]}{\Delta x[1]} = \frac{f[3] - f[1]}{2}$$

- It's the simplest difference, but there are more complex differential equations in our problems, I'll introduce them in the future.



Definition of the integration

- Give a function $f(x)$, f is continuous in $[a,b]$, and $x \in [a,b]$.
- Integration is the area enclosed by the function and the axis

$$\int_a^b f(x)dx = \lim_{\Delta x \rightarrow 0} \sum_{i=0}^n f(x_i)\Delta x$$

- The figure shows in right, the area of the bigger one is U , the other is L , assume the Riemannian sum of f is I , then its integral exists when

$$\bullet L \leq I \leq U$$



Integration in computer

- As the previous slide, we can use rectangular to calculate the integral, its length and width are $f[0]$ and Δx . Suppose $\Delta x = 2$, then the integral is

$$\bullet f[0]*2 + f[2]*2 + \dots + f[n]*2$$

- In program, it's convenient to use function, and set start point at a , end point at b , slice into n part, then

$$\Delta x = \frac{b-a}{n}, I = \sum f(x + \Delta x) \Delta x$$



- Obviously, the error is very high unless Δx is very small, it costs computing source, so there are so many numerical methods for Integration.

Trapezoidal method

- As the figure shows, when we connect the points of $f(x_i)$ and $f(x_{i+1})$, then the error reduces very fast, it forms a trapezoid, $f(x_i)$ and $f(x_{i+1})$ are its base, Δx is height, so

$$\Delta x = \frac{b-a}{n}, I = \sum \frac{[f(x) + f(x + \Delta x)]}{2} \Delta x$$

- Give any curve, the error of trapezoidal method is a little high, so next method we use 3 given points to find a parabola to reduce the error.



Simpson's 1/3 method


- A general form of a parabola is

$$y = ax^2 + bx + c$$

- The integration of y from $-h$ to h is

$$\begin{aligned} A &= \int_{-h}^h y dx = \int_{-h}^h (ax^2 + bx + c) dx = \left[\frac{a}{3} x^3 + \frac{b}{2} x^2 + cx + d \right]_{-h}^h \\ &= \frac{2a}{3} h^3 + 2ch = \frac{h}{3} (2ah^2 + 6c) \end{aligned}$$

- Suppose there are n point in $[a, b]$, then use the first 3 points a, x_0, x_1 , so the integration of y is




Simpson's 1/3 method

- Suppose the parabolic pass through $(-h, y_0)$, $(0, y_1)$, (h, y_2)
- $y_0 = ah^2 - bh + c$
- $y_1 = c$
- $y_2 = ah^2 + bh + c$
- Then solve the c and ah^2 , we get
- $2ah^2 + 2c = (y_0 + y_2)$, $c = y_1$

$$A = \frac{2a}{3}h^3 + 2ch = \frac{h}{3}(2ah^2 + 6c) = \frac{1}{3}(2ah^2 + 6c) = \frac{h}{3}(y_0 + 4y_1 + y_2)$$

- So $I = \frac{h}{3}(y_0 + 4y_1 + 2y_2 + 4y_3 + 2y_4 + \dots + y_n) = \frac{h}{3}(\text{first} + 4 \times \text{odd} + 2 \times \text{even} + \text{last})$



Simpson's 1/3 method

- In the summation, the factor 4 is from the coefficient, but where is from? As we mentioned before, simpson's $1/3$ method needs 3 point.
- Suppose we have n points, the 0, 1, 2 are in the group a, 2, 3, 4 are in the group b, 4, 5, 6 are in the group c, and so on. There is a point plus 2 times

Diagram illustrating the addition of two numbers represented by blocks. The first number is 141, and the second number is 141. The sum is 14242424242424241.

Example

$$I = \int_0^1 e^x dx = e - 1 \cong 1.718281828.....$$

n	result	error	n	result	error
3	1.718861	5.79E-04	12	1.639652	-7.86E-02
4	1.466244	-2.52E-01	13	1.718282	4.60E-07
5	1.718319	3.70E-05	14	1.651262	-6.70E-02
6	1.55514	-1.63E-01	15	1.718282	2.48E-07
7	1.718289	7.34E-06	16	1.659889	-5.84E-02
8	1.598074	-1.20E-01	17	1.718282	1.46E-07
9	1.718284	2.33E-06	18	1.666549	-5.17E-02
10	1.623194	-9.51E-02	19	1.718282	9.09E-08
11	1.718283	9.53E-07	20	1.671847	-4.64E-02

The result is as the table, as you can see simpson's 1/3 method is very accuracy, but there is some problem when n is even number. Because the method we derive needs 3 points for integration, it's lack of one point to do that, so the error is so high. Then how do we solve it?

Simpson's 3/8 method

- Its derivation is like 1/3 method, from quadra form

$$f(x) = ax^3 + bx^2 + cx + d$$

- 4 unknown parameters, at least 4 distinct points to find y . Here we use Lagrange interpolation to
- construct y and solve a, b, c, d .
- Suppose there are 4 distinct points
- $(x_0, f(x_0)), (x_1, f(x_1)), (x_2, f(x_2)), (x_3, f(x_3)), x_i = x_0 + ih$

Simpson's 3/8 method

$$\begin{aligned} f(x) &= \frac{(x-x_1)(x-x_2)(x-x_3)}{(x_0-x_1)(x_0-x_2)(x_0-x_3)} + \frac{(x-x_0)(x-x_2)(x-x_3)}{(x_1-x_0)(x_1-x_2)(x_1-x_3)} \\ &= \frac{(x-x_1)(x-x_0)(x-x_3)}{(x_2-x_1)(x_2-x_0)(x_2-x_3)} + \frac{(x-x_0)(x-x_1)(x-x_2)}{(x_3-x_1)(x_3-x_2)(x_3-x_0)} \\ &= \frac{(x-x_1)(x-x_2)(x-x_3)}{(-h)(-2h)(-3h)} + \frac{(x-x_0)(x-x_2)(x-x_3)}{(h)(-h)(-2h)} \\ &= \frac{(x-x_1)(x-x_0)(x-x_3)}{(h)(2h)(-h)} + \frac{(x-x_0)(x-x_1)(x-x_2)}{(2h)(h)(3h)} \end{aligned}$$

Simpson's 3/8 method

- After integration, the coefficient are
- $a = 3/8, b = 9/8, c = 9/8, d = 3/8$

$$I = \frac{3h}{8} (y_0 + 3y_1 + 3y_2 + 2y_3 + 3y_4 + \dots + y_n)$$

$$= \frac{3h}{8} (\textit{first} + 3 \times \textit{others} + 2 \times (\textit{multiple of 3}) + \textit{last})$$

- Combine 1/3 and 3/8 method, so that it can be applied in n points, $n \geq 3$
- If there just 2 points, use the trapezoidal method.