GGS 662 AGENTIC GEOAI · RESEARCH PROGRESS

Seeing Through Clouds

Handling clouds in agentic change detection

TEST CASE: HURRICANE IDA 2021 · SPACENET 8 · 0.5 M EO

Justin Rogers · George Mason University · 30 October 2026

Preliminary results on development data · Imagery: Maxar via SpaceNet 8



Clouds hide what our change detector must see

The system. We are building an agentic change detection system: a vision foundation model finds where two satellite images differ, an agent describes what changed, and code checks every fact.

ONE DIFFICULTY · CLOUDS IN EO IMAGERY

"Clouds can hide target areas in images taken after disasters. This can lead to labels being inaccurate compared to ground truth due to obscured parts of the imagery. How should we detect clouds in this imagery, handle them, and report them? The goal is results, such as maps of flooded roads, that people can trust."

Expected contribution: a tested rule for what a change map may say under cloud, a measured label gap, and a test of whether a second date can correct thin cloud.

REPORT

What can a flood-road map reliably tell users about roads hidden by cloud?

HANDLE

When only the before image is cloudy, does the after image alone find change better than both images together?

DETECT · THE TEST BED

Can we use synthetic clouds to represent real-life blockages in satellite imagery?

Proposed method: detect, handle, report

GGS 680 · CLASSICAL

1 · Detect

Greyscale masks on each date: bright cores (our rule), the dark channel (He et al. 2011), and its rise from before to after (our rule).

GGS 662 · THE AGENT GATE

2 · Handle, feature by feature

ASSESS	clear sky
CORRECT	thin cloud + clear second date
CANNOT ASSESS	opaque cloud

Correct = two-date de-hazing inside the cloud mask (our rule, after He et al. 2011), then the detector again (flood calls: result 3). Filled pixels count as not observed.

WHAT THE MAP MAY SAY

3 · Report

Each statement about a hidden feature gets a tier:

- RELIABLE
- SYNTHETIC ONLY
- RANGE ONLY

CHANGE DETECTOR

Random forest (Breiman 2001) on classical, texture (Haralick 1979) and DINOv2 (Oquab et al. 2024) cues: 83.0% on 798 roads.


BASELINES AND CHECKS

Tile cloud rule; no correction. Tile-blocked folds (Roberts et al. 2017), McNemar tests (McNemar 1947), one sealed check.

AGENTS PROPOSE, I DECIDE

Role agents draft and critique. Decided by hand: the problem statement, the questions, the per-feature cloud rule.

Study area and data




clear thin cloud or haze opaque

South scene (102 tiles) and west scene (115 tiles), after images. Black bar = 2 km.

Imagery	Maxar before and after, 0.5 m, Hurricane Ida 2021; used in greyscale
Roads	1,037 OpenStreetMap roads with SpaceNet 8 flood labels (Hänsch et al. 2022), on 217 of the 226 development tiles
Split	186 train · 40 validation · 40 test tiles, plus a 333-tile north negative set; the test tiles, the negative set, Germany and Louisiana-West stay sealed
Cloud labels	HRC_WHU (Li et al. 2019): 150 Google Earth images, 0.5–15 m, expert cloud masks; 120 development, 30 sealed; academic use only
Quality issues	Flood labels are “yes” or missing, never “no” · 17 tiles under opaque cloud · one after image for all development tiles


Under cloud, roads rarely get a flood label



labelled “flooded” thin cloud

no flood label (reads as dry)


ROADS LABELLED “FLOODED”, BY CLOUD ON THE ROAD




Analysts “only attribute what was visible” (Hänsch et al. 2022), so a flooded road under cloud reads as dry.

Part of the gap is geography: the south scene is dry around its cloudy tiles. Left: tile 2_15_45, after image · Imagery: Maxar via SpaceNet 8


A second image lifts some thin cloud




BEFORE · CLEAR



AFTER · CLOUD



SINGLE IMAGE



TWO-DATE

Road picked by a fixed rule: the most metres under thin cloud (481 m, tile 2_23_44).

Red = labelled flooded · blue = no flood label · orange = opaque, cannot assess.

Haze model as used by He et al. (2011), whose single-image method is the baseline; two-date de-hazing is our rule. Imagery: Maxar via SpaceNet 8.

SYNTHETIC HAZE · 94 CLEAR TILES

	none	He 2011	two-date
grey error	57	27	37
detail kept	0.82	0.84	1.02
clear changed	0%	85%	0%

He 2011 = single-image de-hazing (the baseline).


CAVEAT

The synthetic haze follows the two-date model (favours it); faint haze outside the mask is left alone. Here, little road detail returns.

FLOOD CALLS · EXPLORATORY

Flooded roads under thin cloud found: 20 of 24 two-date, 17 as it is, 21 He 2011. More false alarms too: result 3.

More flooded roads found, and more false alarms



Production flood model, 5 folds by tile, trained on clear roads. Test written down before it ran; the 24 flooded roads had been looked at before. Dry labels under cloud cannot be trusted.

PAIRED · TWO-DATE VS NONE

Flooded found: 4 fixed, 1 broken (McNemar p = 0.38). Clear roads: 5 fixed, 15 broken (p = 0.04).

LIKELY WHY · NOT YET TESTED

The flood model was trained on uncorrected images; de-hazing changes the after image, and the change cues read that as change.

NEXT

Train the flood model on de-hazed images too, then test once on unseen roads.

Feasibility, limits and next steps

Feasible

- Data, masks, de-hazer and the held-out harness all run
- Dark-channel mask: F1 0.65 against 0.41 for the old rule (HRC_WHU)
- Label gap measured: 55%, 26%, 2% labelled flooded (partly geography)

Limits

- One event, one sensor, roads only; all real cloud in one after image
- Sealed test tiles sit next to development tiles
- Only 24 trusted flooded roads under thin cloud
- De-hazing raises false flood calls; its 24 roads were seen before
- The method critic's verdict on the plan: revise

Next steps


- Pre-register; freeze the cloud generator and de-hazer
- Train the flood model on de-hazed images
- Realism checks and the synthetic test bed
- Before-image cloud
- Report tiers and maps
- One look at the sealed data
- Write the paper

QUESTION FOR DISCUSSION

When a map cannot see the ground, should it say “cannot assess”, show a range, or fill it in from another date?

Cloud masks against human cloud labels

F1 AGAINST EXPERT CLOUD MASKS, BY LAND COVER



0.65

F1, dark-channel mask (precision 0.62, recall 0.68)

0.41


F1, bright rule (precision 0.81, recall 0.28)

120 HRC_WHU development images (Li et al. 2019; 30 sealed). Single-date masks only: the road rule's two-date mask needs a before image.

MASK-BIAS CHECK

Synthetic haze on 769 clear roads: under thin haze the mask flags flooded and dry roads alike; under thick haze it flags flooded roads more (up to +13 points), not less.


How the two images are layered



1 · BEFORE

Clear reference: the ground's darkest pixels


+



2 · AFTER

Ground dimmed by t, plus airlight A(1 – t)


→



3 · HAZE t

Inside the cloud mask only; houses stay at 1

→



4 · CORRECTED

Orange: opaque (t below 0.2), cannot assess

```
t = (A − dark after) ÷ (A − tone(dark before)) · corrected = (after − A) ÷ t + A
```

Standard haze model and dark channel as used by He et al. (2011). Our rules: the before image's darkest pixels in place of He's zero assumption, the tone curve, a 2,500 m² cloud minimum (so a changed roof is not haze), a 20 m fade past the mask, and t below 0.2 left as opaque.

References, code and AI use

CITED ON THESE SLIDES · JIF ABOVE 4 OR COMPANY-LED

Li et al. 2019	HRC_WHU cloud masks · ISPRS J. Photogramm. Remote Sens.
He, Sun & Tang 2011	Dark channel prior · IEEE TPAMI
Oquab et al. 2024	DINOv2 (Meta AI) · TMLR
Haralick 1979	Texture features · Proc. IEEE (not yet read)
Roberts et al. 2017	Blocked cross-validation · Ecography

BACKGROUND

Canty & Nielsen 2008 (Remote Sensing of Environment) · Portalés-Julià et al. 2023 (Scientific Reports) · Qu, Li & Huang 2026 (Remote Sensing of Environment) · Shen et al. 2015 (IEEE Geoscience and Remote Sensing Magazine) · Bruzzone & Bovolo 2013 (Proc. IEEE)

METHOD ATTRIBUTIONS · BELOW THE JIF RULE

Breiman 2001, random forests (Machine Learning) · McNemar 1947, paired proportions (Psychometrika)

Data: SpaceNet 8, Hänsch et al. 2022 (licence to confirm) · HRC_WHU (academic use only)

CODE

github.com/Jrog378/GeoAI-Roads (private): cloud masks, both de-hazing methods, the HRC_WHU check, stage notebooks

AI USE

Claude Code (Anthropic) agents drafted questions, critiques, literature notes, code, figures and these slides.

Justin chose the problem, the questions and the cloud rule. Results come from code on development data.

A validation agent checked the claims against the papers; agreement between agents is not validation.