

# Estimation of species richness

Peter Sutovsky

School of Mathematics & Statistics,  
Newcastle University, UK

26th May, 2017

Estimation of  
species  
richness

Peter  
Sutovsky

Introduction

Inference  
Models

Real data  
results

- Biodiversity ("biological diversity," ) – variety and variability of life on Earth.
- Method: metagenomics (environmental genomics, ecogenomics, or community genomics)
  - Study of genetic material recovered directly from environmental samples (Wikipedia)

**PROBLEM: METAGENOMIC SURVEYS RECOVER ONLY A SMALL FRACTION OF THE EXTANT DIVERSITY. NONETHELESS, MANY METHODS TREAT THE OBSERVED SAMPLE AS THE POPULATION.**

Estimation of  
species  
richness

Peter  
Sutovsky

Introduction

Inference  
Models

Real data  
results

- Microbes are required to sustain almost every other form of life and influence
  - Climate, health, and agricultural productivity and the fate of pollutants
  - Sometimes unanticipated modulators
- Laboratory cultures different from real life microbial communities
- How do these microbial communities form?

- Diversity – the number of taxa or species richness as well as their relative abundance.
- Operational taxonomic units (OTUs) – based on differences of 16S rDNA sequences
- Taxa – clusters of sequences that differ by at most 3% of sites
- Microbial world -  $10^{30}$  organisms
  - Vast, diverse and largely unexplored
  - Observed through relatively small sample size
- Sample size - Technological and financial limitations
- How big a sample is big enough?

- Microbial ecology (or environmental microbiology) is the ecology of microorganisms
  - their relationship with one another and with their environment.
- It concerns the three major domains of life – Eukaryota, Archaea, Bacteria, and viruses
  - fingerprinting of microbial communities or assessing biodiversity
- Microbial ecology and biotechnology provide tools to address environmental and economic challenges
  - e.g. for fingerprinting, assessing biodiversity, and tracking the changes of microbial communities

Estimation of  
species  
richness

Peter  
Sutovsky

Introduction

Inference  
Models

Real data  
results

- Relative species abundance and species richness describe key elements of biodiversity
- Relative species abundance
  - component of biodiversity and
  - refers to how common or rare a species is relative to other species in a defined location or community

Estimation of  
species  
richness

Peter  
Sutovsky

Introduction

Inference  
Models

Real data  
results


Estimation of  
species  
richness

Peter  
Sutovsky

Introduction

Inference  
Models

Real data  
results



Cluster sequences at some % “identity,” typically 97%  
 $\{\text{clusters}\} = \{\text{OTUs}\}$  OTU = “operational taxonomic unit”



Estimation of  
species  
richness

Peter  
Sutovsky

Introduction

Inference  
Models

Real data  
results

- Comprised of:
  - *species richness*: number of species present
  - *heterogeneity of species*
    - relative abundance of each species present in the community

- Estimate total population diversity – number of species, classes, taxa, OTUs – based on frequency count data
- Data =
  - # of units observed exactly once in sample (singletons);
  - # observed exactly twice (doubletons);
  - # observed exactly three times; ...

Global Ocean Survey (GOS)  
data from the upper oceans  
given by Rusch et al. (2007)

Abundance	No. of species
1	311
2	213
3	61
4	38
5	33
...	...
34	1
36	2
38	1
39	1
...	...
365	1
1163	1

# Analysis of species abundance data - whole population

11


Estimation of  
species  
richness

Peter  
Sutovsky

Introduction

Inference  
Models

Real data  
results



**Figure:** Abundance of species - whole population

# Analysis of species abundance data - 0.02% of community sampled

12

Estimation of  
species  
richness

Peter  
Sutovsky

Introduction

Inference  
Models

Real data  
results





Figure: Abundance of species - 0.02% of community sampled



**Figure:** TAD of TARA ocean data (ERR315852)

$i$  – abundance

- $P_0 = P(n = 0 | \theta')$
- $f_k$  – number of taxa with abundance  $k$
- $S$  – total number of taxa in the community
- $D = \sum_{k=1}^L f_k$  – number of observed taxa in sample
- $f_0 = S - D$  – number of unobserved taxa in sample

$$\hat{S} = D + \hat{E}[f_0]$$

$$E[f_0] = \sum_{i=1}^S P(n_i = 0)$$

Estimation of  
species  
richness

Peter  
Sutovsky

Introduction

Inference  
Models

Real data  
results

Diversity-estimator methods:

- 1 Parametric
- 2 Non-parametric
- 3 Coverage based



Estimation of  
species  
richness

Peter  
Sutovsky

Introduction

Inference  
Models

Real data  
results

- Based on specific assumptions about the probability distributions of species densities
- Maximize the Likelihood of the observed  $f_k$  as a function of  $S$  and the parameters of the probability distributions of species densities.

- $S$  classes/taxa/species/OTUs in population.
- Assumption: Each species independently contributes Poisson-distributed number of representatives to the sample:
  - Sample:  $X_1, X_2, X_3, X_4, \dots, X_S$
- Counts  $\sim$  zero-truncated mixed Poisson:

$$X_1 \sim \text{Poisson}(\lambda_1), X_2 \sim \text{Poisson}(\lambda_2), \\ \text{Poisson}(\lambda_3), \dots, X_S \sim \text{Poisson}(\lambda_S)$$

- Species (taxon)  $i$  contributes a Poisson-distributed number  $X_i$  of replicates to the sample – i.e., taxon  $i$  appears in the sample  $X_i$
- Units appear independently in the sample.
- Fundamental problem: heterogeneity, i.e., unequal Poisson means  $\lambda_i$
- standard approach: model  $\lambda_i$ 's as i.i.d. replicates from some mixing distribution  $F$
- Frequency counts  $f_k$  are then marginally i.i.d.  $F$ -mixed Poisson random variables Zero-truncated since zero counts  $X_i$  are unobservable

- Mixing distribution  $F$ , i.e., distribution of sampling intensities  $\lambda$ , is also called *species abundance distribution* (**SAD**) or *taxon abundance distribution* (**TAD**)
- Assumptions: Each species contribution to the sample is *independent and identically distributed*
- Both assumptions are probably wrong

# Taxon/species abundance (mixing) distribution 20

Estimation of  
species  
richness

Peter  
Sutovsky

Introduction

Inference  
Models

Real data  
results

- $T(\lambda|\theta)$  – Normalised TAD, where
- $\theta$  is vector of parameters
  - $\lambda$  taxon abundance
- Quince, Curtis, and Sloan (2008)
  - Assumption: probability that individual sampled (with replacement) is from given taxon is  $\frac{\lambda}{N}$ , where  $N$  is size of population
  - Number of times a taxon appears in the sample will be approximately  $\sim \text{Poisson}(\frac{\lambda L}{N})$ , where  $L$  is the sample size and  $\frac{L}{N}$  is sampling frequency

- Parametric, low-dimensional parameter vector
  - 1 Lognormal
  - 2 Inverse Gaussian
  - 3 Generalized inverse Gaussian (Sichel)
  - 4 Log-t
  - 5 None  $\equiv$  point mass at  $\lambda \equiv$  all equal species sizes Gamma (Fisher, 1943)
  - 6 Pareto
  - 7 Stable
- Finite mixture of exponentials - semiparametric

- Approach based on paper of Quince, Curtis, and Sloan (2008)
- Probability that we will observe a taxon  $n$  times:

$$P_n(r, \theta) = \int_0^{\infty} \frac{e^{-r\lambda}}{n!} (r\lambda)^n T(\lambda|\theta) d\lambda, \quad (1)$$

- $T(\lambda|\theta)$  is taxon abundance (mixing) distribution
- $r = \frac{L}{N}$  is sampling ratio
- $N$  is total population number
- $\theta$  is vector of parameters
- $\lambda$  is taxon abundance

- Substitution  $x = r\lambda$

$$P_n(\theta') = \int_0^{\infty} \frac{e^{-x}}{n!} x^n T\left(\frac{x}{r} | \theta\right) dx,$$

using invariance of most abundance distribution to rescaling

$$T\left(\frac{X}{r} | \theta\right) = T(X | \theta')$$

where  $\theta'$  are rescaled parameters

$$P_n(\theta') = \int_0^{\infty} \frac{e^{-x}}{n!} x^n T(X | \theta') dx$$



$$P(f|\theta', S) = P_0^{S-D} \prod_{i=1}^L \frac{P_i^{f_i}}{f_i!} \frac{S!}{(S-D)!}$$

- $i$  – abundance
- $P_0 = P(n = 0|\theta')$
- $f_i$  – number of taxa with abundance  $i$
- $S$  – total number of taxa in the community
- $D = \sum_{i=1}^L f_i$  – number of observed taxa in sample
- $f_0 = S - D$  – number of unobserved taxa in sample

# Estimating parameters of TAD (lognormal distribution)

25

Estimation of  
species  
richness

Peter  
Sutovsky

Introduction

Inference  
Models

Real data  
results

- Estimated parameters: #taxa in community, mean and variance of lognormal distribution
- TAD as lognormal distribution numerical integration to calculate  $P_n$
- Bayesian parameter estimation using **Metropolis-Hastings MCMC** with quasi-noninformative priors
- Run length 400 000 to 1 200 000 steps with 180 000 burn-in period

# Estimating sampling effort with log-normal TAD 26

Estimation of  
species  
richness

Peter  
Sutovsky

Introduction

Inference  
Models

Real data  
results

- From community abundance  $\lambda$  to sample abundance  $x = r\lambda$
- For  $\ln(\lambda = X \frac{N}{L}) \sim N(\mu, \sigma^2)$ ,  
 $\ln(X = \lambda \frac{L}{N}) \sim N(M, V)$ , since  $M = \mu - \ln(\frac{L}{N})$
- Using the fact that  $M^{new} = \ln(\frac{L^{new}}{L}) + M$ .
- **Observed fraction of taxa**  $c^* = \frac{E[D]}{S} = 1 - P_0(M^{new}, V)$
- For chosen  $c^*$ , observed  $L$  and sample of parameters  $M$  and  $V$  from posterior distribution
  - 1 Find  $M^{new}$  s.t.  $c^* = 1 - P_0(M^{new}, v)$
  - 2 Calculate  $L^{new} = \exp(M^{new} - M)$

- Depend on no assumptions about the probability distributions of species densities e.g.
- Chao estimator 1 (Chao 1984):

$$\hat{S}_{min} = D + \hat{E}[f_0], \text{ where}$$

$$\hat{E}[f_0] = \frac{(f_1)^2}{2f_2}$$

and  $f_i$  is number of species with abundance  $i$ .

- First order Jackknife (Burnham and Overton 1979):

$$\hat{E}[f_0] = \frac{L-1}{L} f_1,$$

where  $L$ - number of sampled units

- Coverage is the sum of the proportions of total density accounted for by all species encountered in the sample.
- If all species had equal density
  - $c = \frac{D}{S}$  and therefore  $\hat{S} = \frac{D}{\hat{c}}$
- Chao (Chao 1987, Chao and Lee 1992) has developed coverage-based estimators (**ACE**) by for the general case of unequal densities based on the coverage of infrequent species

# breakaway: Estimating species richness based on ratios of frequency counts

29

Estimation of  
species  
richness

Peter  
Sutovsky

Introduction

Inference  
Models

Real data  
results

- (Willis & Bunge 2015)
- Idea:
- ratios  $r(j) = \frac{(j+1)f_{j+1}}{f_j}$  are  $\sim$  linear :
  - $r(j) = \frac{(j+1)f_{j+1}}{f_j} = \alpha + \beta_j$
- Project line downward to obtain  $f_0 = \#$  of unobserved species

# Estimation of number of species $S$ in community for various microbial datasets

30


Estimation of  
species  
richness

Peter  
Sutovsky

Introduction

Inference  
Models

Real data  
results



# Estimation of number of species in community for selected dataset

31


Estimation of  
species  
richness

Peter  
Sutovsky

Introduction

Inference  
Models

Real data  
results





- Recent results look promising - more testing on real and synthetic data from various environments needed
- Future work
  - Fitting other TAD distributions: inverse Gaussian, log-Student's t, Sichel
  - Model comparison
  - Including into EBI pipeline

This is joint work with Darren Wilkinson and Tom Curtis , funded jointly with the EBI by the **BBSRC** BBR grant: “EBI Metagenomics Portal” led by Rob Finn at the EBI