



Tabla completa – Drogas vasoactivas

	Preparación	Dosis indicación	Estabilidad	Dosis adm
	10mg/100cc	µg/Kg/min	24 horas a temperatura ambiente	0,01-1 µg/Kg/min
	4mg/250cc 8mg/250cc 16mg/250cc 32mg/250cc	µg/Kg/min	24 horas a temperatura ambiente	0,01-2 µg/Kg/min
	500mg/125cc 500mg/250cc	µg/Kg/min	24 horas a temperatura ambiente	0,5-20 µg/Kg/min
	20mg/125cc 20mg/250cc	µg/Kg/min	24 horas a temperatura ambiente	0,125-0,75 µg/Kg/min
	400mg/125cc 400mg/250cc	µg/Kg/min	24 horas a temperatura ambiente	2-50 µg/Kg/min
	600mg/250cc Depende de indicación medica	ml/hr	24 horas a temperatura ambiente	Bolo: 50, 100, 150mg administrar e
n S...	1mg/250cc	µg/min	24 horas a temperatura ambiente, es fotosensible	1-4 µg/min
	50mg/250cc	µg/min	24 horas a temperatura ambiente, es fotosensible	10-200 µg/ min
	50mg/250cc	µg/Kg/min	Seguir recomendación de fabricante	0,05-8 µg/Kg/min
	1mg/1ml	Mg/min	24 horas a temperatura ambiente	0,5-3 mg/min