

Assignment 4: RDD

Juan Martín Londoño

June 14, 2020

1. Create a new github repo named “RDD”.

The link for the GitHub repository is the following: <https://github.com/JuanMartinL/RDD>.

2. Briefly summarize Hansen paper (2015): ”Punishment and Deterrence: Evidence from Drunk Driving”

Benjamin Hansen paper has as principal question if there is a ”(...) effect of harsher punishments and sanctions on driving under the influence (DUI).” (Hansen, 2015, pg. 1581), specially on recidivism over time. To prove his hypothesis, the author uses a Regression Discontinuity methodology on administrative records on 512.964 DUI stops from the state of Washington, with which he suggests that having a blood alcohol content (BAC) above the DUI threshold and aggravated DUI threshold reduces recidivism (Hansen, 2015, pg. 1581). Specifically, 2 percentage points for the first one and one extra percentage point for the second one.

Furthermore, his research design, mainly based in the RDD, is heavily tested: checking balanced controls, heterogeneity of the dependent variable, robustness and three primary channels that could affect the causal effect in the process. Nevertheless, the regression has some problems in fully passing the testes.

3. Create a dummy equaling 1 if bac1_i = 0.08 and 0 otherwise in your do file or R file.

A dummy is created with the following Stata code:

```
gen dbac = 1 if bac1 >= 0.08
replace dbac = 0 if bac1 < 0.08
```

Nevertheless, if more details needed, please go to <https://github.com/JuanMartinL/RDD/Codes/Assignment4.do>.

4. Recreate Figure 1 using the bac1 variable as your measure of blood alcohol content. Do you find evidence for sorting on the running variable?

Sorting in RDD is a situation in which the observed units are selecting to get the treatment or not endogenously. The presence of this situation can cause bias in the estimations. For this reason, it is important to evaluate if the observed units are sorting in any way.

Nevertheless, Hansen recognizes that the decision of drinking more is endogenous. Even so, he defends that the BAC around the threshold is purely random, if seen closely enough.

To prove this assumption, a McCray Test is made, testing if there is any evidence for sorting. In this way, the Figure 1 shows the results of the tests:




Figure 1: McCray Test for BAC

The plot suggest a piece of evidence for rejecting no sorting in the data. Therefore, the test is made to corroborate this graphical analysis, with the following results:

```
rddensity bac1, c(0.08)
```

```
Computing data-driven bandwidth selectors.
```

```
RD Manipulation Test using local polynomial density estimation.
```

Cutoff c = .08	Left of c	Right of c	Number of obs =	214558
-----+-----			Model =	unrestricted
Number of obs	23010	191548	BW method =	comb
Eff. Number of obs	8895	13730	Kernel =	triangular
Order est. (p)	2	2	VCE method =	jackknife
Order bias (q)	3	3		
BW est. (h)	0.011	0.012		

```
Running variable: bac1.
```

-----+-----		
Method	T	P> T
-----+-----		
Robust	2.2032	0.0276
-----+-----		

As seen, there is evidence of sorting around the cutoff taken by the author (0.08). This suggests the observed units are capable of deciding whether get treated or not in the specific cutoff.

Moreover, Placebo Tests in Table 1 shows no evidence of sorting around different cutoffs, excluding the 0.08 cutoff. These Placebo confirms the data is not sorted in any cutoff, but the specific threshold that can cause the treatment or not.

Nevertheless, this may be explained by the use of a different data set than the author used.

5. **Covariate Balance: Recreate Table 2 but only white male, age and accident as dependent variables**

Table 1: Placebo Tests

	0.04	0.05	0.06	0.07	0.08	0.09	1
McCray Test P-Value	0.667	0.922	0.995	0.365	0.027	0.879	0.142

The author checked in his paper some demographic factors such as age, race and gender are stable across the DUI punishment thresholds (Hansen, 2015, pg. 1589). In this sense, he regressed the control variables as dependent variable on the DUI punishment, BAC and its interaction. He expected a stability of predetermined characteristics, as they were controls, to avoid bias.

Given this, the same process is replicated and the results are shown in the Table 2. In this case, the regressions were made with White-Huber Robust Standard Errors to guarantee proper inference.

Table 2: Regression Discontinuity Estimates for the Effect of Exceeding BAC Thresholds on Predetermined Characteristics table

	Driver Demographic Characteristics			
	White (1)	Male (2)	Age (3)	Accident (4)
Panel A. DUI Threshold				
DUI	0.00445 (0.0175)	-0.0184 (0.0198)	-6.224*** (0.586)	-0.154*** (0.0153)
Mean				
Controls	No	No	No	No
Observations	89967	89967	89967	89967

Robust standard errors in parentheses

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

Contrary to Hansen results, some estimators are statistically significant. In his paper, Hansen defends his no-significant estimators of the control variables gives credibility to the regression discontinuity design. Nevertheless, there is evidence of statistical significance at least of two control variables. Therefore, the covariates aren't balanced at the cutoff.

However, a disclaimer must be made: his regressions use a Kernel approach, which is not taken into account here. Although a Kernel with lpoly was considered, no similar results were gotten.

6. Recreate Figure 2 panel A-D. Fit both linear and quadratic with confidence intervals.

Following Hansen's paper, non parametric histograms are calculated for control variables: accident at scene, male, age and white. These are made using linear and quadratic fits. In first place, the linear fit plots are shown in the Figure 2 and the quadratic fit plots are shown in the Figure 3.

In this case, the linear fits are similar to the plots found in Hansen. Nevertheless, Hansen doesn't observe in the paper the quadratic fits, which seems to adjust better for certain variables than the linear one (as seen in age and male). Though this is not considered by Hansen, this observed behaviour of some variables helps the researcher to understand better the analysis that should be done for each control variable.

7. Estimate equation (1) in Hansen's paper with recidivism (recid) as the outcome.




Figure 2: Non parametric fits