

FAT LOSS MASTERY

FAT LOSS AND PRESERVE MUSCLE

THE SCIENCE OF SUSTAINABLE FAT LOSS

TABLE OF CONTENTS

01	WHAT IS BODY FAT AND ITS FUNCTIONS	03	07	THE KETOGENIC DIET (OPTIONAL STRATEGY)	15
02	TYPES OF BODY FAT	05	08	FASTING AND OTHER EFFICIENT OR COMPLEMENTARY STRATEGIES	17
03	BODY FAT PERCENTAGES AND HOW TO DETERMINE THEM	07	09	HOW TO CALCULATE YOUR CALORIC REQUIREMENTS	19
04	OVERVIEW OF FAT-LOSS STRATEGIES	09	10	THE IMPORTANCE OF MEETING PROTEIN REQUIREMENTS	21
05	CREATING A CALORIC DEFICIT BY EATING LESS	11	11	COMMON MISTAKES	23
06	CREATING AN ENERGY DEFICIT THROUGH EXERCISE — HIIT AND CARDIO	13	12	KEY CONSIDERATIONS FOR SAFE AND SUSTAINABLE RESULTS	25

01

WHAT IS BODY FAT AND ITS FUNCTIONS

ENERGY · STRUCTURE · ENDOCRINE ACTIVITY

CHAPTER 01

WHAT IS BODY FAT AND ITS FUNCTIONS

THE BIOLOGICAL TRUTH

Body fat, or adipose tissue, is a specialized connective tissue made of fat cells (adipocytes) that store energy primarily as triglycerides. It is not merely "extra weight"—it is a dynamic organ with multiple essential roles.

ENERGY FUNCTIONS

Fat is the body's most efficient long-term energy store. One gram of fat provides roughly 9 calories, more than double the energy of carbohydrate or protein. When you eat more calories than you burn, excess energy is converted into triglycerides and stored in adipocytes. During energy shortage (fasting, exercise, or a calorie deficit), these triglycerides are broken down into fatty acids and glycerol, which are released into the bloodstream to fuel muscles, organs, and other tissues. This process is tightly regulated by hormones such as insulin, glucagon, catecholamines, and cortisol.

STRUCTURAL AND MATERIAL FUNCTIONS

Fat serves as physical material in several ways:

- Insulation against cold and heat.
- Cushioning and protection for vital organs (heart, kidneys, liver, etc.).
- Structural support around blood vessels, joints, and bones.
- Contribution to cell membranes and hormone production (sex hormones, for example).

ENDOCRINE ACTIVITY

Adipose tissue secretes hormones and signaling molecules (adipokines) such as leptin (satiety) and adiponectin that help regulate hunger, insulin sensitivity, inflammation, and energy balance.

Healthy amounts of body fat are necessary for normal physiology. Excess or insufficient fat can disrupt these functions and increase health risks.

02

TYPES OF BODY FAT

LOCATION · CELL TYPE · DISTRIBUTION

CHAPTER 02

TYPES OF BODY FAT

SUBCUTANEOUS, VISCERAL, AND BEYOND

Body fat is classified by location and by cell type.

BY LOCATION

- **Subcutaneous fat** – Located just under the skin. This is the soft, pinchable fat found on the abdomen, thighs, hips, arms, and face. It makes up the majority (often ~80–90%) of total body fat in healthy people. It provides insulation and padding and is generally less metabolically harmful than deeper fat. Facial fat is primarily a form of subcutaneous fat that influences facial appearance and volume.
- **Visceral fat** – Stored deep in the abdominal cavity around internal organs (liver, pancreas, intestines, kidneys). It is firmer and not easily pinched. Excess visceral fat is strongly linked to higher risk of insulin resistance, type 2 diabetes, cardiovascular disease, and inflammation because the blood from these fat depots drains directly to the liver and carries inflammatory and metabolic signals.
- Other depots include bone-marrow fat and smaller specialized pads (e.g., around the eyes or in the cheeks).

BY CELL TYPE

- White adipose tissue (the main storage type in adults).
- Brown and beige adipose tissue (more metabolically active, involved in heat production/thermogenesis; more abundant in infants and present in smaller amounts in adults).

DISTRIBUTION NOTE

Men tend to accumulate more visceral fat (apple shape); women typically store more subcutaneous fat in the lower body (pear shape), though this can change with age and hormones.

03

BODY FAT PERCENTAGES AND HOW TO DETERMINE THEM

MEASURING WHAT MATTERS

CHAPTER 03

BODY FAT PERCENTAGES AND HOW TO DETERMINE THEM

REFERENCE RANGES AND MEASUREMENT METHODS

Body fat percentage is a more useful measure of health and composition than scale weight or BMI alone, because it distinguishes fat from lean mass (muscle, bone, organs, water).

GENERAL RANGES (AMERICAN COUNCIL ON EXERCISE / SIMILAR GUIDELINES)

CATEGORY	WOMEN	MEN
Essential fat	10–13%	2–5%
Athletes	14–20%	6–13%
Fitness	21–24%	14–17%
Average / Acceptable	25–31%	18–24%
Obese	32%+	25%+

Ranges can shift slightly with age; body fat tends to increase modestly as people get older even at stable weight. Essential fat is the minimum needed for basic physiological function (higher in women due to reproductive needs).

HOW TO MEASURE

- **DEXA scan** – Gold standard for accuracy and regional (including visceral) data.
- **Bioelectrical impedance** (BIA scales or devices) – Convenient for home or gym use; accuracy varies with hydration and conditions (±3–8%). Best for tracking trends under consistent conditions.
- **Skinfold calipers** – Measures subcutaneous fat at specific sites; accuracy depends heavily on the skill of the person taking the measurements.
- **Circumference methods** (e.g., U.S. Navy formula using neck, waist, and hip measurements) – Simple and reasonably accurate for estimates (±3–4%).

- Visual estimation or online calculators based on height/weight are the least precise.

PRACTICAL ADVICE

Track changes over time rather than obsessing over a single number. Progress photos, measurements, and how clothes fit are also useful practical indicators.

04

OVERVIEW OF FAT-LOSS STRATEGIES

THE CORE PRINCIPLE: ENERGY BALANCE

CHAPTER 04

OVERVIEW OF FAT-LOSS STRATEGIES

THE ENERGY BALANCE EQUATION

Sustainable fat loss requires a consistent energy deficit: you must expend more energy than you consume over time.

This can be achieved by reducing calorie intake, increasing energy expenditure through activity, or (most effectively) a combination of both. Additional tools such as specific dietary patterns or meal timing can support adherence and metabolic health, but the fundamental driver remains the deficit.

KEY TAKEAWAY

There is no magic bullet. All effective fat-loss strategies work by creating a sustained energy deficit. The best approach is the one you can maintain consistently without sacrificing health or performance.

The following chapters detail the main approaches, from caloric restriction and exercise to ketogenic diets and intermittent fasting.

Continue to Chapter 5 to learn how to create a practical caloric deficit through nutrition.

05

CREATING A CALORIC DEFICIT BY EATING LESS

HOW TO DO IT – STEP BY STEP

CHAPTER 05

CREATING A CALORIC DEFICIT BY EATING LESS

PRACTICAL NUTRITION STRATEGIES

A caloric deficit is created when daily energy intake is lower than total daily energy expenditure (TDEE).

A moderate, sustainable deficit is typically 300–600 calories per day below maintenance (roughly 15–25% below TDEE for most people). This usually produces fat loss of about 0.5–1% of body weight per week (roughly 0.5–1 kg / 1–2 lb per week for many individuals). Larger deficits accelerate loss but increase the risk of muscle loss, metabolic adaptation, nutrient deficiencies, fatigue, and poor adherence.

STEP 1: CALCULATE YOUR TDEE AND SET YOUR DEFICIT

Use the Mifflin-St Jeor equation to estimate your BMR, then multiply by an activity factor. Let's do an example:

EXAMPLE CALCULATION

Person: 30-year-old male, 80 kg (176 lb), 180 cm (5'11"), moderately active (exercises 3–5 days/week).

BMR = $(10 \times 80) + (6.25 \times 180) - (5 \times 30) + 5 = 800 + 1125 - 150 + 5 = \mathbf{1780 \text{ kcal/day}}$.

TDEE = 1780×1.55 (moderately active) = **2759 kcal/day**.

Deficit: Subtract 500 kcal → **2259 kcal/day** (about 18% below maintenance). This will yield ~0.5 kg (1 lb) of fat loss per week.

STEP 2: PLAN YOUR MEALS – EXAMPLE DAY

Here's a sample day at ~2200 kcal with high protein and fiber to keep you full:

- **Breakfast (07:30):** 3 whole eggs + 2 egg whites scrambled with spinach, 1 slice whole-grain toast, 1 tbsp olive oil. *~450 kcal*
- **Snack (10:30):** Greek yogurt (200 g) with 50 g berries and 15 g almonds. *~300 kcal*
- **Lunch (13:30):** Grilled chicken breast (200 g) with quinoa (150 g cooked) and roasted broccoli (200 g). *~600 kcal*
- **Snack (16:30):** Apple with 1 tbsp peanut butter. *~200 kcal*
- **Dinner (20:00):** Salmon fillet (150 g) with sweet potato (200 g) and asparagus. *~650 kcal*

Total: ~2200 kcal, ~160 g protein, ~200 g carbs, ~70 g fat.

STEP 3: TRACK AND ADJUST

- Use a food scale and an app (MyFitnessPal, Cronometer) for at least 2 weeks to learn true portion sizes.
- Weigh yourself daily (same time, after bathroom) and take weekly averages. If weight loss is too fast (>1% per week) or you feel lethargic, add 100–200 kcal. If no loss after 2 weeks, reduce by 100–200 kcal.
- Re-calculate TDEE after every 5 kg lost, as your BMR drops with weight.

CAUTION

Avoid extreme restriction. Very low-calorie approaches (<1200 kcal for women, <1500 for men) should only be used short-term under professional supervision.

06

CREATING AN ENERGY DEFICIT THROUGH EXERCISE

HIIT AND CARDIO – HOW TO IMPLEMENT

CHAPTER 06

CREATING AN ENERGY DEFICIT THROUGH EXERCISE — HIIT AND CARDIO

SAMPLE SESSIONS AND WEEKLY PLAN

Increasing energy expenditure through exercise helps enlarge the deficit and supports muscle preservation and metabolic health when combined with strength training and adequate protein.

EXAMPLE HIIT SESSION (20 MINUTES TOTAL)

HIIT PROTOCOL

- **Warm-up (5 min):** Light jog or dynamic stretches.
- **Intervals (4 rounds):** 30 seconds all-out sprint (or burpees, kettlebell swings, rower) followed by 90 seconds active recovery (walking or slow pedaling). Repeat 4 times.
- **Cool-down (3 min):** Walk and static stretches.

This session burns ~250–300 kcal and elevates metabolism for several hours post-workout (EPOC).

EXAMPLE STEADY-STATE CARDIO SESSION

45 minutes of brisk walking, jogging, cycling, or swimming at 60–75% of max heart rate (you can talk but not sing). Burns ~300–400 kcal depending on body weight and intensity.

WEEKLY EXERCISE PLAN (COMBINING STRENGTH, HIIT, AND CARDIO)

SAMPLE WEEK

DAY	ACTIVITY	DURATION
Monday	Strength training (full body) + 20 min LISS (walk)	60 min
Tuesday	HIIT session + core work	30 min
Wednesday	Strength training (upper body) + 30 min jog	70 min
Thursday	Active recovery (light yoga or long walk)	40 min
Friday	Strength training (lower body) + 20 min LISS	60 min
Saturday	HIIT session	25 min
Sunday	Rest or gentle stretching	—

Adjust frequency based on recovery. Beginners can start with 2 strength + 1 HIIT + 2 LISS days.

KEY PRINCIPLE

Consistency beats intensity. Choose activities you enjoy and can sustain for weeks. Walking is highly underrated – 10,000 steps/day can add 300–400 kcal expenditure.

07

THE KETOGENIC DIET

OPTIONAL STRATEGY – HOW TO DO IT

CHAPTER 07

THE KETOGENIC DIET (OPTIONAL STRATEGY)

VERY LOW-CARB, HIGH-FAT – IMPLEMENTATION GUIDE

A ketogenic diet is a very low-carbohydrate, high-fat, moderate-protein approach (typically ≤ 20 –50 g carbs per day) that induces nutritional ketosis, in which the body uses ketones from fat as a primary fuel source.

STEP 1: CALCULATE YOUR KETO MACROS

EXAMPLE FOR A 75 KG PERSON

- **Calories:** 2000 kcal/day (deficit).
- **Protein:** 1.6 g/kg = 120 g (480 kcal).
- **Carbs:** 25 g (100 kcal).
- **Fat:** Remaining calories = $2000 - 480 - 100 = 1420$ kcal $\rightarrow 1420 / 9 =$ **158 g fat**.

Macro ratio: ~70% fat, 25% protein, 5% carbs.

STEP 2: SAMPLE KETO MENU (ONE DAY)

EXAMPLE DAY

- **Breakfast:** 3-egg omelet with 50 g cheese, cooked in 1 tbsp butter, plus 1 cup spinach.
- **Lunch:** Salad with 150 g grilled chicken, 2 tbsp olive oil, 50 g avocado, and 30 g walnuts.
- **Snack:** 30 g macadamia nuts and 1 celery stick with cream cheese.
- **Dinner:** 200 g salmon with 2 cups broccoli sautéed in coconut oil.

Total: ~2000 kcal, 20 g carbs, 120 g protein, 158 g fat.

STEP 3: TRANSITION AND MANAGE SIDE EFFECTS

- **Start gradual:** Reduce carbs by 30–50 g per week to minimize "keto flu" (headache, fatigue).
- **Hydrate:** Drink 3–4 L water and increase salt intake (sodium, potassium, magnesium) to offset electrolyte loss.
- **Test ketosis:** Use urine strips or blood ketone meter (optional).
- **Adjust protein:** Too much protein can kick you out of ketosis; stick to 1.6–2.0 g/kg lean mass.

IMPORTANT

Keto is optional. If you feel unwell or cannot sustain it, switch to a moderate low-carb or balanced diet. The key remains the calorie deficit, not the carb count.

08

FASTING AND OTHER EFFICIENT OR COMPLEMENTARY STRATEGIES

INTERMITTENT FASTING · NEAT · SLEEP

CHAPTER 08

FASTING AND OTHER EFFICIENT OR COMPLEMENTARY STRATEGIES

INTERMITTENT FASTING · NEAT · SLEEP

INTERMITTENT FASTING (IF)

IF involves cycling between periods of eating and fasting. Common protocols:

- **Time-restricted eating** (e.g., 16:8 — eat within an 8-hour window, fast 16 hours).
- **5:2** (normal eating five days, very low calories two non-consecutive days).
- **Alternate-day** approaches.

Evidence shows IF produces fat loss similar to continuous daily calorie restriction, primarily because it often leads to a spontaneous reduction in overall calorie intake. It can improve insulin sensitivity and may help some people manage hunger or simplify meal planning. Early time-restricted eating (eating earlier in the day) appears particularly beneficial for metabolic markers in some studies.

GETTING STARTED WITH IF

Start gradually (e.g., 12:12 or 14:10) and ensure protein and overall nutrition targets are still met within the eating window. Not everyone responds well; women or those with high training loads sometimes need more flexible approaches.

OTHER COMPLEMENTARY STRATEGIES

- Prioritize sleep and stress management (poor sleep and high cortisol impair fat loss and increase appetite).
- Increase daily non-exercise activity (NEAT) — walking, standing, light movement.
- Strength training (as already noted).
- Adequate fiber and protein for satiety.

- Mindful eating and reducing ultra-processed foods.

IMPORTANT REMINDER

None of these replace the need for an overall energy deficit.

09

HOW TO CALCULATE YOUR CALORIC REQUIREMENTS

THE NUMBERS BEHIND FAT LOSS

CHAPTER 09

HOW TO CALCULATE YOUR CALORIC REQUIREMENTS

FROM BMR TO TDEE – A STEP-BY-STEP GUIDE

Knowing your numbers is the foundation of any fat-loss plan. Without an accurate estimate of your energy needs, you are guessing – and guessing usually leads to frustration.

Your **Basal Metabolic Rate (BMR)** is the number of calories your body burns at complete rest, just to keep your heart beating, lungs breathing, and brain functioning. Your **Total Daily Energy Expenditure (TDEE)** is your BMR multiplied by your activity level.

STEP 1: ESTIMATE YOUR BMR (MIFFLIN-ST JEOR EQUATION)

EQUATIONS

Men: $\text{BMR} = (10 \times \text{weight in kg}) + (6.25 \times \text{height in cm}) - (5 \times \text{age}) + 5$

Women: $\text{BMR} = (10 \times \text{weight in kg}) + (6.25 \times \text{height in cm}) - (5 \times \text{age}) - 161$

STEP 2: MULTIPLY BY AN ACTIVITY FACTOR TO GET TDEE

ACTIVITY MULTIPLIERS

- **Sedentary** (little/no exercise): $\times 1.2$
- **Lightly active** (1–3 days/week): $\times 1.375$
- **Moderately active** (3–5 days/week): $\times 1.55$
- **Very active** (6–7 days/week hard exercise): $\times 1.725$
- **Extra active** (physical job + intense training): $\times 1.9$

Be honest here – most people overestimate their activity level. If in doubt, choose the lower multiplier.

STEP 3: APPLY A MODERATE DEFICIT

Subtract **300–600 kcal** (or 15–25% of TDEE) to create your daily target. This will yield ~0.5–1% body weight loss per week.

WORKED EXAMPLE

Person: 35-year-old woman, 72 kg, 165 cm, moderately active (3–5 days/week).

BMR = $(10 \times 72) + (6.25 \times 165) - (5 \times 35) - 161 = 720 + 1031 - 175 - 161 = \mathbf{1415 \text{ kcal}}$.

TDEE = $1415 \times 1.55 = \mathbf{2193 \text{ kcal}}$.

Deficit target = $2193 - 400 = \mathbf{\sim 1800 \text{ kcal/day}}$.

After 4 weeks, if weight loss stalls, reduce by another 100 kcal or increase daily steps by 2000.

WHEN TO ADJUST

- **Every 5 kg (11 lb) lost:** Recalculate BMR – your body now needs fewer calories.
- **If you lose >1% body weight per week:** Increase calories slightly (muscle loss risk).
- **If you lose <0.5% per week after 3 weeks:** Reduce calories by 100–150 or add a 20-min cardio session.

DIET BREAKS

After 8–12 weeks of continuous dieting, consider 1–2 weeks at maintenance calories (TDEE). This helps restore leptin levels, reduce metabolic adaptation, and improve adherence for the next fat-loss phase.

10

THE IMPORTANCE OF MEETING PROTEIN REQUIREMENTS

MUSCLE PRESERVATION DURING A DEFICIT

CHAPTER 10

THE IMPORTANCE OF MEETING PROTEIN REQUIREMENTS

LEUCINE, SATIETY, AND LEAN MASS

In a calorie deficit, your body is in a catabolic state – it will break down muscle tissue for energy if protein intake is insufficient. High protein intake is the single most important nutritional lever to **preserve muscle, increase satiety, and boost the thermic effect of food** (you burn ~20–30% of protein calories during digestion).

HOW MUCH PROTEIN DO YOU NEED?

Evidence-based targets during fat loss: **1.6–2.2 g per kg of body weight** (or ~0.7–1 g per pound). For leaner individuals or those doing intense resistance training, aim for the higher end.

EXAMPLE

A person weighing 80 kg (176 lb) should aim for **128–176 g of protein per day**.

Distribute this across 3–5 meals, with **~30–40 g per meal** to maximise muscle protein synthesis (MPS).

THE LEUCINE THRESHOLD

Leucine is the key amino acid that triggers MPS. You need **~2.5–3 g of leucine** per meal to maximise the anabolic response. This is found in roughly 30 g of high-quality protein (whey, chicken, beef, eggs, fish).

BEST PROTEIN SOURCES (PROTEIN PER 100 G / PROTEIN-TO-CALORIE RATIO)

- **Chicken breast** – 31 g protein, 165 kcal (high ratio)
- **Turkey breast** – 30 g, 135 kcal (very high)
- **Lean beef (95/5)** – 29 g, 170 kcal
- **Salmon** – 25 g, 208 kcal (high in omega-3)
- **Egg whites** – 11 g, 52 kcal (per 100 g)
- **Greek yogurt (0%)** – 10 g, 59 kcal (per 100 g)
- **Tofu** – 17 g, 145 kcal
- **Whey protein** – ~80–90 g per 100 g powder

SAMPLE DISTRIBUTION ACROSS MEALS

DAILY PLAN (150 G PROTEIN)

- **Breakfast:** 3 eggs + 3 egg whites = ~30 g
- **Lunch:** 200 g chicken breast = ~60 g
- **Snack:** 200 g Greek yogurt = ~20 g
- **Dinner:** 150 g salmon = ~40 g

Total: 150 g protein. Adjust portions to your target.

DON'T FORGET

Protein sources also contain calories from fat and carbs. Account for them in your total daily target. Prioritise whole foods over supplements, but use whey or plant-based powders to hit your numbers if needed.

11

COMMON MISTAKES

WHAT TO AVOID – AND HOW TO FIX IT

CHAPTER 11

COMMON MISTAKES

PITFALLS THAT DERAIL PROGRESS – AND SOLUTIONS

Even with the best intentions, small errors can compound and stall your results. Here are the most frequent mistakes, why they happen, and exactly how to correct them.

1. UNDERESTIMATING CALORIE INTAKE

Why it happens: Portion distortion, forgetting cooking oils, condiments, and "small bites" (tasting while cooking). Studies show people under-report intake by 30–50%.

FIX

Use a **food scale** for 2–4 weeks. Weigh everything that goes into your mouth. Log it in an app (MyFitnessPal, Cronometer). You will be shocked at the difference between "eyeballing" and actual weight.

2. STARTING TOO AGGRESSIVELY

Why it happens: Impatience and the desire for rapid results. Severe deficits (>30% of TDEE) trigger a cascade of hormonal adaptations – leptin drops, cortisol rises, hunger skyrockets, and NEAT (non-exercise activity) plummets.

FIX

Start with a **moderate deficit (300–500 kcal)**. Focus on consistency over perfection. If you stall, add activity rather than slashing more calories. Think of fat loss as a marathon, not a sprint.

3. IGNORING RESISTANCE TRAINING AND PROTEIN

Why it happens: Many people focus solely on cardio and cutting calories. Without the stimulus of resistance training and adequate protein, up to 25–30% of weight lost can be muscle, lowering your metabolism.

FIX

Commit to **2–4 strength sessions per week**. Hit your protein target daily (1.6–2.2 g/kg). Treat muscle as your metabolic shield – the more you keep, the easier it is to maintain your results.

4. RELYING SOLELY ON THE SCALE

Why it happens: The scale is easy and immediate. But weight fluctuates daily due to water, glycogen, food mass, and hormonal cycles.

FIX

Use **multiple metrics**: weekly average weight, tape measurements (waist, hips), progress photos every 4 weeks, and how your clothes fit. A stagnant scale but shrinking waist means you are losing fat and gaining muscle – which is a win.

5. NEGLECTING SLEEP AND STRESS

Why it happens: These are often overlooked in favour of diet and exercise. Poor sleep raises ghrelin (hunger) and lowers leptin (satiety). High cortisol encourages visceral fat storage.

FIX

Aim for **7–9 hours of quality sleep**. Manage stress with walks, meditation, or breathwork. These are non-negotiable for optimal hormonal function.

KEY MINDSET SHIFT

Consistency over perfection. One bad meal or missed workout does not ruin your progress – it's the following three days of "I already messed up" that do. Get back on track immediately.

12

KEY CONSIDERATIONS FOR SAFE AND SUSTAINABLE RESULTS

LONG-TERM SUCCESS

CHAPTER 12

KEY CONSIDERATIONS FOR SAFE AND SUSTAINABLE RESULTS

BEYOND THE DEFICIT – THE BIG PICTURE

You have the tools – calorie targets, protein goals, training plans, and strategies. But fat loss is not just about the numbers. It is about **long-term adherence, metabolic health, and psychological well-being**.

METABOLIC ADAPTATION – WHY IT HAPPENS

As you lose weight, your BMR drops – not just because you weigh less, but because your body becomes **more efficient**. Hormones like leptin decrease, thyroid output can reduce, and NEAT often declines unconsciously. This is why progress slows over time.

WHAT TO DO

- **Recalculate TDEE every 5 kg lost.**
- **Incorporate refeed days** (1–2 days at maintenance) every 4–6 weeks to restore leptin and glycogen.
- **Increase activity** (steps or light cardio) rather than continually lowering calories.

MANAGING PLATEAUS

A true plateau is **3+ weeks with no change** in weight or measurements, despite perfect adherence. Solutions:

- Re-check your tracking accuracy (are you weighing everything?).
- Increase daily steps by 2,000–3,000.
- Add 1 extra cardio session or increase intensity.
- Take a 1-week diet break at maintenance – this often "resets" the system and allows for a smoother next phase.

THE IMPORTANCE OF MAINTENANCE PHASES

Many people lose weight and then return to old habits, rebounding. Plan a **maintenance period of 4–8 weeks** after reaching your goal. During this time:

- Eat at your new TDEE (calculated at your new weight).
- Continue training and tracking to solidify habits.
- This allows your hormones and hunger signals to recalibrate, drastically reducing the risk of weight regain.

MEDICAL CONSIDERATIONS

If you have underlying conditions (thyroid, diabetes, PCOS), consult a healthcare professional before starting. Very low body fat levels (<10% men, <18% women) can disrupt hormones – know your limits.

FINAL REFLECTION

Fat loss is a journey of self-discovery. It is about building **habits that last a lifetime**, not just reaching a number on the scale. Celebrate non-scale victories: better energy, improved mood, stronger performance, and greater confidence. You are forging a stronger version of yourself – one consistent day at a time.

— End of Guide —