ABBA2 Midpoint

Explicit duplicated-space step with arithmetic projection

Explicit palindromic operations

All ABBA and BM4 presets inherit `PalindromicMethod`. For signed $d = ch$, the elementary operations are

$$A_d(u, v, t) = (u + df(t, v), v, t), \qquad B_d(u, v, t) = (u, v + df(t, u), t), \qquad T_d(u, v, t) = (u, v, t + d).$$

Only T advances time. A coefficient $c = 1/2$ means duration $h/2$; there is no additional half inside A or B. The optional exact C flow rotates the copy difference and requires an independent coupling object. Both BM4 presets omit C.

In chronological order, standard ABBA2 is $A_{h/2}, B_{h/2}, T_{h/2}, T_{h/2}, B_{h/2}, A_{h/2}$. The distinct symmetric variant $T_{h/2}, A_{h/2}, B_{h/2}, B_{h/2}, A_{h/2}, T_{h/2}$ evaluates every spatial shear at the midpoint time. Both exhibit second order on the tested nonautonomous rotation. Arbitrary palindromes do not inherit fourth or sixth order, and an arithmetic projection need not be reversible or symplectic.

`Composition.from_half` reflects a supplied half sequence. The generic `PalindromicImplicit` and `PalindromicMidpoint` methods select one exterior Hairer projection or a final arithmetic mean. The standard signed ABBA/BM4 recipes, grouped observations and passive momentum normalization remain unchanged. NumPy and JAX execute the same explicit operations and differentiate their retained shear inputs. See `docs/models/extended/simulation/palindromic-operations.md`.

1 Definition

For $\dot{z} = f(t, z)$, $z \in \mathbb{R}^m$, duplicate the accepted state on the diagonal, $u_0 = v_0 = z_n$, and write $s = h/2$. The endpoint-time ABBA stages are

$$u_1 = u_0 + sf(t_n, v_0), \qquad v_1 = v_0 + sf(t_n, u_1), \tag{1}$$

$$v_f = v_1 + sf(t_n + h, u_1), \qquad u_f = u_1 + sf(t_n + h, v_f). \tag{2}$$

The physical output is the midpoint of the copies,

$$z_{n+1} = \frac{u_f + v_f}{2}.$$

"Midpoint" therefore refers to this arithmetic projection; the method is not the implicit-midpoint Runge–Kutta rule.

2 State and energy configurations

One endpoint-time A-B-B-A map followed by a spatial arithmetic mean. Energy tracking is optional and passive. The four accepted shear stages update the normalized passive momentum; no nonlinear solve is used. The common state conventions are specified below.

3 Symmetry, order, and geometry

The four shears form an explicit self-adjoint map on the duplicated phase space when the signed step reverses together with its endpoint times. Starting from the diagonal and taking the arithmetic

mean yields a consistent symmetric method with designed global order two for sufficiently smooth dynamics.

The duplicated ABBA map is symplectic for its split Hamiltonian. Arithmetic projection, however, is not Hairer's symmetric projection and does not in general make the closed physical map symplectic. The copy-separation quantity

$$d_n = \|u_f - v_f\|_\infty$$

is therefore retained as a diagnostic. It is neither an error estimator nor a convergence residual.

4 Work and time dependence

One step performs four vector-field evaluations. The evaluations at t_n and $t_n + h$ are part of the non-autonomous definition. The common fixed-grid runner chooses a uniform main grid independently of requested output times; off-grid output uses shadow steps that do not alter the accepted main trajectory or its diagnostics.

5 Implementation correspondence

The public class is in `src/methods/extended/abba.py`. It uses the common `midpoint_step`, the uncoupled ABBA pair in `composition.py`, and the same passive energy quadrature as implicit ABBA and BM4.

The public names, constructor defaults and exports from `simulation` remain unchanged. Internal imports use the defining modules; old import adapters have been removed. The ordinary implicit path is `advance`, `solve_projection`, `compose`, then explicit $A/B/T/C$ operations. Arithmetic midpoint variants use one explicit average instead of a nonlinear projection. Adjoint stages evaluate at the start of their signed substep; direct stages evaluate at its end.

Newton and Broyden share the spatial equations and convergence services. Exact particle-block tangents, passive energy and optional observation use retained shear inputs; energy is evaluated only after accepting the spatial map, with the existing factor $1/2$ for physical κ. Time and momentum never enter the spatial root. The new internal family name does not enable the retired full time/momentum projection.

Current module responsibilities and diagrams are documented in `docs/models/extended/simulation/extended-simulation-architecture.md`.

6 Four state formulations and passive energy tracking

For one planar particle the supported internal representations are

physical	$z = (x, y)$	$\mathbb{R}^2$,
physical with energy	(z, t, κ)	$\mathbb{R}^4$,
duplicated	$(u, v), \quad u, v \in \mathbb{R}^2$	$\mathbb{R}^4$,
duplicated with energy	(u, v, t, κ)	$\mathbb{R}^6$.

Classical methods use the physical rows; ABBA and BM4 use the duplicated rows. `track_energy` selects the energy variant. For N planar particles the respective dimensions are $2N$, $4N$, $4N$, and $6N$: one time entry and one momentum per particle. All time entries equal the integration time. Classical full-cyclotron runs instead use $4N$ or $6N$ entries. Accepted spatial copies coincide; only one physical copy is exported. Every coordinate block contains N values in component-major order. The formulation validates particle times and aligns output round-off; the integrator does not interpret coordinate indices.

The normalized passive balance is

$$\dot{\kappa} = -\partial_t H(t, z), \qquad \kappa(t_0) = 0, \qquad \Delta K(t) = H(t, z(t)) + \kappa(t) - H(t_0, z_0).$$

Splitting accumulators from both copies are divided by two. The physical Hamiltonian may vary with time; ΔK measures its computed balance. Tracking leaves physical trajectories, nonlinear stopping rules and adaptive acceptance unchanged. Dynamics must supply the momentum derivative, including zero for an autonomous Hamiltonian.

Hairer's constraint is solely $u - v = 0$: the reduced multiplier has $2N$ entries and ABBA's simultaneous spatial solve has $6N$ unknowns. Neither t nor κ is projected. The former eight-component projection is retired; `state_extension='fully_extended'` raises migration guidance.

Dynamics, formulation, method and integration remain separate. The shared state classes own layouts, diagonal embedding and history extraction in

`src/formulations/state.py`

Each integration initializes a fresh method instance. Observers receive physical states; energy histories are in `Solution.diagnostics`. The `extended_time` history has shape (N, n_{samples}). Fixed-step shadow samples leave main states and counters unchanged. Adaptive energy quadrature uses accepted dense output; its backend state and Jacobian remain physical.

7 Optional JAX execution

An optional JAX driver executes the same numerical map on one selected CPU or GPU, with particle operations batched inside a compiled sequential time loop. It reuses the canonical coefficients and prepared potential splines. Physical-state tolerance scaling, signed stage times, passive energy weights and independent shortened output maps retain their definitions above. Applicable Newton systems use independent particle blocks; an explicitly selected Broyden iteration retains its global secant matrix. Nonlinear iteration limits and failure checks remain active on the device.

Both execution routes use double precision and return the same physical solution layout. Floating-point reassociation can produce round-off differences without changing the designed order or geometric properties. First-call compilation is separate from warmed integration timings, which include transfers and output construction. The sibling runtime document and shared execution contract specify supported callbacks and exact device scope.