ABBA and BM4 in Extended Space

Common composition, spatial projection and passive energy

Explicit palindromic operations

All ABBA and BM4 presets inherit `PalindromicMethod`. For signed $d = ch$, the elementary operations are

$$A_d(u, v, t) = (u + df(t, v), v, t), \qquad B_d(u, v, t) = (u, v + df(t, u), t), \qquad T_d(u, v, t) = (u, v, t + d).$$

Only T advances time. A coefficient $c = 1/2$ means duration $h/2$; there is no additional half inside A or B. The optional exact C flow rotates the copy difference and requires an independent coupling object. Both BM4 presets omit C.

In chronological order, standard ABBA2 is $A_{h/2}, B_{h/2}, T_{h/2}, T_{h/2}, B_{h/2}, A_{h/2}$. The distinct symmetric variant $T_{h/2}, A_{h/2}, B_{h/2}, B_{h/2}, A_{h/2}, T_{h/2}$ evaluates every spatial shear at the midpoint time. Both exhibit second order on the tested nonautonomous rotation. Arbitrary palindromes do not inherit fourth or sixth order, and an arithmetic projection need not be reversible or symplectic.

`Composition.from_half` reflects a supplied half sequence. The generic `PalindromicImplicit` and `PalindromicMidpoint` methods select one exterior Hairer projection or a final arithmetic mean. The standard signed ABBA/BM4 recipes, grouped observations and passive momentum normalization remain unchanged. NumPy and JAX execute the same explicit operations and differentiate their retained shear inputs. See `docs/models/extended/simulation/palindromic-operations.md`.

1 Scope and public methods

The directory `src/methods/extended/` contains the common implementation of ABBA, BM4 and generic palindromic methods. It introduces no public method named Extended. Public class names and exports from `simulation` are retained. ABBA6 now uses the same single outer projection as ABBA4. Its observed substeps become unprojected pairs and its work arrays have one column per step. Internal imports use the defining modules. Old ABBA and BM4 import adapters and historical `simulation.methods` routes have been removed.

For N independent planar particles the accepted physical state lies in $\mathbb{R}^{2N}$. Scalar time and optional N physical momenta are separate. Two temporary copies in $\mathbb{R}^{4N}$ are used only inside the composition and projection; they are discarded before acceptance.

2 Direct and adjoint compositions

Let χ_s denote a first-order map on the extended spatial workspace, with adjoint $\chi_s^* = \chi_{-s}^{-1}$, including the appropriate time translation for nonautonomous dynamics. A recipe with $2r$ stages is

$$\Psi_h = \chi_{b_{2r}h} \circ \chi_{b_{2r-1}h}^* \circ \cdots \circ \chi_{b_2 h} \circ \chi_{b_1 h}^*.$$

In execution order the stages alternate adjoint, direct. The weights obey $b_j = b_{2r+1-j}$ and $\sum_j b_j = 1$. Reflection exchanges each map with its adjoint. The direct/adjoint builder checks these weights and expands them into elementary operations. The resulting recipe checks reflected operation pairs and unit A, B and T totals; higher-order conditions remain necessary.

With stage clock τ and signed duration $s = b_j h$, the adjoint evaluates at τ, the direct map at $\tau + s$, and the clock subsequently becomes $\tau + s$. Negative coefficients reverse the clock as well as the subflow. For uncoupled GC maps, the pair with weights $(1/2, 1/2)$ reproduces the endpoint-time

A–B–B–A construction. Each direct or adjoint map contains two spatial shears. Both BM4 variants also omit harmonic coupling. The exact C flow remains independently available through `HarmonicCoupling`, outside BM4.

ABBA4 uses three unprojected ABBA pairs scaled by

$$\gamma = \frac{1}{2 - 2^{1/3}}, \qquad \delta = -\frac{2^{1/3}}{2 - 2^{1/3}}, \qquad (\gamma, \delta, \gamma).$$

Its six direct/adjoint weights are $(\gamma/2, \gamma/2, \delta/2, \delta/2, \gamma/2, \gamma/2)$. BM4 retains its twelve mirrored weights and its own fourth-order recipe. ABBA6 uses seven *unprojected* ABBA2 pairs with the Yoshida weights. Each weight is repeated twice after division by two, producing fourteen direct/adjoint stages. One symmetric projection encloses the complete composition. This deliberately replaces the former seven-projection map.

3 One common spatial projection equation

Write $m = 2N$, $Ez = (z, z)$, $G = [I, -I]$ and $N_0 = G^{\mathsf{T}}$. The reduced Hairer projection finds $\mu \in \mathbb{R}^m$ such that

$$r(\mu) = G\Psi_h(Ez + N_0\mu) + 2\mu = 0.$$

After convergence the two corrected outputs are averaged. With $P = \frac{1}{2}[I, I]$, the accepted state is

$$z_+ = P\big(\Psi_h(Ez + N_0\mu) + N_0\mu\big).$$

The same equation applies to ABBA2, the complete ABBA4 and ABBA6 compositions, and BM4; only Ψ_h changes. All have one nonlinear solve per complete step.

The simultaneous ABBA formulation solves for $(u_+, v_+, \mu) \in \mathbb{R}^{3m}$:

$$\begin{pmatrix} u_+ - \mu - [\Psi_h(Ez + N_0\mu)]_u \\ v_+ + \mu - [\Psi_h(Ez + N_0\mu)]_v \\ u_+ - v_+ \end{pmatrix} = 0.$$

Arithmetic midpoint variants instead start from Ez, apply the complete recipe, and average once. They do not solve a nonlinear equation; arithmetic projection does not imply exact symplecticity or reversibility.

4 Nonlinear solution and exact tangents

All spatial roots use the unchanged stopping threshold

$$\mathrm{atol} + \mathrm{rtol}\max(1, \|z\|_\infty).$$

The equation classes `ReducedProjection` and `SimultaneousProjection` in `src/methods/projection.py` construct these residuals and their Jacobians. The numerical method supplies its composition; the algebraic solver receives the resulting system. Newton and good Broyden share the convergence machinery in `src/solvers/nonlinear.py`. Broyden's initial reduced Jacobian is $4I$, the exact zero-step value of Dr. The simultaneous zero-step block matrix is

$$\begin{pmatrix} I_{2m} & -2N_0 \\ G & 0 \end{pmatrix}.$$

Broyden subsequently uses residual secants and evaluates no analytic Jacobian.

If $M = D\Psi_h$, Newton's reduced Jacobian is $GMN_0 + 2I$. The common composition retains the input to each spatial shear. Exact stage tangents are evaluated at those inputs and multiplied in execution order, $M = M_{2r} \cdots M_1$, without replaying vector-field stages. Independence of the particles gives 4×4 blocks for M, 2×2 reduced Newton blocks, and 6×6 simultaneous blocks. BM4 also retains its optional dense centered-difference differentiation of the complete map.

5 Passive energy from accepted stages

Execution supplies $H(t, z)$ and $-\partial_t H(t, z)$ to an energy observer. The method supplies only numerical quadrature data. For the accepted spatial trace, let (t_j, s_j, z_j) denote the evaluation time, signed duration and physical source state of each individual shear. The shared quadrature is

$$\Delta\kappa = \frac{1}{2}\sum_j s_j \left[-\partial_t H(t_j, z_j)\right].$$

There are four shear points for ABBA2, twelve for ABBA4, 28 for ABBA6, and 24 for BM4. The factor $1/2$ converts summed doubled momentum to the physical variable κ. Spatial harmonic coupling does not change this passive accumulator. Energy derivatives are evaluated only after the spatial root converges, from its retained trace; trial roots do not evaluate energy. The accepted diagnostic is

$$H(t_n, z_n) + \kappa_n - H(t_0, z_0), \qquad \kappa_0 = 0.$$

This is an energy-balance diagnostic, not a claim that the time-dependent physical Hamiltonian is conserved. Tracking does not modify spatial states, nonlinear unknown dimensions or stopping scales.

6 Execution and organization

Execution injects the vector field and its optional Jacobian into a fresh run; no method retains the dynamics object. Initialization binds the method's recipe, `DoubleFormulation` and configured algebraic solver once. Solver controls include tolerances, iteration limits, differentiation and iteration callbacks. The independent energy observer owns momentum and its sampled history. The ordinary implicit execution is

$$\texttt{advance} \longrightarrow \texttt{solve_projection} \longrightarrow \texttt{compose} \longrightarrow \texttt{A/B/T/C}.$$

All ABBA and BM4 presets inherit their advance from the shared palindromic classes; the recipe selects their elementary operations. Public configurations live in `abba.py` and `bm4.py`; common numerical operations live in `core/`, and optional event conversion lives in `observations.py`. The general integration coordinator remains responsible for main steps, saved times, shadow output samples, collection and observer dispatch. Observers receive independent snapshots in the existing public event types; BM4 events reuse accepted traces without repeating the spatial composition.

Current diagrams and module responsibilities are documented in `docs/models/extended/simulation/extended-simulation-architecture.md`. Tests verify public identity, coefficient and projection controls, signed nonautonomous maps, exact tangents and passive tracking. ABBA6 tests also verify sixth-order convergence on a nonlinear nonautonomous field with both root formulations and solvers. Historical ABBA6 results with seven separate projections represent a different map and must be recomputed for comparisons.