

BM4Midpoint: Arithmetic Diagonal Projection

GC2D numerical methods

1 Complete-cycle definition

For $\dot{z} = f(t, z)$, embed the accepted state as $Ez = (z, z)$ and define $P(u, v) = (u + v)/2$. One step is

$$z_{n+1} = P\Psi_h(Ez_n),$$

where Ψ_h is the same unprojected twelve-stage BM4 composition used by BM4Implicit. No projection is applied between stages and no multiplier is solved. “Midpoint” refers to the final arithmetic mean.

The coefficient vector is $(b_1, \dots, b_6, b_6, \dots, b_1)$, where

$$\begin{aligned} b_1 &= 0.0792036964311957, & b_2 &= 0.1303114101821663, \\ b_3 &= 0.2228614958676077, & b_4 &= -0.3667132690474257, \\ b_5 &= 0.3246481886897062, & b_6 &= 0.1096884778767498. \end{aligned}$$

The sum is one to floating-point accuracy. Starting with the adjoint, stages alternate adjoint and direct maps with signed duration $d_j = b_j h$. If $s_j = t_n + \sum_{i < j} d_i$, an adjoint stage uses s_j and a direct stage uses $s_j + d_j$. The negative coefficient also moves the stage clock backwards.

At the chosen evaluation time, let

$$A_d(u, v) = (u, v + df(t, u)), \quad B_d(u, v) = (u + df(t, v), v).$$

Let C_d be the exact GC harmonic mixing flow: it preserves the copy mean and rotates the spatial difference through angle $2\omega d$. The direct map is $C_d \circ B_d \circ A_d$; its adjoint is $B_d \circ A_d \circ C_d$. Each stage evaluates the vector field twice. Thus one complete step uses 24 evaluations and one arithmetic projection.

2 State and energy conventions

Twelve direct/adjoint stages followed by one spatial arithmetic mean. Energy tracking is optional and passive. The two shear contributions in each stage update a summed momentum. Dividing this accumulator by two gives the physical kappa normalization. The common state conventions are specified below.

3 Order and geometric limitations

On the invariant diagonal of the exact extended flow, the two copies follow the original dynamics and the harmonic coupling vanishes. For a smooth field, the fourth-order unprojected composition therefore satisfies

$$\Psi_h(Ez) = E\varphi_h(z) + O(h^5).$$

Since $PE = I$ and P is linear, the projected method has local error $O(h^5)$ and global order four under the usual stability assumptions. The passive energy coordinates do not change the physical map or its order.

Arithmetic projection discards the copy difference. Consequently, symmetry or symplecticity of a base composition alone does not establish exact reversibility or symplecticity of the accepted physical map. This method makes neither claim. The recorded infinity norm $\|u_f - v_f\|_\infty$ measures the discarded separation before projection.

4 Executable checks

The test suite compares the physical map with an independent signed-stage replay and checks fourth-order convergence against an exact rotation and a tightly integrated nonautonomous reference with energy tracking both disabled and enabled. It also checks momentum normalization against an augmented reference ODE, particle batching, persistent observer callbacks, and independence from off-grid output sampling. Nonlinear iterations are absent by construction.

5 Four state formulations and passive energy tracking

For one planar particle the supported internal representations are

physical	$z = (x, y)$	$\mathbb{R}^2$,
physical with energy	(z, t, κ)	$\mathbb{R}^4$,
duplicated	$(u, v), \quad u, v \in \mathbb{R}^2$	$\mathbb{R}^4$,
duplicated with energy	(u, v, t, κ)	$\mathbb{R}^6$.

Classical methods use the physical rows; ABBA and BM4 use the duplicated rows. `track_energy` selects the energy variant. For N planar particles the respective dimensions are $2N$, $4N$, $4N$, and $6N$: one time entry and one momentum per particle. All time entries equal the integration time. Classical full-cyclotron runs instead use $4N$ or $6N$ entries. Accepted spatial copies coincide; only one physical copy is exported. Every coordinate block contains N values in component-major order. The formulation validates particle times and aligns output round-off; the integrator does not interpret coordinate indices.

The normalized passive balance is

$$\dot{\kappa} = -\partial_t H(t, z), \quad \kappa(t_0) = 0, \quad \Delta K(t) = H(t, z(t)) + \kappa(t) - H(t_0, z_0).$$

Splitting accumulators from both copies are divided by two. The physical Hamiltonian may vary with time; ΔK measures its computed balance. Tracking leaves physical trajectories, nonlinear stopping rules and adaptive acceptance unchanged. Dynamics must supply the momentum derivative, including zero for an autonomous Hamiltonian.

Hairer’s constraint is solely $u - v = 0$: the reduced multiplier has $2N$ entries and ABBA’s simultaneous spatial solve has $6N$ unknowns. Neither t nor κ is projected. The former eight-component projection is retired; `state_extension='fully_extended'` raises migration guidance.

Dynamics, formulation, method and integration remain separate. The shared state classes own layouts, diagonal embedding and history extraction in

```
src/formulations/state.py
```

Each integration initializes a fresh method instance. Observers receive physical states; energy histories are in `Solution.diagnostics`. The `extended_time` history has shape (N, n_{samples}) . Fixed-step shadow samples leave main states and counters unchanged. Adaptive energy quadrature uses accepted dense output; its backend state and Jacobian remain physical.

6 Shared extended-space implementation

The public method names and constructor defaults are unchanged. Their code now lives in `src/methods/extended/`. Public exports from `simulation` remain available; former ABBA and BM4 import adapters have been removed. The shared execution is `advance`, `solve_projection`, `compose`, then `direct_map` or `adjoint_map`. Arithmetic midpoint variants replace the nonlinear projection by one final average.

The composition retains signed shear inputs for exact particle-block tangents, passive energy quadrature, and existing observation records. Newton and Broyden share the spatial projection equations; energy is accumulated only from the accepted trace, with the existing factor $1/2$ for physical κ . Adjoint stages evaluate at the start of their signed substep, direct stages at its end. The refactoring preserves the coefficients and projection placement: ABBA4 and BM4 have one outer projection, while ABBA6 retains seven projected ABBA2 pairs. This internal family name does not enable the retired full time/momentum projection.

The implementation architecture and current diagrams are documented in `docs/models/extended/simulation/extended-simulation-architecture.md`.