

ABBA and BM4 in Extended Space

Common composition, spatial projection and passive energy

1 Scope and unchanged public methods

The directory `src/methods/extended/` contains the common implementation of the existing ABBA and BM4 methods. It introduces no public method named `Extended`. Public class names, constructor defaults, exports from `simulation`, observation records and diagnostic keys are retained. Internal imports use the defining modules. Old ABBA and BM4 import adapters and historical `simulation.methods` routes have been removed.

For N independent planar particles, the physical state $z \in \mathbb{R}^{2N}$ is packed in component-major order. The splitting acts on two copies $Y = (u, v) \in \mathbb{R}^{4N}$. Accepted copies coincide. Optional energy tracking adds N time entries and N physical conjugate momenta κ , giving a $6N$ -component accepted internal workspace. These auxiliary variables never enter a spatial projection equation. Physical output remains in $\mathbb{R}^{2N}$.

2 Direct and adjoint compositions

Let χ_s denote a first-order map on the extended spatial workspace, with adjoint $\chi_s^* = \chi_{-s}^{-1}$, including the appropriate time translation for nonautonomous dynamics. A recipe with $2r$ stages is

$$\Psi_h = \chi_{b_{2r}h} \circ \chi_{b_{2r-1}h}^* \circ \cdots \circ \chi_{b_2h} \circ \chi_{b_1h}^*.$$

In execution order the stages alternate adjoint, direct. The weights obey $b_j = b_{2r+1-j}$ and $\sum_j b_j = 1$. Reflection exchanges each map with its adjoint. These properties are checked by the recipe constructor; additional order conditions remain necessary for higher-order accuracy.

With stage clock τ and signed duration $s = b_j h$, the adjoint evaluates at τ , the direct map at $\tau + s$, and the clock subsequently becomes $\tau + s$. Negative coefficients reverse the clock as well as the subflow. For uncoupled GC maps, the pair with weights $(1/2, 1/2)$ reproduces the endpoint-time A–B–B–A construction. Each direct or adjoint map contains two spatial shears. BM4 can additionally apply the existing exact harmonic coupling of the two spatial copies, in reversed operation order for the adjoint.

ABBA4 uses three unprojected ABBA pairs scaled by

$$\gamma = \frac{1}{2 - 2^{1/3}}, \quad \delta = -\frac{2^{1/3}}{2 - 2^{1/3}}, \quad (\gamma, \delta, \gamma).$$

Its six direct/adjoint weights are $(\gamma/2, \gamma/2, \delta/2, \delta/2, \gamma/2, \gamma/2)$. BM4 retains its twelve mirrored weights and its own fourth-order recipe. ABBA6 retains seven *projected* ABBA2 steps with the existing Yoshida weights. A single projection around fourteen unprojected stages would be a different method and is not introduced by this refactoring.

3 One common spatial projection equation

Write $m = 2N$, $Ez = (z, z)$, $G = [I, -I]$ and $N_0 = G^T$. The reduced Hairer projection finds $\mu \in \mathbb{R}^m$ such that

$$r(\mu) = G\Psi_h(Ez + N_0\mu) + 2\mu = 0.$$

After convergence the two corrected outputs are averaged. With $P = \frac{1}{2}[I, I]$, the accepted state is

$$z_+ = P(\Psi_h(Ez + N_0\mu) + N_0\mu).$$

The same equation applies to ABBA2, the complete ABBA4 composition, and BM4; only Ψ_h changes. ABBA6 solves it separately around each base pair.

The simultaneous ABBA formulation solves for $(u_+, v_+, \mu) \in \mathbb{R}^{3m}$:

$$\begin{pmatrix} u_+ - \mu - [\Psi_h(Ez + N_0\mu)]_u \\ v_+ + \mu - [\Psi_h(Ez + N_0\mu)]_v \\ u_+ - v_+ \end{pmatrix} = 0.$$

Arithmetic midpoint variants instead start from Ez , apply the complete recipe, and average once. They do not solve a nonlinear equation; arithmetic projection does not imply exact symplecticity or reversibility.

4 Nonlinear solution and exact tangents

All spatial roots use the unchanged stopping threshold

$$\text{atol} + \text{rtol} \max(1, \|z\|_\infty).$$

Newton and good Broyden share the convergence machinery in `src/methods/_nonlinear.py`. Broyden's initial reduced Jacobian is $4I$, the exact zero-step value of Dr . The simultaneous zero-step block matrix is

$$\begin{pmatrix} I_{2m} & -2N_0 \\ G & 0 \end{pmatrix}.$$

Broyden subsequently uses residual secants and evaluates no analytic Jacobian.

If $M = D\Psi_h$, Newton's reduced Jacobian is $GMN_0 + 2I$. The common composition retains the input to each spatial shear. Exact stage tangents are evaluated at those inputs and multiplied in execution order, $M = M_{2r} \cdots M_1$, without replaying vector-field stages. Independence of the particles gives 4×4 blocks for M , 2×2 reduced Newton blocks, and 6×6 simultaneous blocks. BM4 also retains its optional dense centered-difference differentiation of the complete map.

5 Passive energy from accepted stages

The dynamics supply $H(t, z)$ and $-\partial_t H(t, z)$. For the accepted spatial trace, let (t_j, s_j, z_j) denote the evaluation time, signed duration and physical source state of each individual shear. The shared quadrature is

$$\Delta\kappa = \frac{1}{2} \sum_j s_j [-\partial_t H(t_j, z_j)].$$

There are four shear points for an ABBA2 pair, twelve for ABBA4 and 24 for BM4. The factor $1/2$ converts summed doubled momentum to the physical variable κ . Spatial harmonic coupling does not change this passive accumulator. Energy derivatives are evaluated only after the spatial root converges, from its retained trace; trial roots do not evaluate energy. The accepted diagnostic is

$$H(t_n, z_n) + \kappa_n - H(t_0, z_0), \quad \kappa_0 = 0.$$

This is an energy-balance diagnostic, not a claim that the time-dependent physical Hamiltonian is conserved. Tracking does not modify spatial states, nonlinear unknown dimensions or stopping scales.

6 Execution and compatibility

Initialization binds the method's recipe and projection options once. The ordinary implicit execution is

`advance` $\longrightarrow$ `solve_projection` $\longrightarrow$ `compose` $\longrightarrow$ `direct_map` / `adjoint_map`.

ABBA6 repeats its complete projected-pair operation seven times. The general integration coordinator remains responsible for main steps, saved times, shadow output samples, collection and observer dispatch. Observers receive independent snapshots in the existing public event types; BM4 events reuse accepted traces without repeating the spatial composition.

Current diagrams and module responsibilities are documented in `docs/models/extended/simulation/extended-simulation-architecture.md`. Tests verify public identity, coefficient and projection controls, signed nonautonomous maps, exact tangents, passive tracking and unchanged numerical contracts. Floating-point roundoff differences can arise from equivalent algebraic evaluation order; no mathematical recipe is changed.