

Two-stage Gauss–Legendre Method

Exact one-step Jacobian, symplecticity, and symmetry

Fourth-order collocation method for canonical Hamiltonian systems

1 Hamiltonian convention

Let $z \in \mathbb{R}^{2d}$ and consider the possibly time-dependent canonical Hamiltonian system

$$\dot{z} = f(t, z) = \Omega^{-1} \nabla_z H(t, z), \quad \Omega = \begin{pmatrix} 0 & I_d \\ -I_d & 0 \end{pmatrix}. \quad (1)$$

For the component-major guiding-centre state

$$z = (x_1, \dots, x_d, y_1, \dots, y_d)^T,$$

this convention gives

$$f(t, z) = \begin{pmatrix} -\nabla_y H(t, z) \\ \nabla_x H(t, z) \end{pmatrix}.$$

At any fixed time and state, let $F = D_z f(t, z)$. Since $\nabla_z^2 H$ is symmetric,

$$F^T \Omega + \Omega F = 0. \quad (2)$$

This identity is the infinitesimal Hamiltonian condition used below. The proof remains valid for a time-dependent Hamiltonian because the stage times do not depend on the initial state z_n .

2 Gauss–Legendre stages

Set

$$r = \frac{\sqrt{3}}{6}, \quad c_1 = \frac{1}{2} - r, \quad c_2 = \frac{1}{2} + r.$$

The Butcher coefficients of the two-stage Gauss–Legendre method are

$$A = \begin{pmatrix} \frac{1}{4} & \frac{1}{4} - r \\ \frac{1}{4} + r & \frac{1}{4} \end{pmatrix}, \quad b = \begin{pmatrix} \frac{1}{2} \\ \frac{1}{2} \end{pmatrix}. \quad (3)$$

For one step of length h , the stage states satisfy

$$Z_i = z_n + h \sum_{j=1}^2 a_{ij} f(t_n + c_j h, Z_j), \quad i = 1, 2, \quad (4)$$

and the accepted state is

$$z_{n+1} = \Phi_{h,t_n}(z_n) = z_n + h \sum_{i=1}^2 b_i f(t_n + c_i h, Z_i). \quad (5)$$

The method is implicit: equations (4) determine (Z_1, Z_2) simultaneously.

3 Exact Jacobian of the stage root

Introduce the stage field Jacobians and stage sensitivities

$$F_i := D_z f(t_n + c_i h, Z_i), \quad S_i := \frac{\partial Z_i}{\partial z_n}. \quad (6)$$

Differentiating (4) with respect to z_n gives

$$S_i = I_{2d} + h \sum_{j=1}^2 a_{ij} F_j S_j. \quad (7)$$

Equivalently,

$$\underbrace{\begin{pmatrix} I_{2d} - \frac{h}{4} F_1 & -h \left(\frac{1}{4} - r \right) F_2 \\ -h \left(\frac{1}{4} + r \right) F_1 & I_{2d} - \frac{h}{4} F_2 \end{pmatrix}}_{M=D_{(Z_1, Z_2)} R} \begin{pmatrix} S_1 \\ S_2 \end{pmatrix} = \begin{pmatrix} I_{2d} \\ I_{2d} \end{pmatrix}. \quad (8)$$

Here M is the Jacobian of the coupled stage residual. It is also the matrix used by an exact Newton linearization, but it is *not* the Jacobian of the physical one-step map.

Assuming that the converged stage root is regular, the implicit function theorem gives

$$\begin{pmatrix} S_1 \\ S_2 \end{pmatrix} = M^{-1} \begin{pmatrix} I_{2d} \\ I_{2d} \end{pmatrix}. \quad (9)$$

In an implementation, equation (8) should be solved directly; the inverse in (9) is only notation.

4 Exact Jacobian of the physical step

Define the Jacobian of the map (5) by

$$\mathcal{J}_n := D_{z_n} \Phi_{h, t_n}(z_n).$$

Differentiating the accepted update and using (6) yields

$$\boxed{\mathcal{J}_n = I_{2d} + h \sum_{i=1}^2 b_i F_i S_i = I_{2d} + \frac{h}{2} (F_1 S_1 + F_2 S_2)}. \quad (10)$$

Combining (9) and (10), the same Jacobian can be written as

$$\boxed{\mathcal{J}_n = I_{2d} + h \begin{pmatrix} \frac{1}{2} F_1 & \frac{1}{2} F_2 \end{pmatrix} M^{-1} \begin{pmatrix} I_{2d} \\ I_{2d} \end{pmatrix}}. \quad (11)$$

This is the exact tangent of the ideal converged root. It differentiates the mathematical Gauss–Legendre map, not a finite Newton stopping rule.

5 Direct proof of symplecticity with the step Jacobian

The one-step map is symplectic if and only if

$$\mathcal{J}_n^T \Omega \mathcal{J}_n = \Omega. \quad (12)$$

Set

$$K_i := F_i S_i. \quad (13)$$

Then equations (10) and (7) become

$$\mathcal{J}_n = I_{2d} + h \sum_i b_i K_i, \quad I_{2d} = S_i - h \sum_j a_{ij} K_j. \quad (14)$$

Expanding the symplecticity defect gives

$$\begin{aligned} \mathcal{J}_n^T \Omega \mathcal{J}_n - \Omega &= h \sum_i b_i \left(K_i^T \Omega + \Omega K_i \right) \\ &\quad + h^2 \sum_{i,j} b_i b_j K_i^T \Omega K_j. \end{aligned} \quad (15)$$

By the Hamiltonian identity (2),

$$\begin{aligned} K_i^T \Omega S_i + S_i^T \Omega K_i &= S_i^T \left(F_i^T \Omega + \Omega F_i \right) S_i \\ &= 0. \end{aligned} \quad (16)$$

Using the second identity in (14) on both occurrences of I_{2d} therefore gives

$$K_i^T \Omega + \Omega K_i = -h \sum_j a_{ij} \left(K_i^T \Omega K_j + K_j^T \Omega K_i \right). \quad (17)$$

Substitute (17) into (15). Renaming i and j in the second term arising from (17) produces

$$\boxed{\mathcal{J}_n^T \Omega \mathcal{J}_n - \Omega = h^2 \sum_{i,j} (b_i b_j - b_i a_{ij} - b_j a_{ji}) K_i^T \Omega K_j.} \quad (18)$$

Consequently, the defect vanishes whenever the Runge–Kutta coefficients obey

$$\boxed{b_i a_{ij} + b_j a_{ji} = b_i b_j, \quad i, j = 1, 2.} \quad (19)$$

6 Verification for the Gauss–Legendre coefficients

For the diagonal terms in (19),

$$2b_1a_{11} - b_1^2 = 2\left(\frac{1}{2}\right)\left(\frac{1}{4}\right) - \frac{1}{4} = 0, \quad 2b_2a_{22} - b_2^2 = 0. \quad (20)$$

For the off-diagonal terms,

$$\begin{aligned} b_1a_{12} + b_2a_{21} - b_1b_2 &= \frac{1}{2}\left(\frac{1}{4} - r\right) + \frac{1}{2}\left(\frac{1}{4} + r\right) - \frac{1}{4} \\ &= 0. \end{aligned} \quad (21)$$

The case $(i, j) = (2, 1)$ is the same equality with the two summands exchanged. Thus every coefficient multiplying $K_i^T \Omega K_j$ in (18) is zero, and hence

$$\boxed{\mathcal{J}_n^T \Omega \mathcal{J}_n = \Omega.} \quad (22)$$

Therefore the two-stage Gauss–Legendre method is symplectic when its implicit stage equations are solved exactly.

7 Symmetry and time reversibility

Symmetry is a property distinct from symplecticity. The method is symmetric when its backward step is exactly the inverse of its forward step. For the possibly time-dependent problem (1), this means

$$\boxed{\Phi_{-h, t_n+h} \circ \Phi_{h, t_n} = \text{Id}.} \quad (23)$$

For an autonomous problem, equation (23) reduces to $\Phi_{-h} = \Phi_h^{-1}$.

For the forward step, write

$$f_1 = f(t_n + c_1 h, Z_1), \quad f_2 = f(t_n + c_2 h, Z_2). \quad (24)$$

Equations (4) and (5) are then

$$\begin{aligned} Z_1 &= z_n + h(a_{11}f_1 + a_{12}f_2), \\ Z_2 &= z_n + h(a_{21}f_1 + a_{22}f_2), \end{aligned} \quad (25)$$

$$z_{n+1} = z_n + h(b_1f_1 + b_2f_2). \quad (26)$$

Now start from z_{n+1} at time $t_n + h$ and apply the same method with step $-h$. Its stages $\hat{Z}_1, \hat{Z}_2$ satisfy

$$\hat{Z}_i = z_{n+1} - h \sum_{j=1}^2 a_{ij} f(t_n + h - c_j h, \hat{Z}_j). \quad (27)$$

The Gauss nodes are reflected about the midpoint:

$$1 - c_1 = c_2, \quad 1 - c_2 = c_1. \quad (28)$$

Thus the backward stage times are the forward stage times in reverse order. We shall verify that the backward stages are

$$\boxed{\hat{Z}_1 = Z_2, \quad \hat{Z}_2 = Z_1.} \quad (29)$$

Under the identification (29), the first backward stage equation becomes

$$\begin{aligned} \hat{Z}_1 &= z_{n+1} - h(a_{11}f_2 + a_{12}f_1) \\ &= z_n + h[(b_1 - a_{12})f_1 + (b_2 - a_{11})f_2]. \end{aligned} \quad (30)$$

For the coefficients (3),

$$b_1 - a_{12} = a_{21}, \quad b_2 - a_{11} = a_{22}. \quad (31)$$

Consequently,

$$\hat{Z}_1 = z_n + h(a_{21}f_1 + a_{22}f_2) = Z_2.$$

Likewise, the second backward stage satisfies

$$\begin{aligned} \hat{Z}_2 &= z_{n+1} - h(a_{21}f_2 + a_{22}f_1) \\ &= z_n + h[(b_1 - a_{22})f_1 + (b_2 - a_{21})f_2]. \end{aligned} \quad (32)$$

Since

$$b_1 - a_{22} = a_{11}, \quad b_2 - a_{21} = a_{12}, \quad (33)$$

we obtain

$$\hat{Z}_2 = z_n + h(a_{11}f_1 + a_{12}f_2) = Z_1.$$

The proposed reversed stages therefore solve the coupled backward stage equations exactly.

The output of the backward step is now

$$\begin{aligned}
 \hat{z}_n &= z_{n+1} - h(b_1 f_2 + b_2 f_1) \\
 &= z_n + h(b_1 f_1 + b_2 f_2) - h(b_1 f_2 + b_2 f_1) \\
 &= z_n,
 \end{aligned} \tag{34}$$

where the last equality uses $b_1 = b_2 = 1/2$. This proves (23).

More generally, an s -stage Runge–Kutta method is symmetric when its coefficients are invariant under stage reversal:

$$\boxed{c_i = 1 - c_{s+1-i}, \quad b_i = b_{s+1-i}, \quad a_{ij} = b_{s+1-j} - a_{s+1-i, s+1-j}.} \tag{35}$$

The coefficients in (3) satisfy all three identities for $s = 2$.

Differentiating the map identity (23) also gives the Jacobian form of reversibility:

$$\boxed{D\Phi_{-h, t_n+h}(z_{n+1}) D\Phi_{h, t_n}(z_n) = I_{2d}.} \tag{36}$$

Thus the backward-step Jacobian is exactly the inverse of the forward-step Jacobian. As with symplecticity, a finite Newton tolerance and floating-point round-off can produce a small measured reversibility defect even though the ideal implicit method is exactly symmetric.

8 Consequences and numerical interpretation

For one guiding-centre particle, $\mathcal{J}_n \in \mathbb{R}^{2 \times 2}$ and

$$\mathcal{J}_n^T \Omega \mathcal{J}_n = \det(\mathcal{J}_n) \Omega.$$

Equation (22) therefore implies

$$\det(\mathcal{J}_n) = 1,$$

so the one-step map preserves oriented area. In dimensions greater than two, unit determinant is only a consequence of symplecticity, not an equivalent test.

If $\mathcal{J}_n$ denotes the local Jacobian at step n , the accumulated flow tangent after N steps is

$$\mathcal{P}_N = \mathcal{J}_{N-1} \cdots \mathcal{J}_1 \mathcal{J}_0.$$

Since a product of symplectic matrices is symplectic,

$$\mathcal{P}_N^T \Omega \mathcal{P}_N = \Omega.$$

The algebraic result concerns the exact root and the exact tangent (10). In floating-point computations, a finite Newton tolerance, round-off, or a finite-difference approximation of the map Jacobian produces a small nonzero measured defect

$$\left\| \mathcal{J}_n^T \Omega \mathcal{J}_n - \Omega \right\|.$$

That numerical floor does not contradict the exact symplecticity proved above.

References

- J. M. Sanz-Serna, “Runge–Kutta schemes for Hamiltonian systems,” *BIT Numerical Mathematics*, 28 (1988), 877–883.
- E. Hairer, C. Lubich, and G. Wanner, *Geometric Numerical Integration*, second edition, Springer, 2006.