

Intro to Analytics Modeling HW 3

2024-06-05

Question 7.1:

Describe a situation or problem from your job, everyday life, current events, etc., for which exponential smoothing would be appropriate. What data would you need? Would you expect the value of α (the first smoothing parameter) to be closer to 0 or 1, and why?.

Answer: In my current company, we conduct R&D on lithium metal batteries. When we test our battery cells, we charge and discharge them, where one cycle consists of one charging and one discharging step. While there are several metrics to observe, such as voltage, current, capacity, and temperature, using voltage for exponential smoothing can be particularly helpful for monitoring changes over cycles. This can indirectly indicate capacity fade and identify anomalies. Especially, it can sometimes detect safety-critical failures that have a high potential to lead to thermal runaway. Since the time series data using voltage is normally cyclic, it is not expected to have high randomness; therefore, the value of α tends to be closer to 1.

Importing Libraries

```
# library(ggplot2)
library(knitr)
library(tidyr)
library(stats)
library(caret)
```

```
## Loading required package: ggplot2
```

```
## Loading required package: lattice
```

Question 7.2:

Using the 20 years of daily high temperature data for Atlanta (July through October) from Question 6.2 (file temps.txt), build and use an exponential smoothing model to help make a judgment of whether the unofficial end of summer has gotten later over the 20 years. (Part of the point of this assignment is for you to think about how you might use exponential smoothing to answer this question. Feel free to combine it with other models if you'd like to. There's certainly more than one reasonable approach.)

Note: in R, you can use either `HoltWinters` (simpler to use) or the `smooth` package's `es` function (harder to use, but more general). If you use `es`, the Holt-Winters model uses `model="AAM"` in the function call (the first and second constants are used "A"dditively, and the third (seasonality) is used "M"ultiplicatively; the documentation doesn't make that clear).

Read Data

```
temp_data <- read.table(  
  "~/Desktop/ISYE-6501/week 3 Homework-Summer24/week 3 data-summer/temps.txt",  
  stringsAsFactors = FALSE,  
  header=TRUE  
)  
head(temp_data)
```

```
##      DAY X1996 X1997 X1998 X1999 X2000 X2001 X2002 X2003 X2004 X2005 X2006 X2007  
## 1 1-Jul   98    86    91    84    89    84    90    73    82    91    93    95  
## 2 2-Jul   97    90    88    82    91    87    90    81    81    89    93    85  
## 3 3-Jul   97    93    91    87    93    87    87    87    86    86    93    82  
## 4 4-Jul   90    91    91    88    95    84    89    86    88    86    91    86  
## 5 5-Jul   89    84    91    90    96    86    93    80    90    89    90    88  
## 6 6-Jul   93    84    89    91    96    87    93    84    90    82    81    87  
##      X2008 X2009 X2010 X2011 X2012 X2013 X2014 X2015  
## 1      85    95    87    92   105    82    90    85  
## 2      87    90    84    94    93    85    93    87  
## 3      91    89    83    95    99    76    87    79  
## 4      90    91    85    92    98    77    84    85  
## 5      88    80    88    90   100    83    86    84  
## 6      82    87    89    90    98    83    87    84
```

```
years <- names(temp_data[,2:ncol(temp_data)])  
df_temps <- pivot_longer(temp_data, cols=years, names_to="Year", values_to="Temperature")
```

```
## Warning: Using an external vector in selections was deprecated in tidysselect 1.1.0.  
## i Please use `all_of()` or `any_of()` instead.  
## # Was:  
## data %>% select(years)  
##  
## # Now:  
## data %>% select(all_of(years))  
##  
## See <https://tidysselect.r-lib.org/reference/faq-external-vector.html>.  
## This warning is displayed once every 8 hours.  
## Call `lifecycle::last_lifecycle_warnings()` to see where this warning was  
## generated.
```

```
# Data cleansing  
df_temps$Year <- as.integer(gsub("X", "", df_temps$Year))  
df_temps$ts <- as.Date(paste(df_temps$DAY, df_temps$Year), format = "%d-%b %Y")  
df_temps$Month <- as.integer(format(df_temps$ts, "%m"))  
df_temps$Temperature <- as.numeric(df_temps$Temperature)  
  
df_temps <- df_temps[order(df_temps$ts), ]  
df_temps <- df_temps[, c("ts", "Temperature", "Year", "Month")]  
df_temps
```

```
## # A tibble: 2,460 x 4
```

```
##      ts      Temperature Year Month
##      <date>          <dbl> <int> <int>
##  1 1996-07-01          98  1996    7
##  2 1996-07-02          97  1996    7
##  3 1996-07-03          97  1996    7
##  4 1996-07-04          90  1996    7
##  5 1996-07-05          89  1996    7
##  6 1996-07-06          93  1996    7
##  7 1996-07-07          93  1996    7
##  8 1996-07-08          91  1996    7
##  9 1996-07-09          93  1996    7
## 10 1996-07-10          93  1996    7
## # i 2,450 more rows
```

CUSUM Recap

```
years <- unique(df_temps$Year)
end_summer_dates <- as.Date(character(0))
for (year in years) {
  temporary_table <- df_temps[df_temps$Year == year,]
  mean_overall <- mean(temporary_table$Temperature)
  std_overall <- sd(temporary_table$Temperature)
  C <- std_overall
  T <- 3 * std_overall #threshold set to 3 sigma

  temporary_table[1,"St"] <- 0 # Assuming S_0 is 0

  for(t in 2:nrow(temporary_table)) {
    temporary_table[t, "St"] <-
      max(0, (temporary_table[t-1, "St"] + mean_overall - temporary_table[t, "Temperature"] - C)$St)
  }

  end_summer_dates <- c(
    end_summer_dates,
    as.Date(temporary_table[which(temporary_table$St>T),][1,$ts)
  )
}
end_summer_dates <- df_temps[df_temps$ts %in% end_summer_dates, ][, c("ts", "Temperature")]

end_summer_dates$Year <- as.integer(format(end_summer_dates$ts, "%Y"))
end_summer_dates$month_date <- as.integer(format(end_summer_dates$ts, "%m%d"))
end_summer_dates$Year_norm <- end_summer_dates$Year / max(end_summer_dates$Year)
end_summer_dates$month_date_norm <- end_summer_dates$month_date / max(end_summer_dates$month_date)
end_summer_dates
```

```
## # A tibble: 20 x 6
##      ts      Temperature Year month_date Year_norm month_date_norm
##      <date>          <dbl> <int>      <int>      <dbl>          <dbl>
##  1 1996-10-02          72  1996      1002      0.991          0.978
##  2 1997-10-16          57  1997      1016      0.991          0.991
##  3 1998-10-10          73  1998      1010      0.992          0.985
##  4 1999-10-20          60  1999      1020      0.992          0.995
```


##	5	2000-10-08	55	2000	1008	0.993	0.983
##	6	2001-10-18	64	2001	1018	0.993	0.993
##	7	2002-10-16	66	2002	1016	0.994	0.991
##	8	2003-10-02	68	2003	1002	0.994	0.978
##	9	2004-10-13	64	2004	1013	0.995	0.988
##	10	2005-10-24	56	2005	1024	0.995	0.999
##	11	2006-10-16	59	2006	1016	0.996	0.991
##	12	2007-10-25	61	2007	1025	0.996	1
##	13	2008-10-20	66	2008	1020	0.997	0.995
##	14	2009-10-16	61	2009	1016	0.997	0.991
##	15	2010-10-03	68	2010	1003	0.998	0.979
##	16	2011-10-21	63	2011	1021	0.998	0.996
##	17	2012-10-11	75	2012	1011	0.999	0.986
##	18	2013-10-20	70	2013	1020	0.999	0.995
##	19	2014-10-19	73	2014	1019	1.00	0.994
##	20	2015-10-04	70	2015	1004	1	0.980

Explanation: This is a list of unofficial summer end dates for each year based on CUSUM algorithm. Note that mean and standard deviation for C and T are applied dynamically for each year separately as it's described in the for loop.

```
x <- as.vector(end_summer_dates$Year_norm)
y <- as.vector(end_summer_dates$month_date_norm)

linear_model <- lm(y ~ x)
intercept <- linear_model$coefficients[1]
slope <- linear_model$coefficients[2]

plot(end_summer_dates$Year_norm, end_summer_dates$month_date_norm)
# Add linear line
abline(a = intercept, b = slope, col = "red")
```



Explanation: The X-axis represents normalized years, and the Y-axis represents normalized month_date.

```
linear_model
```

```
##
## Call:
## lm(formula = y ~ x)
##
## Coefficients:
## (Intercept)          x
##      0.6246      0.3666
```

Observation: Based on how the CUSUM algorithm is applied (mean and standard deviation for C and T are applied dynamically for each year separately), it produces unofficial summer end dates for each year. When the result is displayed in a scatter plot along with a linear regression line, it indicates that the unofficial end of summer has gotten later over the 20 years.

Exponential Smoothing


```
# 123 days from July to October for the frequency
ts_data_all <- ts(df_temps$Temperature, start = min(df_temps$ts), frequency = 123)
# Plot the time series with continuous date x-axis
plot(
  ts_data_all,
```

```

xlab = "",
ylab = "Temperature",
main = "Time Series Plot for Years Between 1996 and 2015",
xaxt="n"
)

```

Time Series Plot for Years Between 1996 and 2015



Observation: There appears to be a cyclic pattern over the years, but it's not clear whether the overall trend is increasing or decreasing. Additionally, there is some level of randomness in daily high temperatures. Therefore, it makes sense to utilize triple exponential smoothing, specifically the Holt-Winters method.

```

# Define a function for an exponential smoothing model
build_hw_model = function(data, a, b, g, s) {
  hw_model = HoltWinters(
    data,
    alpha=a,
    beta=b,
    gamma=g,
    seasonal=s,
  )
  return(hw_model)
}


```

```


# A model that an exponential smoothing is applied
hw_model = build_hw_model(ts_data_all, 0.9, NULL, NULL, "m")
smoothed_values <- fitted(hw_model)

```


```
plot(ts_data_all)
lines(smoothed_values[,1], col="pink")
```



```
# A model that an exponential smoothing is applied
hw_model = build_hw_model(ts_data_all, 0.5, NULL, NULL, "m")
smoothed_values <- fitted(hw_model)
plot(ts_data_all)
lines(smoothed_values[,1], col="pink")
```



```
# A model that an exponential smoothing is applied
hw_model = build_hw_model(ts_data_all, 0.1, NULL, NULL, "m")
smoothed_values <- fitted(hw_model)
plot(ts_data_all)
lines(smoothed_values[,1], col="pink")
```

Observation: By applying different values for alpha (0.1, 0.5, and 0.9), it appears that 0.1 is the optimal value for exponential smoothing of the time series data.

```
print(paste("alpha:", hw_model$alpha, " beta:", hw_model$beta, "gamma:", hw_model$gamma))
```

```
## [1] "alpha: 0.1 beta: 0 gamma: 0.279818248767374"
```

Observation: With the model, there's no trend identified but a bit of seasonality identified.

```
summary(hw_model)
```

```
##          Length Class  Mode
## fitted      9348   mts    numeric
## x           2460   ts     numeric
## alpha         1  -none-  numeric
## beta          1  -none-  numeric
## gamma         1  -none-  numeric
## coefficients  125  -none-  numeric
## seasonal      1  -none-  character
## SSE           1  -none-  numeric
## call          6  -none-  call
```

```
hw_model$coefficients[1:2]
```

```
##          a          b
## 85.544930095 -0.004362918
```

Observation: Based on the current estimate of the baseline coefficient, which can be interpreted in a linear model, 'b' indicates the slope of the fitted curve. It appears to be close to zero, suggesting that there may be no significant increase or decrease in daily high temperatures over the 20-year period after the data is smoothed by exponential smoothing. However, it's important to note that this observation captures only the overall trend. We are interested in determining if there exists a trend specifically over the unofficial end-of-summer dates during the 20-year period. Let's recompute the unofficial summer end dates against the smoothed data by applying CUSUM algorithm.

CUSUM using Smoothed temperature

```
# exclude 1996 data since there's no smooth data for the first year because the seasonality coefficient
df_temps_new <- df_temps[df_temps$Year > 1996,]

# down sample the smoothed temperature by applying a frequency of 123
# smoothed_values comes from hw_model with an alpha of 0.1
downsampled_smoothed_data <- downSample(x=smoothed_values, factor(df_temps_new$ts))
df_temps_new$t_hat <- downsampled_smoothed_data$xhat

# Compute CUSUM for each year against the smoothed data
years <- unique(df_temps_new$Year)
end_summer_dates <- as.Date(character(0))
for (year in years) {
  temporary_table <- df_temps_new[df_temps_new$Year == year,]
  mean_overall <- mean(temporary_table$t_hat)
  std_overall <- sd(temporary_table$t_hat)
  C <- std_overall
  T <- 3 * std_overall #threshold set to 3 sigma

  temporary_table[1,"St"] <- 0 # Assuming S_0 is 0

  for(t in 2:nrow(temporary_table)) {
    temporary_table[t, "St"] <-
      max(0, (temporary_table[t-1, "St"] + mean_overall - temporary_table[t, "t_hat"] - C)$St)
  }

  end_summer_dates <- c(
    end_summer_dates,
    as.Date(temporary_table[which(temporary_table$St>T),][1,$ts)
  )
}
end_summer_dates <- df_temps_new[df_temps_new$ts %in% end_summer_dates, ][, c("ts", "t_hat")]

end_summer_dates$Year <- as.integer(format(end_summer_dates$ts, "%Y"))
end_summer_dates$month_date <- as.integer(format(end_summer_dates$ts, "%m%d"))
end_summer_dates$Year_norm <- end_summer_dates$Year / max(end_summer_dates$Year)
end_summer_dates$month_date_norm <- end_summer_dates$month_date / max(end_summer_dates$month_date)
end_summer_dates

## # A tibble: 19 x 6
```

##	ts	t_hat	Year	month_date	Year_norm	month_date_norm
##	<date>	<dbl>	<int>	<int>	<dbl>	<dbl>
##	1 1997-10-01	66.9	1997	1001	0.991	0.978
##	2 1998-10-20	74.7	1998	1020	0.992	0.997
##	3 1999-10-07	68.2	1999	1007	0.992	0.984
##	4 2000-10-06	70.2	2000	1006	0.993	0.983
##	5 2001-10-07	68.1	2001	1007	0.993	0.984
##	6 2002-10-19	69.3	2002	1019	0.994	0.996
##	7 2003-10-08	68.5	2003	1008	0.994	0.985
##	8 2004-10-15	68.8	2004	1015	0.995	0.992
##	9 2005-10-11	76.3	2005	1011	0.995	0.988
##	10 2006-10-17	71.2	2006	1017	0.996	0.994
##	11 2007-10-14	74.4	2007	1014	0.996	0.991
##	12 2008-10-23	68.8	2008	1023	0.997	1
##	13 2009-10-18	66.3	2009	1018	0.997	0.995
##	14 2010-10-10	75.7	2010	1010	0.998	0.987
##	15 2011-10-15	70.5	2011	1015	0.998	0.992
##	16 2012-10-16	73.9	2012	1016	0.999	0.993
##	17 2013-10-23	71.3	2013	1023	0.999	1
##	18 2014-10-17	74.9	2014	1017	1.00	0.994
##	19 2015-10-15	69.2	2015	1015	1	0.992


```

x <- as.vector(end_summer_dates$Year_norm)
y <- as.vector(end_summer_dates$month_date_norm)

linear_model <- lm(y ~ x)
intercept <- linear_model$coefficients[1]
slope <- linear_model$coefficients[2]

plot(end_summer_dates$Year_norm, end_summer_dates$month_date_norm)
# Add linear line
abline(a = intercept, b = slope, col = "red")

```



Explanation: The X-axis represents normalized years, and the Y-axis represents normalized month_date.

```
linear_model
```

```
##
## Call:
## lm(formula = y ~ x)
##
## Coefficients:
## (Intercept)          x
##      -0.1339       1.1300
```

Observation: Based on how the CUSUM algorithm is applied (mean and standard deviation for C and T are applied dynamically for each year separately) against the smoothed data, it produces unofficial summer end dates for each year again. When the result is displayed in a scatter plot along with a linear regression line, it clearly indicates that the unofficial end of summer has gotten later over the 20 years.

Question 8.1:

Describe a situation or problem from your job, everyday life, current events, etc., for which a linear regression model would be appropriate. List some (up to 5) predictors that you might use.

I have a small rechargeable desk fan, and I'm interested in assessing its battery health, specifically in terms of the capacity for a full charge. To achieve this, I believe that applying a linear regression model would be appropriate, considering several potential predictors:

- daily temperature
- hours of desk usage
- hours of daily charging time
- hours of daily operating time
- average speed of the fan (1 through 4)
- cumulative hours of usage

Question 8.2:

Using crime data from <http://www.statsci.org/data/general/uscrime.txt> (file `uscrime.txt`, description at <http://www.statsci.org/data/general/uscrime.html>), use regression (a useful R function is `lm` or `glm`) to predict the observed crime rate in a city with the following data:

- $M = 14.0$
- $So = 0$
- $Ed = 10.0$
- $Po1 = 12.0$
- $Po2 = 15.5$
- $LF = 0.640$
- $M.F = 94.0$
- $Pop = 150$
- $NW = 1.1$
- $U1 = 0.120$
- $U2 = 3.6$
- $Wealth = 3200$
- $Ineq = 20.1$
- $Prob = 0.04$
- $Time = 39.0$

Show your model (factors used and their coefficients), the software output, and the quality of fit.

Note that because there are only 47 data points and 15 predictors, you'll probably notice some overfitting. We'll see ways of dealing with this sort of problem later in the course.

Read Data

```
crime_data <- read.table("~/Desktop/ISYE-6501/week 3 Homework-Summer24/week 3 data-summer/uscrime.txt",
crime_data
```

##	M	So	Ed	Po1	Po2	LF	M.F	Pop	NW	U1	U2	Wealth	Ineq	Prob
## 1	15.1	1	9.1	5.8	5.6	0.510	95.0	33	30.1	0.108	4.1	3940	26.1	0.084602
## 2	14.3	0	11.3	10.3	9.5	0.583	101.2	13	10.2	0.096	3.6	5570	19.4	0.029599
## 3	14.2	1	8.9	4.5	4.4	0.533	96.9	18	21.9	0.094	3.3	3180	25.0	0.083401
## 4	13.6	0	12.1	14.9	14.1	0.577	99.4	157	8.0	0.102	3.9	6730	16.7	0.015801
## 5	14.1	0	12.1	10.9	10.1	0.591	98.5	18	3.0	0.091	2.0	5780	17.4	0.041399
## 6	12.1	0	11.0	11.8	11.5	0.547	96.4	25	4.4	0.084	2.9	6890	12.6	0.034201
## 7	12.7	1	11.1	8.2	7.9	0.519	98.2	4	13.9	0.097	3.8	6200	16.8	0.042100
## 8	13.1	1	10.9	11.5	10.9	0.542	96.9	50	17.9	0.079	3.5	4720	20.6	0.040099
## 9	15.7	1	9.0	6.5	6.2	0.553	95.5	39	28.6	0.081	2.8	4210	23.9	0.071697

##	10	14.0	0	11.8	7.1	6.8	0.632	102.9	7	1.5	0.100	2.4	5260	17.4	0.044498
##	11	12.4	0	10.5	12.1	11.6	0.580	96.6	101	10.6	0.077	3.5	6570	17.0	0.016201
##	12	13.4	0	10.8	7.5	7.1	0.595	97.2	47	5.9	0.083	3.1	5800	17.2	0.031201
##	13	12.8	0	11.3	6.7	6.0	0.624	97.2	28	1.0	0.077	2.5	5070	20.6	0.045302
##	14	13.5	0	11.7	6.2	6.1	0.595	98.6	22	4.6	0.077	2.7	5290	19.0	0.053200
##	15	15.2	1	8.7	5.7	5.3	0.530	98.6	30	7.2	0.092	4.3	4050	26.4	0.069100
##	16	14.2	1	8.8	8.1	7.7	0.497	95.6	33	32.1	0.116	4.7	4270	24.7	0.052099
##	17	14.3	0	11.0	6.6	6.3	0.537	97.7	10	0.6	0.114	3.5	4870	16.6	0.076299
##	18	13.5	1	10.4	12.3	11.5	0.537	97.8	31	17.0	0.089	3.4	6310	16.5	0.119804
##	19	13.0	0	11.6	12.8	12.8	0.536	93.4	51	2.4	0.078	3.4	6270	13.5	0.019099
##	20	12.5	0	10.8	11.3	10.5	0.567	98.5	78	9.4	0.130	5.8	6260	16.6	0.034801
##	21	12.6	0	10.8	7.4	6.7	0.602	98.4	34	1.2	0.102	3.3	5570	19.5	0.022800
##	22	15.7	1	8.9	4.7	4.4	0.512	96.2	22	42.3	0.097	3.4	2880	27.6	0.089502
##	23	13.2	0	9.6	8.7	8.3	0.564	95.3	43	9.2	0.083	3.2	5130	22.7	0.030700
##	24	13.1	0	11.6	7.8	7.3	0.574	103.8	7	3.6	0.142	4.2	5400	17.6	0.041598
##	25	13.0	0	11.6	6.3	5.7	0.641	98.4	14	2.6	0.070	2.1	4860	19.6	0.069197
##	26	13.1	0	12.1	16.0	14.3	0.631	107.1	3	7.7	0.102	4.1	6740	15.2	0.041698
##	27	13.5	0	10.9	6.9	7.1	0.540	96.5	6	0.4	0.080	2.2	5640	13.9	0.036099
##	28	15.2	0	11.2	8.2	7.6	0.571	101.8	10	7.9	0.103	2.8	5370	21.5	0.038201
##	29	11.9	0	10.7	16.6	15.7	0.521	93.8	168	8.9	0.092	3.6	6370	15.4	0.023400
##	30	16.6	1	8.9	5.8	5.4	0.521	97.3	46	25.4	0.072	2.6	3960	23.7	0.075298
##	31	14.0	0	9.3	5.5	5.4	0.535	104.5	6	2.0	0.135	4.0	4530	20.0	0.041999
##	32	12.5	0	10.9	9.0	8.1	0.586	96.4	97	8.2	0.105	4.3	6170	16.3	0.042698
##	33	14.7	1	10.4	6.3	6.4	0.560	97.2	23	9.5	0.076	2.4	4620	23.3	0.049499
##	34	12.6	0	11.8	9.7	9.7	0.542	99.0	18	2.1	0.102	3.5	5890	16.6	0.040799
##	35	12.3	0	10.2	9.7	8.7	0.526	94.8	113	7.6	0.124	5.0	5720	15.8	0.020700
##	36	15.0	0	10.0	10.9	9.8	0.531	96.4	9	2.4	0.087	3.8	5590	15.3	0.006900
##	37	17.7	1	8.7	5.8	5.6	0.638	97.4	24	34.9	0.076	2.8	3820	25.4	0.045198
##	38	13.3	0	10.4	5.1	4.7	0.599	102.4	7	4.0	0.099	2.7	4250	22.5	0.053998
##	39	14.9	1	8.8	6.1	5.4	0.515	95.3	36	16.5	0.086	3.5	3950	25.1	0.047099
##	40	14.5	1	10.4	8.2	7.4	0.560	98.1	96	12.6	0.088	3.1	4880	22.8	0.038801
##	41	14.8	0	12.2	7.2	6.6	0.601	99.8	9	1.9	0.084	2.0	5900	14.4	0.025100
##	42	14.1	0	10.9	5.6	5.4	0.523	96.8	4	0.2	0.107	3.7	4890	17.0	0.088904
##	43	16.2	1	9.9	7.5	7.0	0.522	99.6	40	20.8	0.073	2.7	4960	22.4	0.054902
##	44	13.6	0	12.1	9.5	9.6	0.574	101.2	29	3.6	0.111	3.7	6220	16.2	0.028100
##	45	13.9	1	8.8	4.6	4.1	0.480	96.8	19	4.9	0.135	5.3	4570	24.9	0.056202
##	46	12.6	0	10.4	10.6	9.7	0.599	98.9	40	2.4	0.078	2.5	5930	17.1	0.046598
##	47	13.0	0	12.1	9.0	9.1	0.623	104.9	3	2.2	0.113	4.0	5880	16.0	0.052802

##	Time	Crime
##	1	26.2011 791
##	2	25.2999 1635
##	3	24.3006 578
##	4	29.9012 1969
##	5	21.2998 1234
##	6	20.9995 682
##	7	20.6993 963
##	8	24.5988 1555
##	9	29.4001 856
##	10	19.5994 705
##	11	41.6000 1674
##	12	34.2984 849
##	13	36.2993 511
##	14	21.5010 664
##	15	22.7008 798

```
## 16 26.0991 946
## 17 19.1002 539
## 18 18.1996 929
## 19 24.9008 750
## 20 26.4010 1225
## 21 37.5998 742
## 22 37.0994 439
## 23 25.1989 1216
## 24 17.6000 968
## 25 21.9003 523
## 26 22.1005 1993
## 27 28.4999 342
## 28 25.8006 1216
## 29 36.7009 1043
## 30 28.3011 696
## 31 21.7998 373
## 32 30.9014 754
## 33 25.5005 1072
## 34 21.6997 923
## 35 37.4011 653
## 36 44.0004 1272
## 37 31.6995 831
## 38 16.6999 566
## 39 27.3004 826
## 40 29.3004 1151
## 41 30.0001 880
## 42 12.1996 542
## 43 31.9989 823
## 44 30.0001 1030
## 45 32.5996 455
## 46 16.6999 508
## 47 16.0997 849
```

```
linear_model_crime <- lm(Crime~., data=crime_data)
summary(linear_model_crime)
```

```
##
## Call:
## lm(formula = Crime ~ ., data = crime_data)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -395.74  -98.09   -6.69  112.99  512.67
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept) -5.984e+03  1.628e+03  -3.675 0.000893 ***
## M              8.783e+01  4.171e+01   2.106 0.043443 *
## So            -3.803e+00  1.488e+02  -0.026 0.979765
## Ed             1.883e+02  6.209e+01   3.033 0.004861 **
## Po1            1.928e+02  1.061e+02   1.817 0.078892 .
## Po2           -1.094e+02  1.175e+02  -0.931 0.358830
## LF            -6.638e+02  1.470e+03  -0.452 0.654654
## M.F             1.741e+01  2.035e+01   0.855 0.398995
```

```
## Pop          -7.330e-01  1.290e+00  -0.568  0.573845
## NW           4.204e+00  6.481e+00   0.649  0.521279
## U1          -5.827e+03  4.210e+03  -1.384  0.176238
## U2           1.678e+02  8.234e+01   2.038  0.050161 .
## Wealth       9.617e-02  1.037e-01   0.928  0.360754
## Ineq         7.067e+01  2.272e+01   3.111  0.003983 **
## Prob        -4.855e+03  2.272e+03  -2.137  0.040627 *
## Time        -3.479e+00  7.165e+00  -0.486  0.630708
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 209.1 on 31 degrees of freedom
## Multiple R-squared:  0.8031, Adjusted R-squared:  0.7078
## F-statistic: 8.429 on 15 and 31 DF,  p-value: 3.539e-07
```

Interpretation of the summary information for the linear regression model *I asked ChatGPT the following question to understand the information available in the summary function output for my linear regression model. “How to interpret summary(my linear regression model using lm() function in R)?”*

- **Coefficients:** This section displays the estimated coefficients for each predictor variable in the model. It includes columns for the estimated coefficient, standard error, t-value, and p-value. The coefficient represents the change in the response variable for a one-unit change in the predictor, holding all other predictors constant.
- **Residuals:** This section provides information about the residuals of the model, including the minimum, 1st quartile, median, mean, 3rd quartile, and maximum values. Residuals are the differences between the observed and predicted values of the response variable.
- **Coefficients’ significance:** The p-values associated with each coefficient test the null hypothesis that the corresponding coefficient is zero (i.e., the predictor has no effect on the response). Lower p-values suggest stronger evidence against the null hypothesis and indicate that the predictor is likely to be significant.
- **Multiple R-squared and adjusted R-squared:** These metrics quantify the goodness-of-fit of the model. Multiple R-squared represents the proportion of variance in the response variable explained by the predictors, while adjusted R-squared adjusts for the number of predictors in the model.
- **F-statistic:** This statistic tests the overall significance of the model by comparing the variance explained by the model to the variance not explained. A low p-value for the F-statistic indicates that the model is a significant improvement over a null model with no predictors.
- **Residual standard error:** This is an estimate of the standard deviation of the errors in predicting the response variable. It provides a measure of the model’s accuracy in predicting new observations.
- **Degrees of Freedom:** These values represent the degrees of freedom associated with the model, residual, and total.
- **Significance stars:** Some versions of the summary output use stars to indicate the level of significance of the coefficients, with more stars indicating higher significance.

```
# create a new data set to predict the observed crime rate in a city with the following data
M <- c(14.0)
So <- c(0)
Ed <- c(10.0)
Po1 <- c(12.0)
```



```

Po2 <- c(15.5)
LF <- c(0.640)
M.F <- c(94.0)
Pop <- c(150)
NW <- c(1.1)
U1 <- c(0.120)
U2 <- c(3.6)
Wealth <- c(3200)
Ineq <- c(20.1)
Prob <- c(0.04)
Time <- c(39.0)
new_data <- data.frame(
  M = M,
  So = So,
  Ed = Ed,
  Po1 = Po1,
  Po2 = Po2,
  LF = LF,
  M.F = M.F,
  Pop = Pop,
  NW = NW,
  U1 = U1,
  U2 = U2,
  Wealth = Wealth,
  Ineq = Ineq,
  Prob = Prob,
  Time = Time
)
new_data

```

```

##      M So Ed Po1 Po2 LF M.F Pop NW U1 U2 Wealth Ineq Prob Time
## 1 14  0 10  12 15.5 0.64 94 150 1.1 0.12 3.6  3200 20.1 0.04  39

```

```

predict(linear_model_crime, newdata=new_data)

```

Prediction for the given data

```

##           1
## 155.4349

```

Observation: There are only five predictors obtaining star(s) while there are 15 predictors. This means that the remaining predictors with high p-value are likely to have no effect on the response variable and this potentially indicates that the model is overfitted.